

A nonprofit corporation and independent licensee of the Blue Cross and Blue Shield Association

Hospital-based SNF Coding Tip Sheet: Top 25 codes and ICD-10 Chapter Overview

<u>Chapter 5 - Mental, Behavioral and Neurodevelopmental Disorders (F00-F99)</u>

Key Updates

- Classification improvements (different categories)
- Code expansions:
 - Most notably, Other Isolated or Specific Phobias
- Updates to medical terminology:
 - Bipolar I disorder, single manic episode will change to manic episode.
 - Undersocialized conduct disorders, aggressive will become Conduct disorder childhood-onset type.
- Nicotine dependence updated to identify specific tobacco products (cigarettes, chewing tobacco, and other tobacco).

Chapter 6 - Diseases of the Nervous System (G00-G99)

Key Updates

- ICD-10-CM devotes chapter 6 to Diseases of the Nervous System. The sense organs (eye/adnexa and ear/mastoid processes) have their own chapters in ICD-10-CM.
- Classification improvements (significant changes to sleep disorders)
- Code expansions (e.g. Alzheimer's, headaches)
- Updates to medical terminology (epilepsy, seizures)
- Codes for transient ischemic attack are now included in the nervous system chapter.
- Codes for migraine have been expanded to fifth and sixth characters to indicate if the migraine is intractable and to provide additional specificity within the code description.

Chapter 9 - Diseases of the Circulatory System (100-199)

Key Updates

- A new category has been added for coding a subsequent acute myocardial infarction (MI), which is an MI that occurs within 28 days of a previous acute MI.
- The time frame for assigning the acute MI code is 28 days in ICD-10-CM.
- Terminology used to describe several cardiovascular conditions has been revised to reflect more current medical practice.
- A major change is the classification of hypertension, which in ICD-9-CM was classified by type:
 - Benign, Malignant or Unspecified.
- That classification is not required in ICD-10-CM because there is only one code for hypertension in ICD-10; the code is I10 - Essential (Primary) Hypertension.

Hospital-based SNF Coding Tip Sheet: Top 25 codes and ICD-10 chapter overview, continued

Chapter 10 - Diseases of the Respiratory System (J00-J99)

Key Updates

- Modifications have been made to specific categories that bring the terminology up-to-date with current medical practice.
 - Emphysema now contains codes with panlobular and centrilobular in their description.
 - Asthma is now classified as mild intermittent, mild persistent, moderate persistent and severe persistent.
- Specificity increased for diseases like influenza, acute bronchitis.
- Coding notes updated to require the coder to include information about tobacco use/dependence, where applicable.
- Procedural complications that affect the respiratory system are now included in this chapter in ICD-10-CM.

<u>Chapter 13 - Diseases of the Musculoskeletal System and Connective Tissue (M00-M99)</u> Key Updates

- Most of the codes within this chapter have site and laterality designations. For some conditions where more than one bone, joint or muscle is usually involved, such as osteoarthritis, there is a "multiple sites" code available.
- Many musculoskeletal conditions are a result of previous injury or trauma to a site, or are recurrent conditions. Bone, joint or muscle conditions that are the result of a healed injury are usually found in Chapter 13.
 - ICD-10-CM identifies three different causes for pathological fractures: Neoplastic disease, Osteoporosis, and other specified disease.
 - ICD-10-CM introduces the 7th character that describes type of encounter, or the state
 of a fractures' healing and any sequela. Some codes in this chapter will have the 7th
 character applied.

Chapter 14 - Diseases of the Genitourinary System (N00-N99)

Key Updates

- Procedural complications affecting the genitourinary system are included in chapter 14.
- In some of the categories specificity is based on the gender of the patient.
- Laterality is used to identify conditions under N60 category, benign mammary dysplasia.

<u>Chapter 18 - Symptoms, Signs and Abnormal Clinical and Laboratory Findings (R00-R99)</u> Key Updates

- Repeated falls are coded to a symptom code in ICD-10-CM (was a V code in ICD-9-CM).
- There are codes to identify a patient's coma scale.
- Systemic inflammatory response syndrome (SIRS) due to noninfectious process is moved from the Injury and Poisoning ICD-9-CM chapter to this ICD-10-CM chapter.

<u>Chapter 19 - Injury, Poisoning, and Certain Other Consequences of External Causes (S00-T88)</u>

Key Updates

- Injuries are grouped by body part rather than category of injury as they were in ICD-9-CM.
- Laterality is specified in this chapter.
- ICD-10-CM introduces a 7th character requirement that describes the type of encounter.
 Most categories in this chapter use the 7th character requirement. Most categories in this chapter (see below for fractures) have three 7th character values of:
 - A Initial encounter
 - D Subsequent encounter

Freestanding SNF Coding Tip Sheet: Top 25 codes and ICD-10 chapter overview, continued

- S Sequela
- For traumatic fractures, there are additional 7th character requirements depending upon the type of fracture, and complication. Some of these character descriptions are based on the Gustilo open fracture classification.
- Codes for underdosing are new in ICD-10-CM. Underdosing refers to taking less of a
 medication than is prescribed by a provider or less than the manufacturer's instructions.
 Codes for underdosing should not be assigned as principal or first-listed codes.

<u>Chapter 21 - Factors Influencing Health Status and Contact with Health Services (Z00-Z99)</u>

Key Updates

- These codes are used in any healthcare setting. Z codes may be used as either a first-listed or principal diagnosis code in the inpatient setting, or secondary code, depending on the circumstances of the encounter. Certain Z codes may only be used as first-listed or principal diagnosis in certain situations. Refer to Official Coding Guidelines for details.
- Z codes are not procedure codes. A corresponding procedure code must accompany a Z code to describe any procedure performed.

(Source: 2015 Draft ICD-10-CM Official Guidelines for Coding and Reporting, cms.gov/ICD-10 website http://www.cms.gov/Medicare/Coding/ICD10/Downloads/2016-ICD-10-CM-Guidelines.pdf)

The top 25 codes for hospital-based SNF in ICD-9 to ICD-10 mappings are found in the chart below.

| Top 25 | ICD-9 Code | ICD-9 Description | ICD-10 Code | ICD-10 Description |
|--------|---------------|---|----------------|---|
| 1 | V57.89 | Care involving other specified rehabilitation procedure | Z15.89 | Encounter for other specified aftercare |
| 2 | 728.87 | Muscle weakness (generalized) | M62.81 | Muscle weakness (generalized) |
| 3 | 719.7 | Difficulty in walking | R26.2 | Difficulty in walking, not elsewhere classified |
| 4 | 486 | Pneumonia organism unspecified | J18.8 | Other pneumonia, unspecified organism |
| | | | J18.9 | Pneumonia, unspecified organism |
| 5 | 428.0 | Congestive heart failure, unspecified | 150.20 | Unspecified systolic (congestive) heart failure |
| | | | 150.21 | Acute systolic (congestive) heart failure |
| | | | 150.22 | Chronic systolic (congestive) heart failure |
| | | | 150.23 | Acute on chronic systolic (congestive) heart failure |
| | | | 150.30 | Unspecified diastolic (congestive) heart failure |
| | | | I50.31 | Acute diastolic (congestive) heart failure |
| | | | 150.32 | Chronic diastolic (congestive) heart failure |
| | | _ | 150.33 | Acute on chronic diastolic (congestive) heart failure |
| | | | 150.40 | Unspecified combined systolic (congestive) and diastolic (congestive) heart failure |

| | ICD-9 | | ICD-10 | |
|--------|--------|--|--|--|
| Top 25 | Code | ICD-9 Description | Code | ICD-10 Description |
| | | | I50.41 | Acute combined systolic (congestive) and diastolic (congestive) heart failure |
| | | | 130.41 | Chronic combined systolic (congestive) and |
| | | | 150.42 | diastolic (congestive) heart failure |
| | | | | Acute on chronic combined systolic |
| | | | 150.40 | (congestive) and diastolic (congestive) heart |
| | | | 150.43 | failure |
| | | | 150.9 | Heart failure, unspecified |
| 6 | 342.01 | Flaccid hemiplegia & hemiparesis affecting dominant side | G81.01 | Flaccid hemiplegia affecting right dominant side |
| | | | C94 02 | Flaccid hemiplegia affecting left dominant |
| | | Aftercare following joint | G81.02 | side Aftercare following joint replacement |
| 7 | V54.81 | replacement | Z47.1 | surgery |
| 8 | V43.65 | | 706 651 | |
| | | Knee joint replacement | Z96.651 | Presence of right artificial knee joint |
| | | | Z96.652 | Presence of left artificial knee joint |
| | | | Z96.653 | Presence of artificial knee joint, bilateral |
| | | | Z96.659 | Presence of unspecified artificial knee joint |
| 9 | 482.9 | Bacterial pneumonia, unspecified | J15.9 | Unspecified bacterial pneumonia |
| 10 | 496 | Chronic airway obstruction, not | 1440 | Chronic obstructive pulmonary disease, |
| | | elsewhere classified Care involving other physical | J44.9 | unspecified |
| 11 | V57.1 | therapy | Z51.89 | Encounter for other specified aftercare |
| 12 | 599.0 | Urinary tract infection, site not specified | N39.0 | Urinary tract infection, site not specified |
| 13 | 787.22 | Dysphagia, oropharangeal phase | R13.12 | Dysphagia, oropharangeal phase |
| 14 | 786.05 | Shortness of breath | R06.02 | Shortness of breath |
| 15 | 427.31 | Atrial fibrillation | 148.0 | Paroxysmal atrial fibrillation |
| | | | 148.2 | Chronic atrial fibrillation |
| | | | I48.91 | Unspecified atrial fibrillation |
| 16 | V54.17 | Aftercare for healing traumatic fracture of vertebrae | Code range: S12.000D- S129.XXD; S22.000D- S22.089G; S32.000D- S32.059G | Note: This code maps to 288 ICD-10 codes for various fracture types for the cervical, thoracic and lumbar vertebra |
| 17 | 332.0 | Paralysis agitans | G20 | Parkinson's disease |
| 18 | 438.22 | Late effects of cerebrovascular disease, Hemiplegia affecting nondominant side | 169.053 | Hemiplegia & hemiparesis following nontraumatic subarachnoid hemorrhage affecting right non-dominant side |
| | | | 169.054 | Hemiplegia & hemiparesis following nontraumatic subarachnoid hemorrhage affecting left non-dominant side |
| | | | 169.153 | Hemiplegia & hemiparesis following nontraumatic intracerebral hemorrhage affecting right non-dominant side |

| | ICD-9 | | ICD-10 | |
|--------|--------|---------------------------|----------|---|
| Top 25 | Code | ICD-9 Description | Code | ICD-10 Description |
| | | | | Hemiplegia & hemiparesis following |
| | | | 100 454 | nontraumatic intracerebral hemorrhage |
| | | | 169.154 | affecting left non-dominant side |
| | | | | Hemiplegia & hemiparesis following |
| | | | 100.050 | nontraumatic intracraniall hemorrhage |
| | | | 169.253 | affecting right non-dominant side |
| | | | | Hemiplegia & hemiparesis following |
| | | | 100.054 | nontraumatic intracraniall hemorrhage |
| | | | 169.254 | affecting left non-dominant side |
| | | | | Hemiplegia & hemiparesis following |
| | | | 169.353 | cerebral infarction affecting right non- dominant side |
| | | | 109.333 | Hemiplegia & hemiparesis following |
| | | | | cerebral infarction affecting left non- |
| | | | 169.354 | dominant side |
| | | | 109.554 | Hemiplegia & hemiparesis following other |
| | | | | cerebrovascular disease affecting right non- |
| | | | 169.853 | dominant side |
| | | | 1.00.000 | Hemiplegia & hemiparesis following other |
| | | | | cerebrovascular disease affecting left non- |
| | | | 169.854 | dominant side |
| | | | | Hemiplegia & hemiparesis following |
| | | | | unspecified cerebrovascular disease |
| | | | 169.553 | affecting right non-dominant side |
| | | | | Hemiplegia & hemiparesis following |
| | | | | unspecified cerebrovascular disease |
| | | | 169.554 | affecting left non-dominant side |
| | | Cerebral artery occlusion | | Cerebral infarction due to unspecified |
| 19 | 434.91 | unspecified w/ cerebral | | occlusion or stenosis of unspecified cerebral |
| | | infarction | 163.50 | artery |
| | | | | Cerebral infarction due to unspecified |
| | | | 100: | occlusion or stenosis of right middle |
| | | | 163.551 | cerebral artery |
| | | | | Cerebral infarction due to unspecified |
| | | | IGO E40 | occlusion or stenosis of left middle cerebral |
| | | | 163.512 | artery Cerebral infarction due to unspecified |
| | | | | occlusion or stenosis of unspecified middle |
| | | | l63.519 | cerebral artery |
| | | | 100.019 | Cerebral infarction due to unspecified |
| | | | | occlusion or stenosis of right anterior |
| | | | 163.521 | cerebral artery |
| | | | | Cerebral infarction due to unspecified |
| | | | | occlusion or stenosis of left anterior cerebral |
| | | | 163.522 | artery |
| | | | | Cerebral infarction due to unspecified |
| | | | | occlusion or stenosis of unspecified anterior |
| | | | 163.529 | cerebral artery |
| | | | | Cerebral infarction due to unspecified |
| | | | | occlusion or stenosis of right posterior |
| | | | I63.531 | cerebral artery |
| | | | | Cerebral infarction due to unspecified |
| | | | | occlusion or stenosis of left posterior |
| | | | 163.532 | cerebral artery |
| | | | | Cerebral infarction due to unspecified |
| | | | 100 705 | occlusion or stenosis of unspecified |
| | | | 163.539 | posterior cerebral artery |

| | ICD-9 | | ICD-10 | |
|--------|--------|---|--|--|
| Top 25 | Code | ICD-9 Description | Code | ICD-10 Description |
| | | | | Cerebral infarction due to unspecified |
| | | | | occlusion or stenosis of right cerebellar |
| | | | I63.541 | artery |
| | | | | Cerebral infarction due to unspecified |
| | | | 163.542 | occlusion or stenosis of left cerebellar artery |
| | | | | Cerebral infarction due to unspecified |
| | | | | occlusion or stenosis of unspecified |
| | | | 163.549 | cerebellar artery |
| | | | | Cerebral infarction due to unspecified |
| | | | | occlusion or stenosis of other cerebral |
| | | | 163.59 | artery |
| | | | 163.8 | Other cerebral infarction |
| | | | 163.9 | Cerebral infarction, unspecified |
| 20 | V54.13 | Aftercare for healing traumatic fracture of hip | Code range: \$32.301D- \$32.669G; \$72.001D- \$72.146J | Note: This code maps to 552 ICD-10 codes for various types of traumatic fractures of the hip |
| 21 | 799.3 | Debility unspecified | R53.81 | Other malaise |
| 22 | 331.0 | Alzheimer's disease | G30.0 | Alzheimer's disease w/ early onset |
| | | | | |
| | | | G30.1 | Alzheimer's disease w/ late onset |
| | | | G30.8 | Other Alzheimer's disease |
| | | | G30.9 | Alzheimer's disease unspecified |
| 23 | 996.63 | Infection & inflammatory reaction due to nervous system device implant & graft | T85.79XA | Infection & inflammatory reaction due to other internal prosthetic device implants & grafts, initial encounter |
| 24 | 715.96 | Osteoarthrosis unspecified whether generalized or localized involving lower leg | M17.9 | Osteoarthritis of knee, unspecified |
| 25 | 436 | Acute, but ill-defined cerebrovascular disease | 167.89 | Other cerebrovascular disease |

The codes highlighted in **orange** indicate the individual ICD-9 code that is being mapped to one or many ICD-10 codes

(Source of ICD-9-CM to ICD-10-CM mappings: CMS.org General Equivalence Mappings (GEMs), 2015)

The information in this document is not intended to impart legal advice. This overview is intended as an educational tool only and should not be relied upon as legal or compliance advice. If you have any legal questions about the information contained herein, you should consult your attorney or other professional legal services provider.