



Nonprofit corporations and independent licensees
of the Blue Cross and Blue Shield Association

Medical benefit drug policies are a source for BCBSM and BCN medical policy information only. These documents are not to be used to determine benefits or reimbursement. Please reference the appropriate certificate or contract for benefit information. This policy may be updated and therefore subject to change.

Effective Date: 10/03/2024

Simponi Aria® (golimumab)

HCPCS: J1602

Policy:

Requests must be supported by submission of chart notes and patient specific documentation.

- A. Coverage of the requested drug is provided when all the following are met:
 - a. FDA approved age
 - b. Diagnosis of rheumatoid arthritis (RA):
 - i. Trial and failure of at least a 3-month trial of one disease-modifying anti-rheumatic drug (DMARD) unless contraindicated or not tolerated. Examples include: methotrexate, hydroxychloroquine, leflunomide, sulfasalazine
 - ii. Used in combination with methotrexate (unless contraindicated)
 - c. Diagnosis of psoriatic arthritis (PsA)
 - d. Diagnosis of ankylosing spondylitis (AS)
 - e. Diagnosis of polyarticular juvenile idiopathic arthritis (pJIA)
 - i. Trial and failure of at least a 3-month trial of one DMARD unless contraindicated or not tolerated. Examples include methotrexate and leflunomide
 - f. Not to be used in combination with other biologics or targeted DMARDs for the same indication
 - g. Trial and failure, contraindication, OR intolerance to the preferred drugs as listed in BCBSM/BCN's utilization management medical drug list
- B. Quantity Limitations, Authorization Period and Renewal Criteria
 - a. Quantity Limit: Align with FDA recommended dosing
 - b. Initial Authorization Period: One year at a time
 - c. Renewal Criteria: Clinical documentation must be provided to confirm that current criteria are met and that the medication is providing clinical benefit

***Note: Coverage and approval duration may differ for Medicare Part B members based on any applicable criteria outlined in Local Coverage Determinations (LCD) or National Coverage Determinations (NCD) as determined by Center for Medicare and Medicaid Services (CMS). See the CMS website at <http://www.cms.hhs.gov/>. Determination of coverage of Part B drugs is based on medically accepted indications which have supported citations included or approved for inclusion determined by CMS approved compendia.

This policy and any information contained herein is the property of Blue Cross Blue Shield of Michigan and its subsidiaries, is strictly confidential, and its use is intended for the P&T committee, its members and BCBSM employees for the purpose of coverage determinations.

Background Information:

- Simponi Aria (golimumab) is a tumor necrosis factor inhibitor (TNFi) indicated for the treatment of adult patients with moderately to severely active rheumatoid arthritis (RA; in combination with methotrexate) or active ankylosing spondylitis, and for pediatric patients 2 years of age and older with active psoriatic arthritis or active polyarticular juvenile idiopathic arthritis.
- The use of Simponi Aria in combination with other biologic agents or targeted disease-modifying antirheumatic drugs is not recommended due to an increased risk of infection and a lack of robust safety and efficacy data to support combination use.
- Ankylosing Spondylitis
 - Axial spondyloarthritis, comprising ankylosing spondylitis (AS) and non-radiographic axial spondyloarthritis (NRAS), is the main form of chronic inflammatory arthritis affecting the axial skeleton. Non-radiographic means that damage to the joints is not visible on X-ray. When changes to the vertebrae (the bones of the spine) or sacroiliac joints don't show any changes on an X-ray, that's known as NRAS. Once the joints are clearly affected on an X-ray, a person can be diagnosed with AS.
 - The 2019 American College of Rheumatology recommendations for AS and NRAS are similar. In adult patients who have active disease despite treatment with nonsteroidal anti-inflammatory drugs (NSAIDs), treatment with tumor necrosis factor inhibitors (TNFi) are recommended. They do not recommend any particular TNFi as the preferred choice for the typical patient.
 - Cosentyx® (secukinumab) or Taltz® (ixekizumab) is recommended over the use of a second TNFi in patients with primary nonresponse to the first TNFi, whereas for patients with a secondary nonresponse (i.e. those who relapse after an initial response) it may be beneficial to switch to a different TNFi rather than immediately switch to a different biologic class.
- Rheumatoid Arthritis
 - The 2021 American College of Rheumatology (ACR) Guidelines for the Treatment of Rheumatoid Arthritis (RA) established recommendations for the care of adult RA patients. The guidelines state that treatment decisions should follow a shared decision-making process and should be reevaluated within a minimum of 3 months based on the efficacy and tolerability of the DMARD(s) chosen.
 - For the initial treatment of symptomatic RA, the guidelines strongly recommend the use of conventional synthetic DMARD (csDMARD) monotherapy in those who are DMARD-naive. csDMARD monotherapy is a less costly first line treatment option with an extensive safety record accompanied by well-documented clinical efficacy and a large body of clinical experience and familiarity among prescribers. csDMARDs in the guidelines refer to methotrexate (MTX), hydroxychloroquine, leflunomide (LEF), and sulfasalazine. Azathioprine, cyclosporine, minocycline, and gold were not included due to their infrequent use in RA and lack of new data since the prior guidelines were published. Oral MTX is recommended as the preferred initial DMARD for patients with moderate-to-high disease activity, and hydroxychloroquine is recommended as the preferred initial DMARD for patients with low disease activity.
 - If disease activity remains moderate or high despite optimal dosing of methotrexate monotherapy, the use of dual therapy with methotrexate plus biologic DMARD (bDMARD; etanercept, adalimumab, infliximab, golimumab, certolizumab pegol, abatacept, tocilizumab, sarilumab, rituximab) or targeted synthetic DMARD (tsDMARD; tofacitinib, baricitinib, upadacitinib) therapy is conditionally recommended over the use of triple therapy (i.e., addition of sulfasalazine and hydroxychloroquine). The guidelines do not inform preference of

This policy and any information contained herein is the property of Blue Cross Blue Shield of Michigan and its subsidiaries, is strictly confidential, and its use is intended for the P&T committee, its members and BCBSM employees for the purpose of coverage determinations.

bDMARD over tsDMARD therapy, or vice-versa, for use in combination with methotrexate. No one agent has been shown to be superior to another. The guidelines do acknowledge the emergence of safety signals for the JAK inhibitor class (tsDMARD), and state that further modification of this recommendation may be necessary as additional data are published.

- A treat-to-target approach is conditionally recommended over usual care for patients who have had an inadequate response to bDMARDs or tsDMARDs. Treat-to-target refers to a systematic approach involving frequent monitoring of disease activity using validated instruments and modification of treatment to minimize disease activity with the goal of reaching a pre-defined target (low disease activity or remission).

- Psoriatic Arthritis

- PsA is a chronic inflammatory disease often associated with psoriasis. Psoriasis is an autoimmune disease affecting the skin, resulting in scaly red and white patches. These patches, called plaques, may appear anywhere on the body. The inflammation may also develop in the joints, which is classified as PsA. PsA occurs in up to 30% of patients with psoriasis, most commonly appearing between the ages of 30 and 50. PsA causes pain, stiffness, and swelling in and around the joints. If not properly treated, progressive joint damage may occur.
- Per the 2018 American College of Rheumatology (ACR)/National Psoriasis Foundation (NPF) guideline for the treatment of psoriatic arthritis, all recommendations for treatment-naïve patients with active PsA are conditional based on low- to very-low quality evidence.
- In treatment-naïve patients, oral systemic medications (OSMs), such as methotrexate, sulfasalazine, cyclosporine, and leflunomide, may be used in patients without severe psoriatic arthritis and without severe psoriasis. OSMs have robust longitudinal safety and efficacy data in patients with PsA. Maximal response to OSMs are most commonly achieved within 3 months of therapy.
- If PsA remains active despite OSM therapy, switching to a TNFi, an IL-17i, or an IL-12/23i biologic is recommended over switching to a different OSM; switching to a TNFi biologic over an IL-17i or IL-12/23i biologic is conditionally recommended in this scenario based on moderate quality evidence. The detailed recommendations for subsequent therapies can be found in the 2018 ACR/NPF guideline for the treatment of psoriatic arthritis.

- Juvenile Idiopathic Arthritis

- Juvenile idiopathic arthritis (JIA) defines a collection of inflammatory arthritides of unknown etiology. Onset is prior to 16 years of age with a minimum duration of 6 weeks and other potential causes of synovitis are excluded. JIA can be subdivided into polyarticular JIA and systemic JIA.
- pJIA is defined by the presence of more than 4 affected joints in the first 6 months of illness and comprises 20-30% of children with JIA. Therapy is directed toward treating the underlying inflammation and preventing JIA-associated complications and adverse effects of its treatment.
- The 2019 American College of Rheumatology/Arthritis Foundation (ACR/AF) guideline for the treatment of JIA strongly recommends initial therapy for pJIA with a disease-modifying anti-rheumatic agent (DMARD) such as methotrexate (MTX) or leflunomide. MTX is conditionally recommended over leflunomide as it has a greater volume of data supporting its effectiveness compared to leflunomide and can be administered subcutaneously (recommended) or orally. The guidelines consider an adequate trial of a DMARD to be 3 months.

- If moderate or high disease activity persists despite DMARD use, the ACR/AF guidelines recommend biologic agents either in combination with a DMARD or as monotherapy in certain situations. Biologic agents FDA approved for pJIA in patients 2 years of age and older include Humira® (adalimumab), Enbrel® (etanercept), Actemra® (tocilizumab), Orencia® (abatacept), Simponi Aria® (golimumab), Xeljanz® (tofacitinib), and Xeljanz (tofacitinib) oral solution.
 - Of note, biologic therapy may be an appropriate initial therapy option in certain pJIA patients with risk factors and involvement of high-risk joints, high disease activity, and/or for those judged to be at high risk of disabling joint damage
- There is the most experience with TNFi (Humira, Enbrel, Simponi Aria) as initial biologic therapy; however, the preferred class of initial biologic is not specified in the guideline recommendations due to a lack of comparative data and the consideration that non-TNFi biologics may be preferred in certain patient-specific scenarios.
- If a TNFi is started as the initial biologic, switching to a non-TNFi (tocilizumab or abatacept) is recommended over switching to a second TNFi. An exception to this is for those who had a good initial response to the first TNFi.

References:

1. Singh JA, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Rheumatol.* 2019 Jan; 71 (1): 5-32.
2. Fraenkel L, et al. 2021 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Care & Research.* 2021 Jul; 73 (7):924-939.
3. Ward, MM, Deodhar, A, Akl, EA, et al. 2019 Update of the American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network Recommendations for the Treatment of Ankylosing Spondylitis and Nonradiographic Axial Spondyloarthritis. *Arthritis Rheumatol.* 2019 Oct;71(10):1599-1613.
4. Ringold et al. 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Treatment of JIA. *Arthritis Care and Research.* Vol 71 No 6 Jun 2019.
5. Cimzia (certrolizumab pegol) [prescribing information]. Smyrna, GA: UCB, Inc. September 2019.
6. Enbrel (etanercept) [prescribing information]. Thousand Oaks, CA: Immunex Corporation; March 2020.
7. Humira (adalimumab) [prescribing information]. North Chicago, IL: AbbVie; March 2020.
8. Inflectra (infliximab-dyyb) [prescribing information]. New York, NY: Pfizer; June 2019.
9. Remicade (infliximab) [prescribing information]. Horsham, PA: Janssen Biotech Inc; May 2020.
10. Renflexis (infliximab-abda) [prescribing information]. Whitehouse Station, NJ: Merc Sharp & Dohme Corp; February 2020.
11. Simponi (golimumab) [prescribing information]. Horsham, PA: Janssen Biotech Inc; September 2019.
12. Simponi Aria (golimumab) [prescribing information]. Horsham, PA: Janssen Biotech Inc; September 2019.

Policy History												
#	Date	Change Description										
1.8	Effective Date: 10/03/2024	Added "for the same indication" to the not to be used in combination with other biologics or targeted DMARDs criteria										
1.7	Effective Date: 12/14/2023	Annual review of criteria performed, no changes were made										
1.6	Effective Date: 12/01/2022	Annual review of criteria performed, no changes were made										
1.5	Effective Date: 12/09/2021	Removed FDA approved indications criteria Removed oral DMARD criteria for psoriatic arthritis indication Updated background information for rheumatoid arthritis to reflect 2021 guidelines; remaining guideline references within this section of the policy are up to date										
1.4	Effective Date: 12/03/2020	Criteria updated to align management between pharmacy and medical benefit for all listed indications										
1.3	Effective Date: 02/06/2020	Annual Review										
1.2	Effective Date: 02/14/2019	Updated criteria for PsA diagnosis Updated language regarding combination therapy with other biologics										
1.1	Effective Date: 09/07/2018	The Treatments for RA Policy is being retired; Adding RA criteria back into this document										
1.0	Effective Date: 05/03/2018	<p>New Criteria</p> <table border="1"> <thead> <tr> <th>Line of Business</th> <th>PA Required in Medical Management System (Yes/No)</th> </tr> </thead> <tbody> <tr> <td>BCBS</td> <td>Yes</td> </tr> <tr> <td>BCN</td> <td>Yes</td> </tr> <tr> <td>MAPPO</td> <td>Yes</td> </tr> <tr> <td>BCNA</td> <td>Yes</td> </tr> </tbody> </table>	Line of Business	PA Required in Medical Management System (Yes/No)	BCBS	Yes	BCN	Yes	MAPPO	Yes	BCNA	Yes
Line of Business	PA Required in Medical Management System (Yes/No)											
BCBS	Yes											
BCN	Yes											
MAPPO	Yes											
BCNA	Yes											

* The prescribing information for a drug is subject to change. To ensure you are reading the most current information it is advised that you reference the most updated prescribing information by visiting the drug or manufacturer website or <http://dailymed.nlm.nih.gov/dailymed/index.cfm>.

**Blue Cross Blue Shield/Blue Care Network of Michigan
Medication Authorization Request Form
SIMPONIA[®] (golimumab) J1602**



This form is to be used by participating physicians to obtain coverage for SIMPONIA[®]. For commercial members only, please complete this form and submit via fax to 1-877-325-5979. If you have any questions regarding this process, please contact BCBSM Provider Relations and Servicing or the Medical Drug Helpdesk at 1-800-437-3803 for assistance.

PATIENT INFORMATION	PHYSICIAN INFORMATION
Name	Name
ID Number	Specialty
D.O.B. <input type="checkbox"/> Male <input type="checkbox"/> Female	Address
Pt weight (in kg) Date recorded: _____	City /State/Zip
Diagnosis	Phone/Fax: P: () - F: () -
Drug Name	NPI
Dose and Quantity	Contact Person
Directions	Contact Person Phone / Ext.
Date of Service(s)	

STEP 1: DISEASE STATE INFORMATION

1. Is this request for: Initiation Continuation *Date patient started therapy:* _____
2. Site of administration? Provider office/Home infusion Other: _____
 Hospital outpatient facility (go to #3) *Reason for Hospital Outpatient administration:* _____
3. **Please specify location of administration if hospital outpatient infusion:** _____
4. **Please provide the NPI number for the place of administration:** _____

5. Initiation AND Continuation of therapy:

- a. Please check patient's diagnosis:
 - Rheumatoid arthritis (RA)
 - Polyarticular juvenile idiopathic arthritis
 - Psoriatic arthritis
 - Ankylosing spondylitis
 - Other, list diagnosis: _____
- b. Has the patient tried and failed therapy with at least one conventional therapy?
 - Methotrexate, Date started: _____ Date ended: _____
 - Sulfasalazine, Date started: _____ Date ended: _____
 - Hydroxychloroquine, Date started: _____ Date ended: _____
 - Leflunomide, Date started: _____ Date ended: _____
 - Cyclosporine, Date started: _____ Date ended: _____
 - Other: _____, Date started: _____ Date ended: _____
- c. **For RA indication:** Will the patient use Methotrexate along with Simponi Aria? yes no, Provide rationale: _____
- d. Will the patient be receiving Simponi Aria with other biologic agents (for example: Remicade, Humira, Stelara, Cosentyx, Entyvio, or Tremfya, etc.) or targeted DMARD medications (for example: Otezla)?
 Yes, Comment: _____ No

6. Continuation Request: Simponi Aria start date: _____

- a. Have the patient's signs and symptoms improved with Simponi Aria?
 Yes No, Comment: _____

Please add any other supporting medical information necessary for our review

Coverage will not be provided if the prescribing physician's signature and date are not reflected on this document.

Request for expedited review: I certify that applying the standard review time frame may seriously jeopardize the life or health of the member or the member's ability to regain maximum function

Physician's Name	Physician Signature	Date
Step 2: Checklist	<input type="checkbox"/> Form Completely Filled Out <input type="checkbox"/> Attached Chart Notes	<input type="checkbox"/> Prior Treatments with traditional DMARD
Step 3: Submit	By Fax: BCBSM Specialty Pharmacy Mailbox 1-877-325-5979	By Mail: BCBSM Specialty Pharmacy Program P.O. Box 312320, Detroit, MI 48231-2320

Confidentiality notice: This transmission contains confidential information belonging to the sender that is legally privileged. This information is intended only for use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution or action taken in reliance on the contents of this document is strictly prohibited. If you have received this in error, please notify the sender to arrange for the return of this document.