

---

## Medical Policy



Nonprofit corporations and independent licensees  
of the Blue Cross and Blue Shield Association

**Joint Medical Policies are a source for BCBSM and BCN medical policy information only. These documents are not to be used to determine benefits or reimbursement. Please reference the appropriate certificate or contract for benefit information. This policy may be updated and is therefore subject to change.**

---

**\*Current Policy Effective Date: 5/1/21**  
(See policy history boxes for previous effective dates)

### **Title: Prostatic Urethral Lift Procedure for the Treatment of BPH**

---

#### **Description/Background**

##### **BENIGN PROSTATIC HYPERPLASIA (BPH)**

BPH is a common disorder among older individuals that results from hyperplastic nodules in the periurethral or transitional zone of the prostate. The clinical manifestations of BPH include increased urinary frequency, nocturia, urgency or hesitancy to urinate, and a weak stream when urinating. The urinary tract symptoms often progress with worsening hypertrophy and may lead to acute urinary retention, incontinence, renal insufficiency, and/or urinary tract infection.

Two scores are widely used to evaluate BPH-related symptoms: the American Urological Association Symptom Index (AUASI) and the International Prostate Symptom Score (IPSS). The AUASI is a self-administered 7-item questionnaire assessing the severity of various urinary symptoms.<sup>2</sup> Total AUASI scores range from 0 to 35, with overall severity categorized as mild ( $\leq 7$ ), moderate (8-19), or severe (20-35).<sup>1</sup> The IPSS incorporates questions from the AUASI and a quality of life question or a "Bother score."<sup>3</sup>

##### **Management**

Evaluation and management of BPH include assessment for other causes of lower urinary tract dysfunction (e.g., prostate cancer), symptom severity, and the degree that symptoms are bothersome to determine the therapeutic approach.

##### **Medical Therapy**

A discussion about medical therapy is generally indicated for patients with moderate-to-severe symptoms (e.g., an AUASI score of  $\geq 8$ ), bothersome symptoms, or both. Available medical therapies for BPH-related lower urinary tract dysfunction include  $\alpha$ -adrenergic blockers (e.g., alfuzosin, doxazosin, tamsulosin, terazosin, silodosin), 5 $\alpha$ -reductase inhibitors (e.g., finasteride, dutasteride), combination  $\alpha$ -adrenergic blockers and 5 $\alpha$ -reductase inhibitors, anti-muscarinic agents (e.g., darifenacin, solifenacin, oxybutynin), and phosphodiesterase-5 inhibitors (e.g., tadalafil).<sup>1</sup> In a meta-analysis of both indirect comparisons from placebo-

controlled studies (including 6333 patients) and direct comparative studies (including 507 patients), Djavan et al (1999) found that the IPSS improved by 30% to 40% and the Qmax score (mean peak urinary flow rate) improved by 16% to 25% in individuals assigned to  $\alpha$ -adrenergic blockers.<sup>4</sup> Combination therapy using an  $\alpha$ -adrenergic blocker and 5 $\alpha$ -reductase inhibitor has been shown to be more effective for improving IPSS than either treatment alone, with median scores improving by more than 40% over 1 year and by more than 45% over 4 years.<sup>5</sup>

### **Surgical and Ablative Therapies**

Patients who do not have sufficient response to medical therapy, or who are experiencing significant side effects with medical therapy, may be referred for surgical or ablative therapies. Various surgical and ablative procedures are used to treat BPH. Transurethral resection of the prostate is generally considered the reference standard for comparisons of BPH procedures.<sup>6</sup> In the perioperative period, transurethral resection of the prostate is associated with risks of any operative procedure (e.g., anesthesia risks, blood loss). Although short-term mortality risks are generally low, a large prospective study with 10654 patients by Reich et al (2008) reported the following short-term complications: “failure to void (5.8%), surgical revision (5.6%), significant urinary tract infection (3.6%), bleeding requiring transfusions (2.9%), and transurethral resection syndrome (1.4%).”<sup>7</sup> Incidental carcinoma of the prostate was diagnosed by histologic examination in 9.8% of patients. In the longer term, transurethral resection of the prostate is associated with increased risk of sexual dysfunction and incontinence.

Several minimally invasive prostate ablation procedures are available, including transurethral microwave thermotherapy, transurethral needle ablation of the prostate, urethromicroablation phototherapy, and photo selective vaporization of the prostate. The minimally invasive procedures were individually compared with transurethral resection of the prostate at the time they were developed, which provided a general benchmark for evaluating those procedures.

### **UroLift® System**

Prostatic Urethral Lift using the UroLift® System was designed to meet the need for a less invasive option. The UroLift System is a straightforward treatment that provides immediate, visible results. The procedure can be done in an outpatient or inpatient setting and under general or local anesthesia. Patients typically can return home the same day without a catheter, and experience rapid symptom relief and recovery with low complication rates. This transurethral BPH treatment does not require ongoing medication, heating, cutting or removal of the prostate tissue.

---

## **Regulatory Status**

The UroLift System received FDA authorization for marketing through a de novo classification approval on September 13, 2013 (K130651), as a Class II device. It is indicated for the treatment of symptoms due to urinary outflow obstruction secondary to benign prostatic hyperplasia (BPH) in men age 45 and above. Per the FDA approval, the UroLift System should *not* be used if the patient has:

- Prostate volume of >100 cc;
- A urinary tract infection;
- Urethra conditions that may prevent insertion of delivery system into bladder;
- Urinary incontinence;

- Current gross hematuria; or
- A known allergy to nickel.

In addition, the UroLift Implant has been shown to be MR Conditional and can be scanned under the following conditions:

- Static magnetic field strength of 3 Tesla or less;
- Maximum spatial gradient magnetic field of 720 Gauss/cm;
- A maximum whole-body-averaged specific absorption rate (SAR) of 4 W/kg for 15 minutes of scanning.

In March 2016, FDA determined that the UL500 was substantially equivalent to existing devices (UL400) for the treatment of symptoms of urinary flow obstruction secondary to benign prostatic hyperplasia in men age 45 years and older. In 2017, the FDA expanded the indication for the UL400 and UL500 to include lateral and median lobe hyperplasia in men 45 years or older. An additional clearance in 2019 (K193269) modified one contraindication from men with prostate volume of >80 cc to men with prostate volume of >100 cc. FDA product code: PEW.

---

## Medical Policy Statement

The safety and efficacy of the prostatic urethral lift procedure for the treatment of benign prostatic hypertrophy (BPH) have been established. It is a useful therapeutic option for men with symptomatic BPH who have failed conventional pharmacologic therapy.

---

## Inclusionary and Exclusionary Guidelines (Clinically based guidelines that may support individual consideration and pre-authorization decisions)

### Inclusions:

Candidates for the prostatic urethral lift procedure must meet **all** of the following guidelines:

- Age 45 years or older
- A documented diagnosis of symptomatic benign prostatic hypertrophy (BPH) of the *lateral* lobes of the prostate, including but not limited to the following symptoms:
  - Difficulty starting and stopping urination (hesitancy and straining).
  - Decreased strength of the urine stream (weak flow).
  - Dribbling after urination.
  - Feeling that the bladder is not completely empty.
  - An urge to urinate again soon after urinating (urgency).
  - Pain during urination (dysuria).
  - Nocturia – waking up several times during the night with the urge to urinate.
  - Frequent urinary tract infections secondary to urinary obstruction.
- Documented failure of, inability to tolerate, or undesirable side effects of pharmacologic intervention for BPH, including, but not limited to
  - Alpha blockers such as Uroxatral, Cardura, Rapaflo, Flomax or Hytrin
  - 5-Alpha Reductase Inhibitors for BPH, such as Avodart or Proscar
  - Combination drugs using both an alpha blocker and a 5-alpha reductase inhibitor.

### Exclusions:

Patients not meeting the patient selection criteria above.

---

**BCBSA/HCPCS Level II Codes** *(Note: The inclusion of a code in this list is not a guarantee of coverage. Please refer to the medical policy statement to determine the status of a given procedure.)*

**Established codes:**

52441            52442

**Other codes (investigational, not medically necessary, etc.):**

N/A

---

**Rationale**

Evidence reviews assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life, and ability to function—including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, two domains are examined: the relevance and the quality and credibility. To be relevant, studies must represent one or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

**PROSTATIC URETHRAL LIFT**

**Clinical Context and Therapy Purpose**

The purpose of PUL in patients who have lower urinary tract symptoms due to benign prostatic hyperplasia (BPH) is to provide a treatment option that is an alternative to or an improvement on existing therapies such as medical management or transurethral resection of the prostate (TURP).

Evaluation and management of BPH include assessment for other causes of lower urinary tract dysfunction (e.g., prostate cancer), symptom severity, and the degree that symptoms are bothersome to determine the therapeutic approach.

For patients with moderate-to-severe symptoms (e.g., an AUASI score of  $\geq 8$ ), bothersome symptoms, or both, a discussion about medical therapy is reasonable. BPH should generally be treated medically first. Available medical therapies for BPH-related lower urinary tract

dysfunction include  $\alpha$ -adrenergic blockers (e.g., alfuzosin, doxazosin, tamsulosin, terazosin, silodosin), 5 $\alpha$ -reductase inhibitors (e.g., finasteride, dutasteride), combination  $\alpha$ -adrenergic blockers and 5 $\alpha$ -reductase inhibitors, anti-muscarinic agents (e.g., darifenacin, solifenacin, oxybutynin), and phosphodiesterase-5 inhibitors (e.g., tadalafil).<sup>1</sup> In a meta-analysis of both indirect comparisons from placebo-controlled studies (including 6333 patients) and direct comparative studies (including 507 patients), Djavan et al (1999) found that the IPSS improved by 30% to 40% and the Qmax score (mean peak urinary flow rate) improved by 16% to 25% in individuals assigned to  $\alpha$ adrenergic blockers.<sup>4</sup> Combination therapy using an  $\alpha$ -adrenergic blocker and 5 $\alpha$ -reductase inhibitor has been shown to be more effective for improving IPSS than either treatment alone, with median scores improving by more than 40% over 1 year and by more than 45% over 4 years.

Patients who do not have sufficient response to medical therapy, or who are experiencing significant side effects with medical therapy, may be referred for surgical or ablative therapies. Various surgical and ablative procedures are used to treat BPH. Transurethral resection of the prostate is generally considered the reference standard for comparisons of BPH procedures.<sup>5</sup> In the perioperative period, transurethral resection of the prostate is associated with risks of any operative procedure (e.g., anesthesia risks, blood loss). Although short-term mortality risks are generally low, a large prospective study with 10654 patients by Reich et al (2008) reported the following short-term complications: "failure to void (5.8%), surgical revision (5.6%), significant urinary tract infection (3.6%), bleeding requiring transfusions (2.9%), and transurethral resection syndrome (1.4%)."<sup>6</sup> Incidental carcinoma of the prostate was diagnosed by histologic examination in 9.8% of patients. In the longer term, transurethral resection of the prostate is associated with increased risk of sexual dysfunction and incontinence.

Several minimally invasive prostate ablation procedures are available, including transurethral microwave thermotherapy, transurethral needle ablation of the prostate, urethromicroablation phototherapy, and photo selective vaporization of the prostate. The minimally invasive procedures were individually compared with transurethral resection of the prostate at the time they were developed, which provided a general benchmark for evaluating those procedures.

The question addressed in this evidence review is: Does PUL improve the net health outcome in individuals with BPH?

The following **PICOs** were used to select literature to inform this review.

### **Patients**

BPH prevalence increases with age and is present in more than 80% of individuals ages 70 to 79.<sup>1</sup>

The relevant population-of-interest are men who are experiencing lower urinary tract symptoms without a history suggesting non-BPH causes of the symptoms and who do not have sufficient response to medical therapy or are experiencing significant side effects with medical therapy.

AUA recommends surgical intervention for patients who have "renal insufficiency secondary to BPH, refractory urinary retention secondary to BPH, recurrent urinary tract infections (UTIs), recurrent bladder stones or gross hematuria due to BPH, and/or with LUTS attributed to BPH refractory to and/or unwilling to use other therapies."<sup>7</sup>

## Interventions

The therapy being considered is PUL. The PUL procedure involves placement of one or more implants in the lateral lobes of the prostate using a transurethral delivery device. The implant device is designed to retract the prostate to allow expansion of the prostatic urethra. The implants are retained in the prostate to maintain an expanded urethral lumen.

One device, the NeoTract UroLift System, has been cleared for marketing by the U.S. Food and Drug Administration (see Regulatory Status section). The device has two main components: the delivery device and the implant. Each delivery device comes preloaded with a UroLift implant.

## Comparators

Various surgical or ablative procedures are used to treat BPH. TURP is generally considered the reference standard for comparisons of BPH procedures. Several minimally invasive prostate ablation procedures have also been developed, including transurethral microwave thermotherapy, transurethral needle ablation of the prostate, urethromicroablation phototherapy, and photo-selective vaporization of the prostate.

## Outcomes

A number of health status measures are used to evaluate symptoms relevant to BPH and adverse events of treatment for BPH, including urinary dysfunction measured by urinary flow rate (Qmax), ejaculatory dysfunction, overall sexual health, and overall quality of life. Some validated patient-reported scales are shown in Table 1.

Of note, prostate volume does not have a direct correlation with severity of urinary symptoms.<sup>8</sup>

**Table 1. Patient-Reported Health Outcome Measures Relevant to Benign Prostatic Hyperplasia**

| Measure   | Outcome Evaluated                            | Description   | Clinically Meaningful Difference (if Known)   |
|---|--|---|---|
| Male Sexual Health Questionnaire for Ejaculatory Dysfunction <sup>9</sup>                             | Ejaculatory function and quality of life     | Patient-administered, 4-item scale. Symptoms rated as absent (15) to severe (0). QOL assessed as no problem (0) to extremely bothered (5).  | NR  |
| Sexual Health Inventory for Men <sup>10</sup>   | Erectile function                            | Patient-administered, 5-item scale. Erectile dysfunction rated as severe (1-7), moderate (8-11), mild to moderate (12-16), or mild (17-21). Fewest symptoms present for patients with score 22-25.  | 5-point change <sup>11</sup>  |
| American Urological Association Symptom Index; International Prostate Symptom Score <sup>1,3,12</sup> | Severity of lower urinary tract symptoms     | <ul style="list-style-type: none"><li>• Patient-administered, 7-item scale. Symptoms rated as mild (0-7), moderate (8-19), or severe (20-35)</li><li>• IPSS asks an additional question, rating QOL as delighted (0) to terrible (6).</li></ul> | <ul style="list-style-type: none"><li>• Minimum of 3-point change<sup>1,12</sup></li><li>• Minimum of 30% change<sup>13</sup></li></ul> |
| Benign Prostatic Hyperplasia Impact Index <sup>6,14</sup>   | Effect of urinary symptoms on health domains | Patient-administered, 4-item scale. Symptoms rated as absent (0) to severe (13).  | Minimum of 0.4-point change <sup>12</sup>   |

QOL: quality of life; IPSS: International Prostate Symptom Score; NR: Not reported.

## Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- a. To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- b. In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- c. To assess longer-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- d. Studies with duplicative or overlapping populations were excluded.

## **Systematic Reviews**

### **Initial Prostatic Urethral Lift Procedure**

Several systematic reviews on PUL have been published. They include a similar set of trials and noncomparative studies. Perera et al (2015) reported on the results of a systematic review and meta-analysis<sup>15</sup> of studies reporting outcomes after the PUL procedure, which included 7 prospective cohort studies,<sup>17-23</sup> a crossover study (Cantwell et al [2014]<sup>16</sup>), and the LIFT RCT (Roehrborn et al [2013],<sup>17</sup> McVary et al [2014]<sup>18</sup>). Shore (2015)<sup>19</sup> performed a systematic review of UroLift studies, which included the LIFT RCT (Roehrborn et al [2013]<sup>17</sup>; Roehrborn et al [2015]<sup>20</sup>; McVary et al [2014]<sup>18</sup>), a crossover study (Cantwell et al [2014]<sup>16</sup>), and 4 prospective cohort studies (Garrido Abad et al [2013]<sup>21</sup>; Chin et al [2012]<sup>22</sup>; Woo et al [2012]<sup>23</sup>; McNicholas et al [2013]<sup>24</sup>).

Jones et al (2016) performed a systematic review of UroLift studies with at least 12 months of follow-up.<sup>25</sup> Seven studies were identified, which included 4 noncomparative studies (Woo et al [2011],<sup>26</sup> Chin et al [2012],<sup>22</sup> McNicholas et al [2013],<sup>24</sup> Bozkurt et al [2016]<sup>27</sup>), a crossover study (Cantwell et al [2014]<sup>16</sup>), and 2 RCTs (LIFT<sup>17</sup> and BPH<sup>11</sup>).

The National Institute for Health and Care Excellence (2016) published technical guidance on prostatic lift procedures.<sup>28</sup> The National Institute for Care Excellence performed a literature search and data synthesis to support the development of the guidance. Studies selected were the same studies included in Perera et al (2015),<sup>15</sup> except for the exclusion of Hoffman et al (2012)<sup>29</sup> in the analysis.

Tanneru et al (2020) published a systematic review and meta-analysis of studies with at least 24 months of follow-up.<sup>30</sup> Five studies were included; 3 noncomparative studies (Chin et al [2012]<sup>22</sup> Rukstalis (2016)<sup>31</sup> Sievert et al [2020]<sup>32</sup> and 2 RCTs (LIFT and BPH).

Perera et al (2015), Shore (2015) and Jones et al (2016) analyzed data from the PUL arms of the studies only and the National Institute for Health and Care Excellence review was published before the BPH6 RCT. Therefore, these systematic reviews s will not be discussed further.

Jung et al (2019) published a Cochrane systematic review of PUL parallel-group RCTs published up to Jan 2019.<sup>33</sup> The 2 included RCTs (n=297) were the LIFT and BPH6 trials described in detail in the following section.<sup>17,34</sup> The two RCTs included different comparators and results were not combined meta-analytically. The authors used the GRADE approach to rate the certainty of the evidence. The conclusions were as follows:

- PUL appears less effective than TURP in improving urological symptoms, both in the short-term and long-term (low-certainty evidence);
- PUL may result in a similar QOL compared to TURP (low-certainty evidence);

- PUL may result in similar erectile function compared to TURP (moderate-certainty evidence);
- PUL may result in better ejaculatory function compared to TURP (moderate-certainty evidence);
- Rates of major adverse events are unclear (very low-certainty evidence);
- Rates of retreatment are unclear (very low-certainty evidence).

### Surgical Reinterventions

Miller et al (2020) reported results of a systematic review and meta-analysis on surgical reintervention rate of PUL.<sup>35</sup> Randomized or nonrandomized controlled studies, prospective and retrospective observational studies published through January 2020 were eligible for inclusion. Eleven studies (9 observational, 2 RCTs) were included with a total of 2,016 patients. The two RCTs (LIFT, BPH6) are described in the following section. There were 153 surgical reinterventions performed (TURP, 51.0%; repeat PUL, 32.7%, device explant, 19.6%). Overall, the annual rate of surgical reintervention was 6.0% per year (95% CI, 3.0% to 8.9%): 4.3% per year in studies with ≤1 year mean follow-up, 10.7% per year in studies with >1 year to 3 years mean follow-up, and 5.8% per year in one study with >3 years mean follow-up. No information was provided on the success of the reinterventions.

### Randomized Controlled Trials

Two RCTs of PUL have been performed. Key trial characteristics and study results are shown below in Tables 2 and 3, 6 and 7. Additionally, a brief description of each trial is provided in the following sections.

**Table 2. PUL Randomized Controlled Trial Characteristics**

| Study; Trial                                | Countries               | Sites | Dates              | Inclusion Criteria   | Baseline Prostate Volume, cm <sup>3</sup> | Interventions, n |            |
|---|-------------------------|-------|--------------------|--|---|------------------|------------|
|   |                         |       |                    |  |   | Active           | Comparator |
| Sonksen et al (2015) <sup>11</sup> ; BPH6   | Denmark, Germany, U.K.  | 10    | Feb 2012- Oct 2013 | Age ≥50 y, IPSS >12, prostate volume ≤60 cm <sup>3</sup>                                   | 16-59                                     | PUL=46           | TURP=45    |
| Roehrborn et al (2013) <sup>17</sup> ; LIFT | U.S., Canada, Australia | 19    | Feb-Dec 2011       | Age ≥50 y, IPSS ≥13, prostate volume 30-80 cm <sup>3</sup> , washed out of BPH medications | 30-77                                     | PUL=140          | Sham=66    |

BPH: benign prostatic hypertrophy; IPSS: International Prostate Symptom Score; PUL: prostatic urethral lift; TURP: transurethral resection of the prostate.

### BPH6 Study

Sonksen et al (2015) reported on the results of a multicenter RCT comparing the PUL procedure with TURP among individuals ages 50 and older with lower urinary tract symptoms, secondary to benign prostatic obstruction.<sup>11</sup> Eligible patients had an IPSS above 12, a Qmax of 15 mL/s or less for a 125-mL voided volume, a post-void residual volume less than 350 mL, and prostate volume of 60 cm<sup>3</sup> or less on ultrasound. Patients were excluded if there was median lobe obstruction in the prostate or signs of active infection. The trial used a novel composite endpoint, referred to as the BPH6, which included the following criteria:



- Lower urinary tract symptom relief: Reduction in IPSS by  $\geq 30\%$  within 12 months, relative to baseline
- Recovery experience: Self-assessed by patients as  $\geq 70\%$  within 1 month, using a visual analog scale
- Erectile function: Reduction in Sexual Health Inventory for Men (SHIM) score by  $\leq 6$  points within 12 months, relative to baseline
- Ejaculatory function: Emission of semen as assessed by question three in the Male Sexual Health Questionnaire for Ejaculatory Dysfunction (MSHQ-EjD)
- Continence preservation: Incontinence Severity Index  $\leq 4$  points at all follow-up visits
- Safety: No treatment-related adverse events exceeding grade 1 on the Clavien-Dindo classification system at time or procedure or any follow-up.

Patients were considered treatment responders if they met all six composite criteria. While this composite endpoint has not been previously validated, core components of the composite score have been independently validated in a clinical setting. The trial used a noninferiority design with a margin of 10% for the primary endpoint, BPH6. Study investigators modified 2 of the original endpoint definitions in the study's analysis, including changing the sexual function element assessment from a single time point (12 months) to assess sustained effects during 12 months of follow-up, and lowering the threshold of quality of recovery on a visual analog scale from 80 to 70.

**Table 3. Summary of Evidence From the BPH6 Study**

| Outcomes                          | 3 Months    |             | 12 Months   |             | 24 Months  |             |
|-----------------------------------|-------------|-------------|-------------|-------------|------------|-------------|
|                                   | PUL         | TURP        | PUL         | TURP        | PUL        | TURP        |
| Mean change in IPSS               |             |             |             |             |            |             |
| n                                 | 42          | 34          | 40          | 32          | 37         | 32          |
| Mean (SD)                         | -11.7 (8.5) | -11.8 (9.5) | -10.9 (7.9) | -15.4 (6.8) | -9.2 (9.2) | -15.3 (7.5) |
| P                                 | <0.001      | <0.001      | <0.001      | <0.001      | <0.001     | <0.001      |
| Comparison (p)                    | 0.978       |             | 0.013       |             | 0.004      |             |
| Change in IPSS QOL                |             |             |             |             |            |             |
| n                                 | 43          | 34          | 40          | 32          | 37         | 32          |
| Mean (SD)                         | -2.6 (1.7)  | -2.4 (2.0)  | -2.8 (1.8)  | -3.1 (1.6)  | -2.5 (1.8) | -3.3 (1.6)  |
| p                                 | <0.001      | <0.001      | <0.001      | <0.001      | <0.001     | <0.001      |
| Comparison (p)                    | 0.55        |             | 0.436       |             | 0.066      |             |
| Change in Qmax                    |             |             |             |             |            |             |
| n                                 | 33          | 25          | 32          | 29          | 27         | 27          |
| Mean (SD)                         | 4.2 (5.0)   | 12.7 (9.8)  | 4.0 (4.8)   | 13.7 (10.4) | 5.0 (5.5)  | 15.8 (16.5) |
| p                                 | <0.001      | 0.003       | <0.001      | 0.003       | <0.001     | 0.002       |
| Comparison (p)                    | <0.001      |             | <0.001      |             | 0.002      |             |
| Change in SHIM score              |             |             |             |             |            |             |
| n                                 | 38          | 27          | 32          | 27          | 29         | 28          |
| Mean (SD)                         | -0.7 (5.2)  | -1.0 (5.2)  | -0.1 (4.7)  | -0.9 (4.3)  | -0.2 (4.3) | -1.8 (4.90) |
| p                                 | 0.386       | 0.328       | 0.940       | 0.29        | 0.832      | 0.067       |
| Comparison (p)                    | 0.861       |             | 0.486       |             | 0.201      |             |
| Change in MSHQ-EjD function score |             |             |             |             |            |             |
| n                                 | 38          | 27          | 32          | 27          | 29         | 27          |
| Mean (SD)                         | -0.7 (2.1)  | -3.0 (4.1)  | 1.3 (3.3)   | -3.7 (4.1)  | 0.3 (3.4)  | -4.0 (4.6)  |
| p                                 | 0.251       | <0.001      |             | <0.001      | 0.666      | <0.001      |
| Comparison (p)                    | <0.001      |             | <0.001      |             | <0.001     |             |
| Change in MSHQ-EjD bother score   |             |             |             |             |            |             |
| n                                 | 38          | 28          | 32          | 27          | 29         | 27          |
| Mean (SD)                         | -0.7 (2.1)  | 0.2 (1.5)   | 0.5 (2.2)   | 0.0 (1.5)   | -0.1 (2.2) | -0.3 (1.9)  |
| p                                 | 0.062       | 0.470       | 0.214       | 0.896       | 0.734      | 0.415       |
| Comparison (p)                    | 0.069       |             | 0.359       |             | 0.771      |             |

|  |    |    |  |               |    |    |
|--|----|----|--|---------------|----|----|
| Composite score Comparison (95% CI); p | NR | NR | Response:52% Difference: 32% (10% to 51%); 0.005 | Response:20%  | NR | NR |
| Clavien-Dindo adverse events           |    |    |  |               |    |    |
| Grade 1, n (%) Adverse events          | NR | NR | 30 (68)<br>60                                    | 26 (74)<br>79 | NR | NR |
| Grade 2, n (%) Adverse events          | NR | NR | 3 (7)<br>3                                       | 4 (11)<br>5   | NR | NR |
| Grade 3, n (%) Adverse events          | NR | NR | 4 (9)<br>4                                       | 5 (14)<br>5   | NR | NR |

Adapted from Gratzke et al (2017).<sup>34</sup>

BPH: benign prostatic hypertrophy; CI: confidence interval; IPSS: International Prostate Symptom Score; MSHQ-EjD: Male Sexual Health Questionnaire for Ejaculatory Dysfunction; NR: not reported; PUL: prostatic urethral lift; Qmax: mean peak urinary flow rate; QOL: quality of life; SD: standard deviation; SHIM: Sexual Health Inventory for Men; TURP: transurethral resection of the prostate.

Ninety-one patients were randomized to TURP (n=45) or PUL (n=46). Ten patients in the TURP group and 1 patient in the PUL group declined treatment, leaving an analysis group of 80 subjects. The analysis was per-protocol, including 35 in the TURP group and 44 in the PUL group (87% of those randomized; 1 patient was excluded for violating the active urinary retention exclusion criterion). Groups were similar at baseline, except for the MSHQ-EjD function score. For procedure recovery, 82% of the PUL group achieved the recovery endpoint by 1 month compared with 53% of the TURP group (p=0.008). For the study's primary outcome, the proportion of participants who met the original BPH6 primary endpoint was 34.9% for the PUL group, and 8.6% for the TURP group (noninferiority p<0.001; superiority p=0.006). The modified BPH6 primary endpoint was met by 52.3% of the PUL group and 20.0% of the TURP group (noninferiority p<0.001; superiority p=0.005). Both groups demonstrated improvements over IPSS, IPSS quality of life score, BPH-II score, and Qmax over time, as described in Table 3. There were 60, grade 1 adverse events in 30 (68%) PUL patients and 79 adverse events in 26 (74%) TURP patients. The number of patients experiencing grade 2 and 3 adverse events was similar between groups. Intention-to-treat analyses were not reported.

Gratzke et al (2017) reported on 2-year results from BPH6.<sup>34</sup> Two additional patients were excluded from analysis: one TURP patient who discontinued participation; and one PUL patient who had a protocol violation. Composite scores for the two groups were not reported. Both groups continued to show significant improvements in IPSS score, IPSS quality of life, BPH-II score, and Qmax during the two-year follow-up, as described in Table 3. Six (14%) PUL patients and 2 (6%) TURP patients had secondary treatment (PUL, intradetrusor botulinum toxin, laser or TURP procedure), showing moderate durability over 2 years.

**Table 4. Relevance Limitations**

| Study | Population <sup>a</sup>              | Intervention <sup>b</sup> | Comparator <sup>c</sup>              | Outcomes <sup>d</sup>                | Follow-Up <sup>e</sup> |
|-------|--------------------------------------|---------------------------|--------------------------------------|--------------------------------------|------------------------|
| BPH6  | 3. Unclear history of BPH treatments |                           |                                      | 4: Primary outcome was not validated |                        |
| LIFT  | 3. Unclear history of BPH treatments |                           | 2: Men were washed out of medication |                                      |                        |

The study limitations stated in this table are those notable in the current review; this is not a comprehensive limitations assessment.

BPH: benign prostatic hypertrophy.

<sup>a</sup> Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

<sup>b</sup> Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest.

<sup>c</sup> Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively.

<sup>d</sup> Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported.

<sup>e</sup> Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

**Table 5. Study Design and Conduct Limitations**

| Study | Allocation <sup>3</sup> | Blinding <sup>b</sup>    | Selective Reporting <sup>c</sup> | Data Completeness <sup>d</sup>  | Power <sup>e</sup> | Statistical <sup>f</sup>                    |
|-------|-------------------------|--------------------------|----------------------------------|---|--------------------|---|
| BPH6  |                         | 1. Blinding not feasible |                                  | 6. Only per-protocol analysis presented   |                    |   |
| LIFT  |                         |                          |                                  | 1, 2, 5. High losses and/or exclusions in extended follow-up, only LOCF sensitivity analyses provided |                    | 3, 4. CI not reported for treatment effects |

The study limitations stated in this table are those notable in the current review; this is not a comprehensive limitations assessment. CI: confidence interval; LOCF: last observation carried forward.

<sup>a</sup> Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

<sup>b</sup> Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

<sup>c</sup> Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

<sup>d</sup> Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

<sup>e</sup> Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

<sup>f</sup> Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

### Subsection Summary: BPH6 Study

In the BPH6 study, PUL was both noninferior ( $p < 0.001$ ) and superior ( $p = 0.005$ ) to TURP for the study's composite endpoint. This endpoint was calculated using the concurrent achievement of validated measures of symptoms and complications and is sufficient to describe patient health outcomes. TURP was associated with greater improvements in urinary tract obstruction symptom outcomes and with greater declines in ejaculatory function compared with PUL.

### LIFT Study

#### Comparative Data

Roehrborn et al (2013) reported on results of the pivotal LIFT study, an RCT comparing PUL with sham control among 206 individuals ages 50 and older with lower urinary tract symptoms secondary to BPH.<sup>17</sup> Eligible patients had an American Urological Association Symptom Index (AUASI) score of 13 or greater, Qmax of 12 mL/s or less for a 125-mL voided volume, and a prostate volume between 30 and 80 mL. Patients were excluded if there was median lobe obstruction in the prostate, post-void obstruction of more than 250 mL, or signs of active infection. Patients underwent washout of BPH medications before enrollment; the washout period was two weeks for  $\alpha$ -blockers and three months for 5 $\alpha$ -reductase inhibitors. Patients were randomized to PUL ( $n = 140$ ) or sham control ( $n = 66$ ) and evaluated at 3 months post-procedure for the trial's primary efficacy endpoint. After that, all patients were unblinded, and sham control patients were permitted to undergo the PUL procedure. Fifty-three control subjects eventually underwent a PUL procedure. The analysis was intention-to-treat. The study met its primary efficacy endpoint, which was that the reduction in AUASI score at 3 months post-procedure had to be at least 25% greater after the PUL than the reduction in AUASI score seen with sham ( $p = 0.003$ ). The AUASI score decreased from 24.4 at baseline to 18.5 at 3-

month follow-up for sham control patients and from 22.2 at baseline to 11.2 at 3-month follow-up for PUL patients (see Table 4). The 3-month change in Qmax was 4.28 mL/s for PUL patients and 1.98 mL/s for sham control patients (p=0.005). Compared with sham control patients, PUL patients had greater improvements in quality of life scores and BPH-II score (see Table 5). Nine serious adverse events in seven patients were reported in the PUL group, and one serious adverse event was reported in the sham group during the first three months of follow-up. Limitations in the trial design are summarized in Tables 4 and 5.

McVary et al (2014) reported on sexual function outcomes in a subset of patients from the LIFT study.<sup>18</sup> At baseline, 53 (38%) PUL subjects and 23 (53%) sham control subjects were sexually inactive or had severe erectile dysfunction and were censored from the primary sexual function analysis. Scores on the SHIM, MSHQ-EjD function scale and the MSHQ-EjD bother scale did not differ significantly between groups.

**Table 6. Summary of LIFT Initial Trial Results**

| Study          | Change in IPSS | Change in IPSS QOL | Change in Qmax | Change in MSHQ-EjD Function | Change in MSHQ-EjD Bother | Any Adverse Events, n (%) | Serious Adverse Events, n (%) |
|----------------|----------------|--------------------|----------------|-----------------------------|---------------------------|---------------------------|-------------------------------|
| <b>LIFT</b>    |                |                    |                |                             |                           |                           |                               |
| N at 3 months  | 206            | 206                | 182            | 144                         | 177                       | 206                       | 206                           |
| PUL            | -11.1 (7.7)    | -2.2 (1.8)         | 4.3 (5.2)      | 2.2 (2.5)                   | -0.8 (1.5)                | 122 (87%)                 | 7 (5%)                        |
| Adverse events |                |                    |                |                             |                           |                           |                               |
| Sham           | -5.9 (7.7)     | -1.0 (1.5)         | 2.0 (4.9)      | 1.7 (2.6)                   | -0.7 (1.6)                | 43 (52%)                  | 1 (1.5%)                      |
| Adverse events |                |                    |                |                             |                           |                           |                               |
| TE (p)         | NR<br>(0.003)  | NR<br>(<0.001)     | NR<br>(0.005)  | NR<br>(0.283)               | NR<br>(0.60)              | NR                        | NR                            |

Adapted from Roehrborn et al (2013).<sup>17</sup>

Values are mean (standard deviation) unless otherwise indicated.

IPSS: International Prostate Symptom Score; MSHQ-EjD: Male Sexual Health Questionnaire for Ejaculatory Dysfunction; NR: not reported; PUL: prostatic urethral lift; Qmax: mean peak urinary flow rate; QOL: quality of life; TE: treatment effect

**Table 7. Summary of Evidence for LIFT Study, Including Participants in the PUL Group**

| Outcomes            | 3 Months        | 1 Year          | 2 Years         | 3 Years         | 5 Years          |
|---------------------|-----------------|-----------------|-----------------|-----------------|------------------|
| n                   | 140             | 129             | 118             | 109             | 87               |
| Death/LTFU          | 0               | 2               | 7               | 2               | 18               |
| Protocol deviations | 3               | 0               | 0               | 1               | 0                |
| Retreatment         | 0               | 6               | 4               | 6               | 4                |
| Change in IPSS      |                 |                 |                 |                 |                  |
| n                   | 136             | 123             | 103             | 93              | 72               |
| Change              | -11.4 (7.72)    | -10.61 (7.51)   | -9.13 (7.62)    | -8.83 (7.41)    | -35.9%           |
| 95% CI              | -12.45 to -9.83 | -11.95 to -9.27 | -10.62 to -7.64 | -10.35 to -7.30 | -44.4% to -27.3% |
| p                   | <0.001          | <0.001          | <0.001          | <0.001          | <0.001           |
| Change in IPSS QOL  |                 |                 |                 |                 |                  |
| n                   | 136             | 123             | 103             | 93              | 72               |
| Change              | -2.22 (1.78)    | -2.31 (1.60)    | 2.19 (1.72)     | -2.25 (1.72)    | -50.3            |
| 95% CI              | -2.52 to -1.92  | -2.59 to -2.02  | -2.53 to -1.86  | -2.60 to -1.89  | -58.4% to -42.2% |
| p                   | <0.001          | <0.001          | <0.001          | <0.001          | <0.001           |
| Change in Qmax      |                 |                 |                 |                 |                  |
| n                   | 122             | 102             | 86              | 69              | 52               |
| Change              | 4.29 (5.16)     | 4.03 (4.96)     | 4.21 (5.09)     | 3.47 (5.00)     | 44.3%            |
| 95% CI              | 3.36 to 5.21    | 3.06 to 5.00    | 3.12 to 5.30    | 2.27 to 4.67    | 29.4% to 59.1%   |

|                                   |                |                |                |                |                 |
|-----------------------------------|----------------|----------------|----------------|----------------|-----------------|
| p                                 | <0.001         | <0.001         | <0.001         | <0.001         | <0.001          |
| Change in SHIM score              |                |                |                |                |                 |
| n                                 | 91             | 87             | 72             | 66             | NR              |
| Change                            | 1.27 (4.65)    | 0.70 (5.12)    | 1.06 (4.78)    | 0.53 (4.41)    | NR              |
| 95% CI                            | 0.31 to 2.24   | -0.39 to 1.79  | -0.07          | To 2.18        | -0.55 to 1.62   |
| p                                 | 0.005          | 0.299          | 0.046          | 0.338          | NR              |
| Change in MSHQ-EjD function score |                |                |                |                |                 |
| n                                 | 91             | 87             | 72             | 66             | 49              |
| Change                            | 2.31 (2.58)    | 1.56 (2.68)    | 1.08 (2.51)    | 0.56 (2.48)    | 9.3%            |
| 95% CI                            | 1.77 to 2.85   | 0.99 to 2.13   | 0.49 to 1.67   | -0.05 to 1.17  | -3.8% to 22.5%  |
| p                                 | <0.001         | <0.001         | <0.001         | 0.013          | 0.096           |
| Change in MSHQ-EjD bother score   |                |                |                |                |                 |
| n                                 | 91             | 87             | 72             | 66             | 49              |
| Change                            | -1.07 (1.44)   | -0.76 (-1.55)  | 0.63 (1.51)    | -0.59 (1.52)   | -6.3%           |
| 95% CI                            | -1.37 to -0.77 | -1.09 to -0.43 | -0.98 to -0.27 | -0.96 to -0.22 | -31.5% to 18.8% |
| p                                 | <0.001         | <0.001         | <0.001         | <0.001         | 0.019           |

Adapted from Roehrborn et al (2015)<sup>36</sup> for data from 3 months to 3 years and Roehrborn et al (2017)<sup>37</sup> for data for 5 years. While not specifically indicated, change values likely represent means and standard deviations. CI: 95% confidence interval; IPSS: International Prostate Symptom Score; LTFU: lost to follow-up; MSHQ-EjD: Male Sexual Health Questionnaire for Ejaculatory Dysfunction; NR: not reported; PUL: prostatic urethral lift; Qmax: mean peak urinary flow rate; QOL: quality of life; SHIM: Sexual Health Inventory for Men.

### Follow-Up of Sham-Assigned Crossover Participants

Cantwell et al (2014) reported on 12-month outcomes for 53 subjects in the LIFT sham control group who underwent PUL after unblinding at 3 months postprocedure.<sup>16</sup> Crossover (unblinded) patients had a change in IPSS from 23.4 to 12.3 at 3 months post-procedure compared with the change in IPSS from 25.2 to 20.2 at 3 months after the sham procedure. Subjects had greater improvements in BPH-II score in the crossover period (-3.3) than in the sham period (-1.9;  $p=0.024$ ) but did not report significant differences in improvement in Qmax. Change in sexual function scores did not differ significantly after the sham procedure compared with after active procedure.

Rukstalis et al (2016) reported on 24-month outcomes for 42 of the 53 participants in the LIFT sham group who underwent PUL after unblinding.<sup>31</sup> During the 24 months, 4 patients were known to have had TURP, and 1 patient required additional PUL implants. The change in IPSS from baseline to 24 months was -9.6 (-35%; 95% CI, not reported;  $p<0.001$ ) and there was significant score improvements in Qmax, BPH-II scores, and quality of life. There were no significant changes compared with baseline for SHIM scores; however, MSHQ-EjD scores improved by 41% ( $p<0.001$ ).

### Follow-Up of PUL-Assigned Participants

Roehrborn et al (2015) reported on 3-year results from patients randomized to PUL in the LIFT study.<sup>20</sup> After exclusion of 11 subjects who were lost to follow-up, 36 subjects with missing data, protocol deviations, medication treatment for BPH, or other prostate procedures, and 15 subjects who underwent surgical retreatment for lower urinary tract symptoms (6 with repeat PUL procedures, 9 with TURP or laser vaporization), the 3-year effectiveness analysis included 93 (66%) of the original 140 subjects. For subjects with follow-up data, change in IPSS was -8.83 (95% CI, -10.35 to -7.30;  $p<0.001$ ). Significant improvements were also reported for the quality of life score, BPH-II score, and Qmax. Sexual function was unchanged. Implants were removed from ten participants. No analyses were performed to assess how sensitive the results were to changes in the assumptions about the considerable amount of missing data.

Roehrborn et al (2016) reported on 4-year results from patients randomized to PUL in the LIFT study.<sup>36</sup> Of the 140 originally randomized patients, 32 were lost by the 4-year follow-up visit (6 losses were deaths). Of the remaining 108 patients for whom data were available, an additional 29 patients were excluded from analysis for BPH retreatment or protocol deviations. For the 79 (56%) of the 140 subjects included in the analysis, change in IPSS score was -8.8 (precision not given) or -41% (95% CI, -49% to -33%;  $p < 0.001$ ). Significant improvements (vs. baseline) were also reported for scores relating to the quality of life, BPH-II, and Qmax. Authors reported that 14% “of the 140 originally enrolled” participants had surgical retreatment at some point during the 4 years; however, the 4-year follow-up included 79 patients, so the denominator for the 14% is not clear, and estimated retreatment rates are likely underestimated since individuals lost to follow-up could also have received retreatment. Attributes of patients who received retreatment were not analyzed. SHIM scores did not differ statistically from baseline.

Roehrborn et al (2017) reported on 5-year results from patients randomized to PUL in the LIFT study.<sup>37</sup> The authors reported two analyses. The first was called a per-protocol analysis, which censored patients who had additional BPH procedures, started a BPH medication or had a protocol deviation. A second analysis was called intention-to-treat analysis, which used last observation carried forward to impute values that were censored in the per-protocol analysis. While there were 104 participants with 5-year data, only 72 patients (approximately 50% of those randomized) were included in the per-protocol analysis after exclusion for protocol violations, additional BPH procedures, or treatment with BPH medication. In the intention-to-treat analysis, change in IPSS was -7.85 at 5 years (-35%; 95% CI, -41% to -29%;  $p < 0.001$ ). In the per-protocol analysis, change in IPSS was -7.56 at 5 years (-35.9%; 95% CI, -44% to -27%). Significant improvements, compared with baseline, continued to be reported for scores associated with quality of life, Qmax, and BPH-II. Of the limited number of patients that remained in the analysis, 13.6% had surgical reintervention by 5 years.

### **Subsection Summary: LIFT Study**

The LIFT RCT compared PUL with a sham procedure in individuals who had completed a washout period for BPH medications before enrollment. The PUL procedure was associated with greater improvements in lower urinary tract symptoms compared with sham; additionally, the PUL procedure was found to have not worsened sexual function after three months of follow-up. After 3 months, patients were given the option to have PUL surgery and about 80% of the sham patients did so. Functional improvements, compared with baseline, appear durable in patients over two years and are consistent with the BPH6 study. Follow-up over three to five years was notable for a high number of patients who were either excluded or lost.

### **Section Summary: Randomized Controlled Trials**

The BPH6 study demonstrated that PUL is noninferior to TURP when assessed by a composite score, which reflects concurrent improvements in validated scales of symptoms, safety, and sexual function. These findings are reflected in the analysis of the individual aspects of the composite score. PUL demonstrates measurable improvements in urinary symptoms to two years and is superior to TURP in preserving sexual function. These findings were confirmed in the LIFT study, which compared PUL with a sham treatment. Prior to crossover at three months, patients were found to have greater improvement in urinary symptoms and preserved sexual function relative to patients receiving sham treatment. After 3 months, 80% of patients who had received a sham treatment chose to have the PUL procedure. Patients treated with PUL had improvement of urinary symptoms with preservation of sexual function, consistent with the BPH6 study. These findings were preserved in a subset

of patients over three to five years; a high number of patients were either excluded or lost to follow-up during this time. The BPH6 and LIFT RCTs excluded men with median lobe obstruction.

### Noncomparative Studies

The approved indications for PUL have expanded since the original approval to include men with median lobe obstruction and those with prostate volume between 80cc and 100cc. Neither of these expansions have supporting RCTs.

### Median Lobe Obstruction

Several noncomparative studies were published including men without median lobe obstruction. Since RCTs with long-term follow-up exist for this population, these noncomparative studies will not be discussed in further detail. Rukstalis et al (2019) reported results of the MedLift study, the study used to support the expansion of the Food and Drug Administration clearance for PUL to include obstructive median lobes.<sup>38</sup> MedLift was a single-arm study enrolling 45 men with eligibility criteria identical to LIFT except requiring obstructive median lobes. Results in the MedLift cohort were compared to the LIFT historical cohort. Characteristics are shown in Table 8 and results are shown in Table 9. One patient required surgical retreatment and no implants were removed over the 12 months of follow-up.

**Table 8. Summary of Characteristics of Key Non-Comparative Studies**

| Study            | Country | Sites | Participants  | Treatment Delivery                                | Follow-up |
|------------------|---------|-------|---|---|-----------|
| Rukstalis (2019) | US      | 9     | n=45<br>Men ages 50+ with IPSS>13, Qmax <=12 mL/s, 30 to 80 cc intraurethral prostatic volume and, OMLa | UroLift PUL procedure with median lobe deployment | 12 months |

<sup>a</sup>OML (Obstructive Median Lobe) was defined as excessive posterior tissue that precludes a normal lateral lobe procedure.

**Table 9. Summary Results of Key Non-Comparative Studies**

| Study                                       | IPSS                   | IPSS QOL              | Qmax                 | SHIM                |
|---|------------------------|-----------------------|----------------------|---------------------|
| Rukstalis (2019)                            | At 12 m                | At 12 m               | At 12 m              | At 12 m             |
| N   | 44                     | 44                    | 37                   | 38                  |
| Change from baseline, mean (SD);<br>p-value | -13.5 (7.7)<br>p<0.001 | -3.0 (1.5)<br>p<0.001 | 6.4 (7.4)<br>p<0.001 | 1.2 (4.3)<br>p=0.04 |

CI: 95% confidence interval; IPSS: International Prostate Symptom Score; Qmax: mean peak urinary flow rate; QOL: quality of life; SHIM: Sexual Health Inventory for Men; SD: standard deviation.

The purpose of the limitation tables (see Tables 10 and 11) is to display notable limitations identified in each study. This information is synthesized as a summary of the body of evidence following each table and provides the conclusions on the sufficiency of the evidence supporting the position statement.

**Table 10. Relevance Limitations**

| Study | Population <sup>a</sup> | Intervention <sup>b</sup> | Comparator <sup>c</sup> | Outcomes <sup>d</sup> | Follow-Up <sup>e</sup> |
|-------|-------------------------|---------------------------|-------------------------|-----------------------|------------------------|
|-------|-------------------------|---------------------------|-------------------------|-----------------------|------------------------|

|                  |                                      |                             |  |                                       |
|------------------|--------------------------------------|-----------------------------|--|---------------------------------------|
| Rukstalis (2019) | 3. Unclear history of BPH treatments | 2: No concurrent comparator | 3: Reporting of adverse events was qualitative; rates not reported | 1, 2: Only 12 m of follow-up reported |
|------------------|--------------------------------------|-----------------------------|--|---------------------------------------|

The study limitations stated in this table are those notable in the current review; this is not a comprehensive limitations assessment. BPH: benign prostatic hypertrophy.

<sup>a</sup> Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

<sup>b</sup> Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest.

<sup>c</sup> Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively.

<sup>d</sup> Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported.

<sup>e</sup> Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

**Table 11. Study Design and Conduct Limitations**

| Study            | Allocation <sup>a</sup> | Blinding <sup>b</sup> | Selective Reporting <sup>c</sup> | Data Completeness <sup>d</sup>      | Power <sup>e</sup> | Statistical <sup>f</sup> |
|------------------|-------------------------|-----------------------|----------------------------------|-------------------------------------|--------------------|--------------------------|
| Rukstalis (2019) | 1,2: Not randomized     | 1,2: No blinding      |                                  | >15% missing data for Qmax and SHIM |                    | 3: CIs not reported      |

The study limitations stated in this table are those notable in the current review; this is not a comprehensive limitations assessment.

CI: confidence interval; Qmax: mean peak urinary flow; SHIM: Sexual Health Inventory for Men.

<sup>a</sup> Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

<sup>b</sup> Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

<sup>c</sup> Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

<sup>d</sup> Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

<sup>e</sup> Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

<sup>f</sup> Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

### Prostate Volume Greater Than 80 mL

Sievert et al (2019) reported results of a noncomparative study that included 5 men with prostate volume greater than 80 mL.<sup>32</sup> Results were not presented stratified by prostate volume.

Eure (2019)<sup>40</sup> included 38 men with prostate volume >80 mL. Although the authors reported that 'no significant differences in symptom response emerged based on prostate volume', results were not presented stratified by prostate volume.

Bozkurt (2016)<sup>27</sup> Woo (2012)<sup>23</sup> and Chin (2012)<sup>22</sup> included men with prostate volume greater than 80 mL but had mean volume in the 40 to 60 range and it is unclear how many patients had volume greater than 80.

Given the limited amount of published data on outcomes for men with prostate volume greater than 80 mL and limited follow-up, the risks and benefits cannot be evaluated.

### Section Summary: Noncomparative Studies

One single-arm study (n=45) including men with obstructive median lobes has been conducted and was used to support the Food and Drug Administration expansion of the PUL indication to include these men. Symptom scores and QOL appeared to improve by statistically and clinically significant amounts and were similar in magnitude to improvements reported in the



original LIFT study. Rates of adverse events were not reported. Design and conduct limitations preclude interpretation.

Noncomparative studies have included a small number of men with larger prostate volume but have generally not reported results stratified by prostate volume. One study presented data for 20 men with less than 6 months of follow-up.

## **SUMMARY OF EVIDENCE**

For individuals who have lower urinary tract obstruction symptoms due to BPH and receive a PUL, the evidence includes systematic reviews, randomized controlled trials, and noncomparative studies. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and treatment-related morbidity. One randomized controlled trial, the BPH6 study, compared the PUL procedure with transurethral resection of the prostate and reported that the PUL procedure was noninferior for the study's composite endpoint, which required concurrent fulfillment of six independently validated measures of symptoms, safety, and sexual health. While transurethral resection of the prostate was superior to PUL in managing lower urinary tract symptoms, PUL did provide significant symptom improvement over two years. PUL was further superior to transurethral resection of the prostate in preserving sexual function. These findings were corroborated by another randomized controlled trial (the LIFT study), which compared PUL with sham control. Patients underwent washout of BPH medications before enrollment. LIFT reported that patients with the PUL procedure, compared with patients who had sham surgery and no BPH medication, had greater improvements in lower urinary tract symptoms without worsened sexual function at three months. After 3 months, patients were given the option to have PUL surgery; 80% of the patients with sham procedures chose that option. Publications from this trial reported that functional improvements were durable over 3-, 4-, and 5-year follow-ups in a subset of patients treated with PUL; there was a high number of exclusions and loss to follow-up in that group. The BPH6 and LIFT RCTs included men with prostate volume up to 80 cm<sup>3</sup> and excluded men with median lobe obstruction. Selection criteria of patients for whom evidence is sufficient to support improvement are derived from clinical trial eligibility criteria, product labeling, and clinical input. The evidence is sufficient to determine the effects of the technology on health outcomes.

## **Ongoing and Unpublished Clinical Trials**

A search of ClinicalTrials.gov did not identify any ongoing or unpublished trials that would likely influence this review.

## **CLINICAL INPUT**

Clinical input is sought to help determine whether the use of prostatic urethral lift (PUL) for individuals with moderate-to-severe lower urinary tract obstruction symptoms due to benign prostatic hyperplasia (BPH) would provide a meaningful clinical benefit, defined as improved net health outcome, and whether this use is consistent with generally accepted medical practice.

## **RESPONDENTS**

Clinical input was provided by the following specialty societies and physician members identified by a specialty society or clinical health system:

- John Lin, MD, Urology; identified by American Urological Association (AUA)<sup>a</sup>
- Anonymous, MD, Urology; identified by AUA<sup>a</sup>

- Anonymous, MD, Urology; identified by University of California San Francisco (UCSF) Medical Center
- Anonymous, MD, Urology; identified by UCSF Medical Center

<sup>a</sup> Indicates that conflicts of interest related to the topic where clinical input is being sought were identified by this respondent

Clinical input provided by the specialty society at an aggregate level is attributed to the specialty society. Clinical input provided by a physician member designated by a specialty society or health system is attributed to the individual physician and is not a statement from the specialty society or health system. Specialty society and physician respondents participating in the Evidence Street® clinical input process provide a review, input, and feedback on topics being evaluated by Evidence Street. However, participation by a specialty society and/or physician member designated by a specialty society or health system in the clinical input process does not imply an endorsement or explicit agreement with the Evidence Opinion published by BCBSA or any Blue Plan.

## CLINICAL INPUT RESPONSES

| Clinical Indication  | Respondent | Identified by | Confidence Level That Clinical Use Expected to Provide Meaningful Clinical Benefit |              |     |     |      |      |              |     |           |      | Confidence Level That Clinical Use is Consistent with Generally Accepted Medical Practice |              |     |     |      |      |              |     |     |      |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
|--|------------|---------------|--|--------------|-----|-----|------|------|--------------|-----|-----------|------|---|--------------|-----|-----|------|------|--------------|-----|-----|------|-----|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|
|  |            |               | NO   |              |     |     |      | YES  |              |     |           |      | NO  |              |     |     |      | YES  |              |     |     |      |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
|  |            |               | High   | Intermediate | Low | Low | High | High | Intermediate | Low | Low       | High | High  | Intermediate | Low | Low | High | High | Intermediate | Low | Low | High |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| Yes or No  | 5          | 4             | 3  | 2            | 1   | 1   | 2    | 3    | 4            | 5   | Yes or No | 5    | 4   | 3            | 2   | 1   | 1    | 2    | 3            | 4   | 5   |      |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| PUL for individuals with moderate to severe lower urinary tract obstruction symptoms (due to benign prostatic hyperplasia)                               | Anonymous* | AUA           | YES  |              |     |     |      |      |              |     |           |      |   |              |     |     |      |      |              |     |     |      |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
|  | Dr. Lin*   | AUA           | YES  |              |     |     |      |      |              |     |           |      |   |              |     |     |      |      |              |     |     |      | YES |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
|  | Anonymous  | UCSF Med Ctr. | YES  |              |     |     |      |      |              |     |           |      |   |              |     |     |      |      |              |     |     |      | NO  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
|  | Anonymous  | UCSF Med Ctr. | YES  |              |     |     |      |      |              |     |           |      |   |              |     |     |      |      |              |     |     |      | YES |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| PUL for individuals with moderate to severe lower urinary tract obstruction symptoms (due to benign prostatic hyperplasia) and failed medical management | Anonymous* | AUA           | YES  |              |     |     |      |      |              |     |           |      |   |              |     |     |      |      |              |     |     | YES  |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
|  | Dr. Lin*   | AUA           | YES  |              |     |     |      |      |              |     |           |      |   |              |     |     |      |      |              |     |     | YES  |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
|  | Anonymous  | UCSF Med Ctr. | YES  |              |     |     |      |      |              |     |           |      |   |              |     |     |      |      |              |     |     | YES  |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
|  | Anonymous  | UCSF Med Ctr. | YES  |              |     |     |      |      |              |     |           |      |   |              |     |     |      |      |              |     |     | YES  |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| PUL for individuals with moderate to severe lower urinary tract obstruction symptoms (due to benign prostatic hyperplasia) and not surgical candidate    | Anonymous* | AUA           | YES  |              |     |     |      |      |              |     |           |      |   |              |     |     |      |      |              |     |     | YES  |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
|  | Dr. Lin*   | AUA           | YES  |              |     |     |      |      |              |     |           |      |   |              |     |     |      |      |              |     |     | YES  |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
|  | Anonymous  | UCSF Med Ctr. | YES  |              |     |     |      |      |              |     |           |      |   |              |     |     |      |      |              |     |     | YES  |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
|  | Anonymous  | UCSF Med Ctr. | YES  |              |     |     |      |      |              |     |           |      |   |              |     |     |      |      |              |     |     | YES  |     |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |

AUA: American Urological Association; UCSF Med Ctr: University of California San Francisco Medical Center; PUL: prostatic urethral lift.

\* Indicates that information on conflicts of interest related to the topic where clinical input is being sought were identified by this respondent (see Appendix 1).

## SUPPLEMENTAL INFORMATION

### CLINICAL INPUT FROM PHYSICIAN SPECIALTY SOCIETIES AND ACADEMIC MEDICAL CENTERS

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

## PRACTICE GUIDELINES AND POSITION STATEMENTS

### American Urological Association

The American Urological Association (2018) published guidelines on the surgical management of LUTS attributed to BPH; the 2018 guidelines were amended in 2019 and 2020.<sup>7</sup> The guidelines made the following recommendations and statements regarding PUL.

- “ PUL [prostatic urethral lift] may be offered as an option for patients with LUTS [lower urinary tract symptoms] attributed to BPH [benign prostatic hyperplasia] provided prostate volume <80g and verified absence of an obstructive middle lobe.”  
“Moderate Recommendation; Evidence Level: Grade C indicating “Benefits > Risks/Burdens (or vice versa); Net benefit (or net harm) appears moderate. Applies to most patients in most circumstances but better evidence is likely to change confidence”
- “PUL may be offered to eligible patients concerned with erectile and ejaculatory function for the treatment of with LUTS attributed to BPH.”  
“Conditional Recommendation; Evidence Level: Grade C indicating “Risks/Burdens unclear; Alternative strategies may be equally reasonable. Better evidence likely to change confidence”
- “Clinicians should inform patients of the possibility of treatment failure and the need for additional or secondary treatments when considering surgical and minimally-invasive treatments for LUTS secondary to BPH.”

### National Institute for Health and Care Excellence (NICE)<sup>41</sup>

The National Institute for Health and Care Excellence (2014) published guidance on urethral lift implants to treat lower urinary tract symptoms (LUTS) secondary to benign prostatic hyperplasia (BPH). The guidance stated:

“Current evidence on the efficacy and safety of insertion of prostatic urethral lift implants to treat lower urinary tract symptoms secondary to benign prostatic hyperplasia is adequate to support the use of this procedure.”

The Institute (2015) published guidance on the use of UroLift for treating LUTS of BPH. The guidance stated: “the UroLift system is effective in relieving symptoms of benign prostatic hyperplasia” and “the UroLift system should be considered as an alternative to current surgical procedures for use in a day-case setting in individuals with lower urinary tract symptoms of benign prostatic hyperplasia who are aged 50 years and older and who have a prostate of less than 100 ml without an obstructing middle lobe.”

---

## Government Regulations

### National/Local:

There is no national coverage determination on this topic. Medicare has established a fee for BCBSA codes 52441 and 52442.

4/14/15-It has been confirmed by WPS staff that 52441 and 52442 are payable if the services are medically necessary and documented as such by the provider in the medical record.

*(The above Medicare information is current as of the review date for this policy. However, the coverage issues and policies maintained by the Centers for Medicare & Medicare Services [BCBSA, formerly HCFA] are updated and/or revised periodically. Therefore, the most current BCBSA information may not be contained in this document. For the most current information, the reader should contact an official Medicare source.)*

---

## **Related Policies**

N/A

---

## **References**

1. Sarma AV, Wei JT. Clinical practice. Benign prostatic hyperplasia and lower urinary tract symptoms. *N Engl J Med.* Jul 19 2012; 367(3): 248-57. PMID 22808960
2. Barry MJ, Fowler FJ, O'Leary MP, et al. Measuring disease-specific health status in men with benign prostatic hyperplasia. Measurement Committee of The American Urological Association. *Med Care.* Apr 1995; 33(4 Suppl): AS145-55. PMID 7536866
3. O'leary MP. Validity of the bother score in the evaluation and treatment of symptomatic benign prostatic hyperplasia. *Rev Urol.* 2005; 7(1): 1-10. PMID 16985801
4. Djavan B, Marberger M. A meta-analysis on the efficacy and tolerability of alpha1-adrenoceptor antagonists in patients with lower urinary tract symptoms suggestive of benign prostatic obstruction. *Eur Urol.* 1999; 36(1): 1-13. PMID 10364649
5. Foster HE, Barry MJ, Dahm P, et al. Surgical Management of Lower Urinary Tract Symptoms Attributed to Benign Prostatic Hyperplasia: AUA Guideline. *J Urol.* Sep 2018; 200(3): 612-619. PMID 29775639
6. Reich O, Gratzke C, Bachmann A, et al. Morbidity, mortality and early outcome of transurethral resection of the prostate: a prospective multicenter evaluation of 10,654 patients. *J Urol.* Jul 2008; 180(1): 246-9. PMID 18499179
7. Foster HE, Barry MJ, Dahm P, et al. Surgical Management of Lower Urinary Tract Symptoms Attributed to Benign Prostatic Hyperplasia: AUA GUIDELINE. 2020. [https://www.auanet.org/guidelines/benign-prostatic-hyperplasia-\(bph\)-guideline](https://www.auanet.org/guidelines/benign-prostatic-hyperplasia-(bph)-guideline). Accessed September 2020.
8. Sundaram D, Sankaran PK, Raghunath G, et al. Correlation of Prostate Gland Size and Uroflowmetry in Patients with Lower Urinary Tract Symptoms. *J Clin Diagn Res.* May 2017; 11(5): AC01-AC04. PMID 28658743
9. Rosen RC, Catania JA, Althof SE, et al. Development and validation of four-item version of Male Sexual Health Questionnaire to assess ejaculatory dysfunction. *Urology.* May 2007; 69(5): 805-9. PMID 17482908
10. Cappelleri JC, Rosen RC. The Sexual Health Inventory for Men (SHIM): a 5-year review of research and clinical experience. *Int J Impot Res.* Jul-Aug 2005; 17(4): 307-19. PMID 15875061
11. Sonksen J, Barber NJ, Speakman MJ, et al. Prospective, randomized, multinational study of prostatic urethral lift versus transurethral resection of the prostate: 12-month results from the BPH6 study. *Eur Urol.* Oct 2015; 68(4): 643-52. PMID 25937539
12. Barry MJ, Williford WO, Chang Y, et al. Benign prostatic hyperplasia specific health status measures in clinical research: how much change in the American Urological Association symptom index and the benign prostatic hyperplasia impact index is perceptible to patients?. *J Urol.* Nov 1995; 154(5): 1770-4. PMID 7563343
13. Roehrborn CG, Wilson TH, Black LK. Quantifying the contribution of symptom improvement to satisfaction of men with moderate to severe benign prostatic hyperplasia: 4-year data from the CombAT trial. *J Urol.* May 2012; 187(5): 1732-8. PMID 22425127

14. McVary KT, Roehrborn CG, Avins AL, et al. American Urological Association Guideline: Management of Benign Prostatic Hyperplasia (BPH). 2010 (affirmed 2014); [http://www.auanet.org/guidelines/benign-prostatic-hyperplasia-\(2010-reviewed-andvalidity-confirmed-2014\)](http://www.auanet.org/guidelines/benign-prostatic-hyperplasia-(2010-reviewed-andvalidity-confirmed-2014)). Accessed September 2020.
15. Perera M, Roberts MJ, Doi SA, et al. Prostatic urethral lift improves urinary symptoms and flow while preserving sexual function for men with benign prostatic hyperplasia: a systematic review and meta-analysis. *Eur Urol.* Apr 2015; 67(4): 704-13. PMID 25466940
16. Cantwell AL, Bogache WK, Richardson SF, et al. Multicentre prospective crossover study of the 'prostatic urethral lift' for the treatment of lower urinary tract symptoms secondary to benign prostatic hyperplasia. *BJU Int.* Apr 2014; 113(4): 615-22. PMID 24765680
17. Roehrborn CG, Gange SN, Shore ND, et al. The prostatic urethral lift for the treatment of lower urinary tract symptoms associated with prostate enlargement due to benign prostatic hyperplasia: the L.I.F.T. Study. *J Urol.* Dec 2013; 190(6): 2161-7. PMID 23764081
18. McVary KT, Gange SN, Shore ND, et al. Treatment of LUTS secondary to BPH while preserving sexual function: randomized controlled study of prostatic urethral lift. *J Sex Med.* Jan 2014; 11(1): 279-87. PMID 24119101
19. Shore N. A Review of the Prostatic Urethral Lift for Lower Urinary Tract Symptoms: Symptom Relief, Flow Improvement, and Preservation of Sexual Function in Men With Benign Prostatic Hyperplasia. *Curr Bladder Dysfunct Rep. NA* 2015; 10(2): 186-192. PMID 25984251
20. Roehrborn CG, Rukstalis DB, Barkin J, et al. Three year results of the prostatic urethral L.I.F.T. study. *Can J Urol.* Jun 2015; 22(3): 7772-82. PMID 26068624
21. Garrido Abad P, Coloma Del Peso A, Sinues Ojas B, et al. [Urolift(R), a new minimally invasive treatment for patients with low urinary tract symptoms secondary to BPH. Preliminary results]. *Arch Esp Urol.* Jul-Aug 2013; 66(6): 584-91. PMID 23985459
22. Chin PT, Bolton DM, Jack G, et al. Prostatic urethral lift: two-year results after treatment for lower urinary tract symptoms secondary to benign prostatic hyperplasia. *Urology.* Jan 2012; 79(1): 5-11. PMID 22202539
23. Woo HH, Bolton DM, Laborde E, et al. Preservation of sexual function with the prostatic urethral lift: a novel treatment for lower urinary tract symptoms secondary to benign prostatic hyperplasia. *J Sex Med.* Feb 2012; 9(2): 568-75. PMID 22172161
24. McNicholas TA, Woo HH, Chin PT, et al. Minimally invasive prostatic urethral lift: surgical technique and multinational experience. *Eur Urol.* Aug 2013; 64(2): 292-9. PMID 23357348
25. Jones P, Rajkumar GN, Rai BP, et al. Medium-term Outcomes of Urolift (Minimum 12 Months Follow-up): Evidence From a Systematic Review. *Urology.* Nov 2016; 97: 20-24. PMID 27208817
26. Woo HH, Chin PT, McNicholas TA, et al. Safety and feasibility of the prostatic urethral lift: a novel, minimally invasive treatment for lower urinary tract symptoms (LUTS) secondary to benign prostatic hyperplasia (BPH). *BJU Int.* Jul 2011; 108(1): 82-8. PMID 21554526
27. Bozkurt A, Karabakan M, Keskin E, et al. Prostatic Urethral Lift: A New Minimally Invasive Treatment for Lower Urinary Tract Symptoms Secondary to Benign Prostatic Hyperplasia. *Urol Int.* 2016; 96(2): 202-6. PMID 26613256
28. Ray A, Morgan H, Wilkes A, et al. The Urolift System for the Treatment of Lower Urinary Tract Symptoms Secondary to Benign Prostatic Hyperplasia: A NICE Medical Technology Guidance. *Appl Health Econ Health Policy.* Oct 2016; 14(5): 515-26. PMID 26832146
29. Hoffman RM, Monga M, Elliott SP, et al. Microwave thermotherapy for benign prostatic hyperplasia. *Cochrane Database Syst Rev.* Sep 12 2012; (9): CD004135. PMID 22972068
30. Tanneru K, Gautam S, Norez D, et al. Meta-analysis and systematic review of intermediate-term follow-up of prostatic urethral lift for benign prostatic hyperplasia. *Int Urol Nephrol.* Jun 2020; 52(6): 999-1008. PMID 32065331

31. Rukstalis D, Rashid P, Bogache WK, et al. 24-month durability after crossover to the prostatic urethral lift from randomised, blinded sham. *BJU Int.* Oct 2016; 118 Suppl 3: 14-22. PMID 27684483
32. Sievert KD, Schonthaler M, Berges R, et al. Minimally invasive prostatic urethral lift (PUL) efficacious in TURP candidates: a multicenter German evaluation after 2 years. *World J Urol.* Jul 2019; 37(7): 1353-1360. PMID 30283994
33. Jung JH, Reddy B, McCutcheon KA, et al. Prostatic urethral lift for the treatment of lower urinary tract symptoms in men with benign prostatic hyperplasia. *Cochrane Database Syst Rev.* May 25 2019; 5: CD012832. PMID 31128077
34. Gratzke C, Barber N, Speakman MJ, et al. Prostatic urethral lift vs transurethral resection of the prostate: 2-year results of the BPH6 prospective, multicentre, randomized study. *BJU Int.* May 2017; 119(5): 767-775. PMID 27862831
35. Miller LE, Chughtai B, Dornbier RA, et al. Surgical Reintervention Rate after Prostatic Urethral Lift: Systematic Review and MetaAnalysis Involving over 2,000 Patients. *J Urol.* May 12 2020: 101097JU0000000000001132. PMID 32396049
36. Roehrborn CG. Prostatic Urethral Lift: A Unique Minimally Invasive Surgical Treatment of Male Lower Urinary Tract Symptoms Secondary to Benign Prostatic Hyperplasia. *Urol Clin North Am.* Aug 2016; 43(3): 357-69. PMID 27476128
37. Roehrborn CG, Barkin J, Gange SN, et al. Five year results of the prospective randomized controlled prostatic urethral L.I.F.T. study. *Can J Urol.* Jun 2017; 24(3): 8802-8813. PMID 28646935
38. Rukstalis D, Grier D, Stroup SP, et al. Prostatic Urethral Lift (PUL) for obstructive median lobes: 12 month results of the MedLift Study. *Prostate Cancer Prostatic Dis.* Sep 2019; 22(3): 411-419. PMID 30542055
39. Shah BB, Tayon K, Madiraju S, et al. Prostatic Urethral Lift: Does Size Matter?. *J Endourol.* Jul 2018; 32(7): 635-638. PMID 29631445
40. Eure G, Gange S, Walter P, et al. Real-World Evidence of Prostatic Urethral Lift Confirms Pivotal Clinical Study Results: 2-Year Outcomes of a Retrospective Multicenter Study. *J Endourol.* Jul 2019; 33(7): 576-584. PMID 31115257
41. National Institute for Health and Care Excellence (NICE). Insertion of prostatic urethral lift implants to treat lower urinary tract symptoms secondary to benign prostatic hyperplasia [IPG475]. 2014; <https://www.nice.org.uk/guidance/ipg475/chapter/1recommendations>. Accessed September 2020.
42. National Institute for Health and Care Excellence (NICE). UroLift for treating lower urinary tract symptoms of benign prostatic hyperplasia [MTG26]. 2015; <https://www.nice.org.uk/guidance/mtg26>. Accessed September 2020.

*The articles reviewed in this research include those obtained in an Internet based literature search for relevant medical references through February 2021, the date the research was completed.*

### Joint BCBSA/BCN Medical Policy History

| Policy Effective Date | BCBSA Signature Date | BCN Signature Date | Comments   |
|-----------------------|----------------------|--------------------|--|
| 9/1/15                | 6/16/15              | 7/16/15            | Joint policy established   |
| 9/1/16                | 6/21/16              | 6/21/16            | Routine policy maintenance. Added Hayes rating, Added BCBSA policy information.  |
| 5/1/17                | 2/21/17              | 2/21/17            | Routine policy maintenance. Added the word "general" to clarify anesthesia on pg 5. Added article to references  |
| 3/1/18                | 12/12/17             | 12/12/17           | Rationale updated, added references 18-20. Removed reference to patient not being a surgical candidate.  |
| 3/1/19                | 12/11/18             |                    | Rationale reorganized, added reference 37 and 38. Added clinical input.  |
| 3/1/20                | 12/17/19             |                    | Rationale updated and reformatted. Reference # 6, 14, 32, and 37. No change in policy status.  |
| 1/1/21                | 10/20/20             |                    | Routine policy maintenance, added references 30, 32, 35 and 40. Added information on UroLift to the background section. No change in policy status.<br>1/6/21: Code C9769 removed as it does not apply to this policy. |
| 5/1/21                | 2/16/21              |                    | Section on Urolift was added to the background section. Code 0619T removed from policy as it does not apply. No change in policy status.   |

Next Review Date: 1<sup>st</sup> Qtr. 2022

## BLUE CARE NETWORK BENEFIT COVERAGE

### POLICY: PROSTATIC URETHRAL LIFT PROCEDURE FOR THE TREATMENT OF BPH

#### I. Coverage Determination:

|  |   |
|--|---|
| <b>Commercial HMO<br/>(includes Self-Funded<br/>groups unless otherwise<br/>specified)</b> | Covered; criteria apply                                     |
| <b>BCNA (Medicare<br/>Advantage)</b>   | See government section                                      |
| <b>BCN65 (Medicare<br/>Complementary)</b>  | Coinsurance covered if primary Medicare covers the service. |

#### II. Administrative Guidelines:

N/A