



Nonprofit corporations and independent licensees
of the Blue Cross and Blue Shield Association

Medical benefit drug policies are a source for BCBSM and BCN medical policy information only. These documents are not to be used to determine benefits or reimbursement. Please reference the appropriate certificate or contract for benefit information. This policy may be updated and therefore subject to change.

Effective Date: 10/12/2023

Tecvayli™ (teclistamab-cqyv)

HCPCS: J9380

Policy:

Requests must be supported by submission of chart notes and patient specific documentation.

- A. Coverage of the requested drug is provided when all the following are met:
 - a. FDA approved indication
 - b. FDA approved age
 - c. Prescribed by or in consultation with an oncologist
 - d. Treatment of patients with relapsed or refractory multiple myeloma after at least 4 prior lines of therapy
 - e. Patients must have been treated with all of the following:
 - i. An immunomodulatory agent
 - ii. A proteasome inhibitor
 - iii. An anti-CD38 antibody
 - f. Must have active disease defined by at least one of the following:
 - i. Serum M-protein greater or equal to 1.0 g/dL
 - ii. Urine M-protein greater or equal to 200 mg/24 h
 - iii. Serum free light chain (FLC) assay greater or equal to 10 mg/dL provided the baseline serum FLC ratio is abnormal
 - g. Patients must meet all of the following
 - i. ECOG performance status of 0 - 2
 - ii. No known central nervous system involvement with myeloma
 - iii. Alanine aminotransferase (ALT) and aspartate aminotransferase (AST) less than or equal to 3 times the upper limit of normal (ULN)
 - iv. Creatinine clearance greater than or equal to 40 mL/min
 - v. No detectable hepatitis B or C viral load
 - vi. No infection that is uncontrolled or requires IV or long-term oral antimicrobial therapy
 - vii. Left ventricular ejection fraction greater than or equal to 40%
 - viii. No stroke event within 6 months of therapy administration
 - ix. No pulmonary disease requiring oxygen dependence
 - x. No seizures within 6 months of therapy administration
 - xi. No active autoimmune disease except vitiligo, type 1 diabetes mellitus, or prior autoimmune thyroiditis

This policy and any information contained herein is the property of Blue Cross Blue Shield of Michigan and its subsidiaries, is strictly confidential, and its use is intended for the P&T committee, its members and BCBSM employees for the purpose of coverage determinations.

- h. Have not received prior treatment with any bispecific B-cell maturation antigen (BCMA)-directed CD3 T-cell engager therapy
 - i. Trial and failure, contraindication, OR intolerance to the preferred drugs as listed in BCBSM/BCN's utilization management medical drug list
- B. Quantity Limitations, Authorization Period and Renewal Criteria
- a. Quantity Limits: Align with FDA recommended dosing
 - b. Authorization Period: Aligns with FDA recommended or guideline supported treatment duration and provided for at least 60 days and up to 6 months at a time
 - c. Renewal Criteria: Treatment may be continued until disease progression or until unacceptable toxicity occurs

***Note: Coverage and approval duration may differ for Medicare Part B members based on any applicable criteria outlined in Local Coverage Determinations (LCD) or National Coverage Determinations (NCD) as determined by Center for Medicare and Medicaid Services (CMS). See the CMS website at <http://www.cms.hhs.gov/>. Determination of coverage of Part B drugs is based on medically accepted indications which have supported citations included or approved for inclusion determined by CMS approved compendia.

Background Information:

- Tecvayli is a bispecific BCMA-directed/CD3 T-cell engager indicated for the treatment of adult patients with relapsed or refractory multiple myeloma who have received at least four prior lines of therapy including a proteasome inhibitor, an immunomodulatory agent, and an anti-CD38 monoclonal antibody.
- Safety and efficacy were evaluated in the MajesTEC-1 study, a single-arm, multi-cohort, open-label, phase I/II trial of 110 patients with relapsed or refractory multiple myeloma who had previously received at least three prior therapies, including a proteasome inhibitor, an immunomodulatory agent, and an anti-CD38 monoclonal antibody. Subjects had not received prior BCMA-targeted therapy. About 78% of patients had received at least 4 prior lines of therapy, with a median number of 5. The study excluded patients who had stroke, seizure, allogeneic stem cell transplantation (SCT) within the past 6 months, an ECOG performance score of 2 or higher, known active CNS involvement or clinical signs of meningeal involvement of multiple myeloma, or active or documented history of autoimmune disease. The primary endpoint was overall response rate (ORR). The median time to first response was 1.2 months (range: 0.2 to 5.5 months). With a median follow-up of 7.4 months among responders, the estimated duration of response (DOR) rate was 90.6% (95% CI: 80.3%, 95.7%) at 6 months and 66.5% (95% CI: 38.8%, 83.9%) at 9 months. The ORR was 61.8% (95% CI: 52.1%, 70.9%).
- Disease should be measured/staged with PET-CT. Focal uptake in nodal and extranodal sites is considered involvement with lymphoma, including spleen, liver, bone, thyroid, and so on. A measurable node must have a longest diameter (LDi) greater than 1.5 cm. A measurable extranodal lesion should have an LDi greater than 1.0 cm. All other lesions (including nodal, extranodal, and assessable disease) should be followed as nonmeasured disease (eg, cutaneous, GI, bone, spleen, liver, kidneys, pleural or pericardial effusions, ascites).
- Due to the risk of cytokine release syndrome and neurological toxicities, the step-up doses of Tecvayli are required to be administered in an inpatient setting.
- Tecvayli has not been studied when given following prior treatment with Tecvayli or following any other bispecific B-cell maturation antigen (BCMA)-directed CD3 T-cell engager therapy.

This policy and any information contained herein is the property of Blue Cross Blue Shield of Michigan and its subsidiaries, is strictly confidential, and its use is intended for the P&T committee, its members and BCBSM employees for the purpose of coverage determinations.

References:

1. Tecvayli [prescribing information]. Horsham, PA: Janssen Biotech, Inc.; August 2023.
2. Moreau P, Garfall AL, van de Donk NWCJ, et al. Teclistamab in relapsed or refractory multiple myeloma. NEJM. 2022 Aug 11; 387 (6): 495 - 505.
3. Usmani SZ, Garfall AL, van de Donk NWCJ, et al. Teclistamab, a b-cell maturation antigen × CD3 bispecific antibody, in patients with relapsed or refractory multiple myeloma (MajesTEC-1): a multicentre, open-label, single-arm, phase 1 study. Lancet. 2021 Aug 21; 398 (10301): 665 - 74.
4. National Comprehensive Cancer Network. Multiple myeloma (Version 4.2023). 2023 August 25. Available at: https://www.nccn.org/professionals/physician_gls/pdf/myeloma.pdf. Accessed on September 13, 2023.

Policy History												
#	Date	Change Description										
1.2	Effective Date: 10/12/2023	Updated to align criteria with the other bispecific BCMA directed CD3 T-cell engager therapies including adding liver enzymes tests to be less than or equal to 3 times the ULN, creatinine clearance greater than or equal to 40 mL/min, left ventricular ejection fraction greater than 40%, no autoimmune disease, and removed the requirement ensuring there is no HIV infection										
1.1	Effective Date: 08/23/2023	UM medical management system update for BCNA, MAPPO, BCN, and BCBS <table border="1" data-bbox="483 808 1365 1018" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>Line of Business</th> <th>PA Required in Medical Management System (Yes/No)</th> </tr> </thead> <tbody> <tr> <td>BCBS</td> <td>Yes</td> </tr> <tr> <td>BCN</td> <td>Yes</td> </tr> <tr> <td>MAPPO</td> <td>Yes</td> </tr> <tr> <td>BCNA</td> <td>Yes</td> </tr> </tbody> </table>	Line of Business	PA Required in Medical Management System (Yes/No)	BCBS	Yes	BCN	Yes	MAPPO	Yes	BCNA	Yes
Line of Business	PA Required in Medical Management System (Yes/No)											
BCBS	Yes											
BCN	Yes											
MAPPO	Yes											
BCNA	Yes											
1.0	Effective Date: 12/01/2022	New policy										

* The prescribing information for a drug is subject to change. To ensure you are reading the most current information it is advised that you reference the most updated prescribing information by visiting the drug or manufacturer website or <http://dailymed.nlm.nih.gov/dailymed/index.cfm>.