

Nonprofit corporations and independent licensees of the Blue Cross and Blue Shield Association

Medical benefit drug policies are a source for BCBSM and BCN medical policy information only. These documents are not to be used to determine benefits or reimbursement. Please reference the appropriate certificate or contract for benefit information. This policy may be updated and therefore subject to change.

Effective Date: 12/12/2024

Adzynma (ADAMTS13, recombinant-krhn)

HCPCS: J7171

Policy:

Requests must be supported by submission of chart notes and patient specific documentation.

- A. Coverage of the requested drug is provided when all the following are met:
 - a. FDA approved indication
 - b. FDA approved age
 - c. Confirmation of diagnosis by serum assay showing less than 10% of normal ADAMTS13 enzyme activity and genetic testing showing a mutation in the ADAMTS13 gene
 - d. Must not be used in combination with any other therapy for the treatment of congenital thrombotic thrombocytopenic purpura (cTTP)
 - e. Trial and failure, contraindication, OR intolerance to the preferred drugs as listed in BCBSM/BCN's utilization management medical drug list.
- B. Quantity Limitations, Authorization Period and Renewal Criteria
 - a. Quantity Limit: Align with FDA recommended dosing
 - b. Initial Authorization Period: One year at a time
 - c. Renewal Criteria: Clinical documentation must be provided to confirm that current criteria are met and that the medication is providing clinical benefit

***Note: Coverage and approval duration may differ for Medicare Part B members based on any applicable criteria outlined in Local Coverage Determinations (LCD) or National Coverage Determinations (NCD) as determined by Center for Medicare and Medicaid Services (CMS). See the CMS website at http://www.cms.hhs.gov/. Determination of coverage of Part B drugs is based on medically accepted indications which have supported citations included or approved for inclusion determined by CMS approved compendia.

Background Information:

- Congenital thrombotic thrombocytopenic purpura (cTTP) is an autosomal recessive disorder caused by biallelic mutations on the ADAMTS13 gene. It is characterized by less than 10% ADAMTS13 enzyme activity and the accumulation of ultra-large von Willebrand factor (VWF). The buildup of ultra-large VWF multimers leads to uncontrolled platelet aggregation, platelet adhesion, and abnormal clotting in the small blood vessels. cTTP often causes seemingly mild and nonspecific symptoms such as lethargy, headache, loss of concentration, and abdominal discomfort. In many patients, symptoms are ongoing without a clear time of onset or resolution. Thrombocytopenia and microangiopathic hemolytic anemia can occur, but neurologic symptoms can occur without thrombocytopenia or anemia, and thrombocytopenia may occur without apparent hemolysis. Acute, severe exacerbations are uncommon but may be life-threatening without appropriate treatment. Exacerbations of the disease can be triggered during the neonatal period, pregnancy, infections, or surgeries although none of these has to occur for an acute attack to happen.
- The 2020 International Society on Thrombosis and Haemostasis (ISTH) good practice statements for the clinical care of patients with thrombotic thrombocytopenic purpura (TPP) state TTP should be considered in individuals presenting with thrombocytopenia and microangiopathic hemolytic anemia. cTTP should be suspected in individuals with any of the following presentations: severe neonatal hyperbilirubinemia, recurrent thrombocytopenia in a child or young adult, transient neurologic symptoms of stroke in a child or young adult, embolic stroke of undetermined source, or new onset TTP and absence of an ADAMTS13 inhibitor. Once suspected, patients should be tested for decreased ADAMTS13 enzyme activity, and if the enzymatic activity is less than 10% of normal without ADAMTS13 antibodies, genetic testing showing biallelic mutations in the ADAMTS13 gene will be confirmatory of cTPP.
- The standard of care for cTTP up until now has been plasma transfusions. Plasma transfusions allow for rapid replacement of ADAMTS13 during an acute attack and can be used prophylactically to minimize the risk of future attacks. The main complications of plasma infusion are allergic reactions. These may be treated or prevented using antihistamines. If allergic reactions are severe, other therapies, such as, a plasma-derived factor VIII concentrate that contains ADAMTS13 may be used.
- Adzynma provides an additional treatment option for cTTP. It is a human recombinant ADAMTS13 indicated for prophylactic or on demand enzyme replacement therapy (ERT) in adult and pediatric patients with cTTP.
- Safety and efficacy of Adzynma were evaluated in a Phase III, randomized, controlled, open-label, crossover, trial of 46 patients with cTTP. Patients were randomized to receive 6 months of treatment with either 40 IU/kg Adzynma or plasma-based therapies during the first period of the trial and then crossed over to receive the other treatment for 6 months during the second period. The mean annualized event rate of thrombocytopenia manifestations was 2.0 for patients receiving Adzynma (9 of 37 patients) compared to 4.44 in patients receiving plasma-based therapies (19 of 38 patients). Pharmacokinetic results showed patients (n = 23) receiving 40 IU/kg Adzynma IV achieved a four- to five-fold increase in ADAMTS13 enzyme activity after a single infusion compared to plasma-based therapies.
- Adzynma has not been studied and there is no data to support use in combination with other medications or treatment modalities used to treat cTTP.

References:

- 1. Adzynma [prescribing information]. Lexington, MA: Takeda Pharmaceuticals U.S.A., Inc.; June 2024.
- 2. Kremer Hovinga JA & George JN. Hereditary thrombotic thrombocytopenic purpura. NEJM. 2019 Oct 24; 381 (17): 1653 62.
- 3. Alwan F, Vendramin C, Liesner R, et al. Characterization and treatment of congenital thrombotic thrombocytopenic purpura. Blood. 2019 Apr 11; 133 (15): 1644 51.
- 4. Borogovac A & George JN. Stroke and myocardial infarction in hereditary thrombotic thrombocytopenic purpura: similarities to sickle cell anemia. Blood Adv. 2019; 3 (23): 3973 6.
- 5. Hamroun A, Prouteau C, & Provôt F. Hereditary thrombotic thrombocytopenic purpura. NEJM. 2020 Jan 23; 382 (4): 392 3.
- 6. Page EE, Kremer Hovinga JA, Terrell DR, et al. Thrombotic thrombocytopenic purpura: diagnostic criteria, clinical features, and long-term outcomes from 1995 through 2015. Blood Adv. 2017 Apr 6; 1 (10): 590 600.
- 7. Zheng XL, Vesely SK, Cataland SR, et al. ISTH guidelines for treatment of thrombotic thrombocytopenic purpura. J Thromb Haemost. 2020 Oct; 18: 2496 502.
- 8. Clinicaltrials.gov. A study of BAX 930 in children, teenagers, and adults born with thrombotic thrombocytopenic purpura (TTP) (NCT03393975). Available at: https://classic.clinicaltrials.gov/ct2/show/NCT03393975. Accessed on November 13, 2023.
- 9. Clinicaltrials.gov. A study of TAK-755 in participants with congenital thrombotic thrombocytopenic purpura (NCT04683003). Available at: https://classic.clinicaltrials.gov/ct2/show/study/NCT04683003. Accessed on November 13, 2023.

Policy	History			
#	Date	Change Description		
1.3	Effective Date: 12/12/2024	Annual review of criteria was performed, no changes were made		
1.2	Effective Date: 03/01/2024	UM medical management system update for MAPPO and BCNA		
		Line of Business	PA Required in Medical Management System (Yes/No)	
		BCBS	Yes	
		BCN	Yes	
		MAPPO	Yes	
		BCNA	Yes	
1.1	Effective Date: 12/14/2023	New policy		
1.0	Effective Date: 11/30/2023	UM medical management system update for BCBS and BCN		
		Line of Business	PA Required in Medical	
			Management System (Yes/No)	
		BCBS	Yes	
		BCN	Yes	
		MAPPO	No	

^{*} The prescribing information for a drug is subject to change. To ensure you are reading the most current information it is advised that you reference the most updated prescribing information by visiting the drug or manufacturer website or http://dailymed.nlm.nih.gov/dailymed/index.cfm.

Blue Cross Blue Shield/Blue Care Network of Michigan Medication Authorization Request Form



This form is to be used by participating physicians to obtain coverage for **drugs covered under the medical benefit**. For <u>commercial members only</u>, please complete this form and submit via fax to 1-877-325-5979. If you have any questions regarding this process, please contact BCBSM Provider Relations and Servicing or the Medical Drug Helpdesk at 1-800-437-3803 for assistance.

Nonprofit corporations and independent licensees of the Blue Cross and Blue Shield Association

Wedical Drug Ti	PATIENT INFORMATION	PHYSICIAN INFORMATION			
Name		Name			
ID Number		Specialty			
D.O.B.		Address			
Diagnosis		City /State/Zip			
Drug Name		Phone/Fax: P: () - F: () -			
Dose and Q	uantity	NPI			
Directions		Contact Person			
Date of Serv	rice(s)	Contact Person Phone / Ext.			
STEP 1: DI	SEASE STATE INFORMATION				
1. Is th	his request for: Initiation Continuation Date patient started therapy:				
2. Admi	inistered by patient or a medical professional? patient (self) health care professional (physician, nurse, etc.)				
3. Site o	Site of administration? Provider office/Home infusion Other:				
	☐ Hospital outpatient facility (go to #4)	Reason for Hospital Outpatient administration:			
	☐ Hospital inpatient facility for Car-T therap	y only (for example: Kymriah, Yescarta, or Tecartus) (go to #5)			
4. Pleas	ase specify location of administration if hospital outpatient infusion:				
5. Pleas	Please specify location of administration if hospital inpatient infusion:				
6. Pleas	6. Please provide the NPI number for the place of administration:				
	itiation AND Continuation of therapy: a. What is the patient's diagnosis?				
	b. What other medication has the patient received for their co	ndition? Please list			
	i. Please describe the response to previous therapies:				
	c. Will the patient be receiving any other treatment for the listed condition while on this medication? Please list:				
	d. Please list any labs values important for diagnosing or monitoring this patient's condition:				
8. Continuation of therapy: a. Has the patient progressed while on this medication? yes no b. How has the patient's condition changed while on this medication? Improved: Please describe: Stable: please describe: Worsened; Please describe: Other; Please describe:					
Chart notes are		supporting medical information necessary for our review (required)			
☐ Request for exp	Coverage will not be provided if the prescribing physicia edited review: I certify that applying the standard review time frame may seriously jeopar	In's signature and date are not reflected on this document. dize the life or health of the member or the member's ability to regain maximum function			
Physician's N Step 2:	ame Physician Signature ☐ Form Completely Filled Out	Date			
Checklist	☐ Provide chart notes	Attach test results			
Step 3: Submit	By Fax: BCBSM Specialty Pharmacy Mailbox 1-877-325-5979	By Mail: BCBSM Specialty Pharmacy Program P.O. Box 312320, Detroit, MI 48231-2320			

Confidentiality notice: This transmission contains confidential information belonging to the sender that is legally privileged. This information is intended only for use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution or action taken in reliance on the contents of this document is strictly prohibited. If you have received this in error, please notify the sender to arrange for the return of this document.