
Medical Policy



Nonprofit corporations and independent licensees
of the Blue Cross and Blue Shield Association

Joint Medical Policies are a source for BCBSM and BCN medical policy information only. These documents are not to be used to determine benefits or reimbursement. When Centers for Medicare and Medicaid (CMS) coverage rules are not fully developed, this medical policy may be used by BCBSM or BCN Medicare Advantage plans 42 CFR § 422.101 (b)(6).

Please reference the appropriate certificate or contract for benefit information. This policy may be updated and is therefore subject to change.

***Current Policy Effective Date: 7/1/26**
(See policy history boxes for previous effective dates)

Title: Remote Electrical Neuromodulation for Migraines (i.e., Nerivio)

Description

Nerivio is a wearable, non-pharmacologic device that uses remote electrical neuromodulation (REN) to treat and prevent migraines by stimulating nerves in the upper arm. The noninvasive device is controlled by a smartphone application. REN has been suggested as an alternative option for treatment of migraine, potentially reducing reliance on abortive or preventive medications and lowering the risk of medication overuse.

Medical Policy Statement

Remote electrical neuromodulation for the treatment and prevention of migraines is established when criteria are met.

Inclusionary and Exclusionary Guidelines

Inclusions:

Initial use of the FDA approved remote electrical neuromodulation device (i.e., Nerivio™) for the treatment and prevention of migraine in individuals who meet **ALL** the following four criteria:

- Are eight years of age or older; **and**
- Have a six-month history of headaches that meet International Classification of Headache Disorders (ICHD 3-beta) diagnostic criteria for migraine with or without aura; **and**
- **One** of the following conditions exists:

- Insufficient response, contraindication, or intolerance to 2 or more guideline-recommended headache medications (e.g., anticonvulsants, antihypertensives, antidepressants, CGRP inhibitors); **or**
- Pregnancy, breastfeeding, or planning to conceive; **or**
- At risk for or have a history of medication overuse headache; **or**
- At risk for drug-drug interactions with medications for comorbid conditions **or**
- Children that have a documented contraindication or intolerance to a medication **and**
- Meets **one** of the following indications:
 - Remote electrical neuromodulation (REN) is used to treat acute migraines; **or**
 - Remote electrical neuromodulation is used to prevent migraine for individuals with 6 to 24 headache days (defined as a calendar day with headache regardless of severity or duration) per 28-day period in each of the 3 months preceding use of the REN device.

Continued use of the FDA approved remote electrical neuromodulation device (i.e., Nerivio™) for the prevention or treatment of acute migraine in individuals when **BOTH** of the following are demonstrated:

- Compliance with use; **and**
- Clinical benefit (for example, fewer migraines or shorter duration of migraines).

Note: Each prescription, whether initial or refill, includes 18 treatments. For acute migraine management, individuals typically use 2 treatments per day, with a maximum of 6 per day. In contrast, preventative migraine treatment involves one treatment every other day. Based on these patterns, it is estimated that an individual would need approximately one prescription per year for acute migraines and two prescriptions per year for prevention.

Exclusions:

- Non-FDA approved REN devices
- Individuals with contraindications to use of REN devices
- All other indications for REN not meeting the above criteria
- Trigeminal nerve stimulation (e.g., Cefaly) devices for migraine treatment

Note: Contraindications include: Individuals with uncontrolled epilepsy and those with an active implantable medical device (e.g., pacemaker, hearing aid implant, or any implanted electronic device). Nerivio has not been evaluated in individuals with congestive heart failure, severe cardiac or cerebrovascular disease or patients under the age of 8 years.

CPT/HCPCS Level II Codes *(Note: The inclusion of a code in this list is not a guarantee of coverage. Please refer to the medical policy statement to determine the status of a given procedure.)*

Established codes:

A4540* E1399*

*When representing FDA approved devices

Other codes (investigational, not medically necessary, etc.):

E1399**

**When representing non-FDA approved devices

Note: Individual policy criteria determine the coverage status of the CPT/HCPCS code(s) on this policy. Codes listed in this policy may have different coverage positions (such as established or experimental/investigational) in other medical policies.

Background

The only currently available REN device (Nerivio™) cleared for use by the Food and Drug Administration (FDA) is worn on the upper arm and stimulates the peripheral nerves to induce conditioned pain modulation (CPM). The conditioned pain in the arm induced by the Nerivio REN device is purported to reduce the perceived migraine pain intensity.(9) Control of the device is accomplished through Bluetooth communication between the device and the individuals smartphone or tablet. For acute treatment, at onset of migraine or aura and no later than within 1 hour of onset, the user initiates use of the device through their mobile application. When used for preventive treatment, the device should be used every other day, controlled by the individual through their smartphone or tablet application. User-controlled stimulation intensity ranges from 0 to 100%, corresponding to 0 to 40 milliamperes (mA) of electrical current. Users are instructed to set the device to the strongest stimulation intensity that is just below their perceived pain level. The device provides stimulation for up to 45 minutes before turning off automatically. The Nerivio manufacturer indicates that the device can be used instead of or in addition to medication.

Migraine

Migraine is a neurologic disease characterized by recurrent moderate to severe headaches with associated symptoms that can include aura, photophobia, nausea, and/or vomiting.(1) Overall migraine prevalence in the United States is about 15% but varies according to population group.(2) Prevalence is higher in women (21%), among American Indian/Alaska Natives (22%), and among 18- to 44-year-olds (19%). Social determinants including low education level (18%), use of Medicaid (27%), high poverty level (23%), and being unemployed (22%) are also associated with higher rates of migraine.

Migraine is categorized as episodic or chronic depending on the frequency of attacks. Generally, episodic migraine is characterized by 14 or fewer headache days per month and chronic migraine is characterized by 15 or more headache days per month.(3) Specific International Classification of Headache Disorders (4) diagnostic criteria are as follows:

- Episodic migraine:
 1. Untreated or unsuccessfully treated headache lasting 4 to 72 hours.
 2. Headache has at least 2 of the following characteristics:
 - a. Unilateral location
 - b. Pulsating quality
 - c. Moderate or severe pain intensity
 - d. Aggravation by or causing avoidance of routine physical activity.
 3. At least 1 of the following during headache:
 - a. Nausea and/or vomiting.
 - b. Photophobia or phonophobia.
- Chronic migraine:
 1. Migraine-like or tension-type headache on 15 or more days per month for more than 3 months

2. At least 5 headache attacks without aura meet episodic migraine criteria 1 to 3, and/or at least 5 headache attacks with aura meet episodic migraine criteria 2 to 3.
3. On more than 8 days per month for more than 3 months, fulfilling any of the following criteria:
 - a. For migraine without aura, episodic migraine criteria 2 and 3
 - b. For migraine with aura, episodic migraine criteria 1 and 2
 - c. Believed by the patient to be migraine at onset and relieved by a triptan or ergot derivative.

Migraine attacks, whether due to episodic or chronic migraine, require acute management. The goal of acute treatment is to provide pain and symptom relief as quickly as possible while minimizing adverse effects, with the intent of timely return to normal function. Pharmacologic interventions for treatment of acute migraine vary according to migraine severity. First-line therapy for an acute episode of mild or moderate migraine includes oral non-steroidal anti-inflammatory drugs (NSAIDs) or acetaminophen. Moderate to severe migraine can be treated through the use of triptans or an NSAID-triptan combination. Antiemetics can be added for migraine accompanied by nausea or vomiting, though certain antiemetic medications used as monotherapy can also provide migraine relief. Other pharmacologic interventions used to treat acute migraine include calcitonin-gene related peptide antagonists, which can be used in patients with an insufficient response or contraindications to triptans, lasmiditan, and dihydroergotamine. Migraine can be managed at home, although acute migraine is a frequently cited reason for primary care and emergency department visits.(5) Regular use of pharmacologic interventions can result in medication overuse, which in turn could lead to rebound headache and increased risk of progression from episodic to chronic migraine.(4)

Many individuals who suffer from migraine may also benefit from preventive migraine therapy, including those with frequent or long-lasting migraines, migraine attacks that diminish quality of life or cause significant disability despite acute treatment, contraindications to or failure of acute therapies, and risk of medication overuse headache.(6,7,8) The main goals of preventive therapy are to reduce future attack frequency, severity, and duration, improve responsiveness to acute treatments, improve function and reduce disability, and prevent progression of episodic migraine to chronic migraine. For most adults with episodic migraines who may benefit from preventive therapy, initial therapy with an antiepileptic drug (divalproex sodium, sodium valproate, topiramate) or beta-blockers (metoprolol, propranolol, timolol) is recommended. Frovatriptan may be beneficial as initial therapy for prevention of menstrually associated migraine. Antidepressants (amitriptyline, venlafaxine), alternative beta-blockers (atenolol, nadolol), and additional triptans (naratriptan, zolmitriptan for menstrually associated migraine prevention) may be considered if initial therapy is unsuccessful. For preventive treatment of pediatric migraine, many children and adolescents who received placebo in clinical trials improved and most preventive medications were not superior to placebo. Possibly effective preventive treatment options for children and adolescents may include amitriptyline, topiramate, or propranolol.

Summary of Evidence

For individuals with acute migraine due to episodic or chronic migraine who receive REN, the evidence includes 2 RCTs and nonrandomized, uncontrolled studies. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Use of an

active REN device resulted in more patients with improved pain and symptoms at 2-hour follow-up compared with a sham device based on 2 small (N=212) RCTs with numerous relevance limitations. However, no significant between-group difference in functional disability or quality of life was noted in a post-hoc analysis of the pivotal RCT. Although, controlled studies in pediatric populations are lacking a non-randomized clinical trial in adolescents showed pain relief at 2 hours and improved functional disability relief. That data is consistent with emerging prospective real-world studies in younger children. Additional information was assessed from clinicians currently practicing at various academic medical centers. Respondents noted that all individuals with migraine may benefit from a nonpharmacologic option for either stand-alone or adjunctive use, particularly among those who have failed other options, who have contraindications or an intolerance to alternatives, who are at risk for or have a history of medication overuse headache, or who are at risk of drug-drug interactions. It was also noted that while some clinicians trial acute treatment with REN first, failure in the abortive setting does not preclude success with preventive use. Respondents additionally emphasize that the discreet nature of the REN device is ideal for use by children and adolescents in the school setting. Our assessment of the clinical input concluded that clinicians generally support use of REN for acute or preventive treatment, particularly in adolescent and pediatric populations where there are limited alternatives with evidence of efficacy or tolerable side effect profiles. The comprehensive evidentiary review which includes the clinical input is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who may benefit from preventive migraine therapy, including those with frequent or long-lasting episodic or chronic migraines, migraine attacks that diminish quality of life or cause significant disability despite acute treatment, contraindications to or failure of acute therapies, and risk of medication overuse headache, who receive REN, the evidence includes 1 RCT and 1 prospective, observational study. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Use of an active REN device resulted in more adults with decreased migraine days per month, regardless of episodic or chronic subtype, when used every other day for 8 weeks compared with a sham device based on 1 small (N=248) RCT with numerous relevance limitations. Prospective, observational data in 2 real world evidence studies using the device for acute treatment of migraine demonstrated a significant reduction in migraine headache days from baseline to months 2 and 3 with device use in adolescent patients. Based on the existing evidence, it is unclear how Nerivio would fit into the current migraine prevention pathway, although it could provide benefit for those who do not receive adequate benefit from pharmacologic first- or second-line therapies, or who may have a contraindication to pharmacologic therapies. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Rationale

Evidence reviews assess the clinical evidence to determine whether the use of a technology improves the net health outcome. Broadly defined, health outcomes are length of life, quality of life, and ability to function including benefits and harms. Every clinical condition has specific outcomes that are important to individuals and to managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health

outcome is a balance of benefits and harms. The following is a summary of the key literature to date.

ACUTE MIGRAINE DUE TO EPISODIC OR CHRONIC MIGRAINE

Clinical Context and Therapy Purpose

The purpose of remote electrical neuromodulation (REN) in individuals who have acute migraine attacks due to episodic or chronic migraine is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following therapies are currently being used to treat acute migraine due to episodic or chronic migraine: medical management or no treatment. A number of medications are used to treat migraine. First-line therapy for mild or moderate migraine includes oral non-steroidal anti-inflammatory drugs (NSAIDs) or acetaminophen. More severe migraine can be treated through the use of triptans or an NSAID-triptan combination through a variety of routes (e.g., oral, nasal spray or powder, subcutaneous). Antiemetics can be added for migraine accompanied by nausea or vomiting. Other pharmacologic interventions used to treat acute migraine include calcitonin-gene related peptide antagonists, which can be used in patients with an insufficient response or contraindications to triptans, lasmiditan, and dihydroergotamine.

The general outcomes of interest are: symptoms, functional outcomes, quality of life, and treatment-related morbidity. Specific important health outcomes include freedom from migraine pain and bothersome symptoms, restored function (e.g., return to normal activities), and patient-assessed global impression of treatment. Examples of relevant outcome measures appear in Table 1.

Follow-up over several hours is needed to monitor for treatment effects.

Table 1. Health Outcome Measures Relevant to Acute Migraine Attack

Outcome	Description
Pain free	No pain at defined assessment time (e.g., 2 hours)
Pain relief	Improvement of pain from moderate to severe at baseline to mild or none or pain scale improved at least 50% from baseline at defined assessment time (e.g., 2 hours)
Sustained pain free	No pain at initial assessment (e.g., 2 hours) and remains at follow-up assessment (e.g., 1 day) with no use of rescue medication or relapse (recurrence) within that time frame
Sustained pain relief	Improvement of pain from moderate to severe at baseline to mild or none or pain scale improved at least 50% from baseline at defined assessment time (e.g., 2 hours) and remains improved at follow-up assessment (e.g., 1 day) with no use of rescue medication or relapse (recurrence) within that time frame
Symptom relief	Improvement of most bothersome symptom(s) from moderate to severe at baseline to mild or none at defined assessment time (e.g., 2 hours)
Function relief	Improvement of function from moderate to severe at baseline to mild or none at defined assessment time (e.g., 2 hours)
Restored function	No restriction to perform work or usual activities at a defined assessment time (e.g., 2 hours)
Global impact of treatment	Patient assessment of functional disability and health-related quality of life using a Likert or other validated scale at a defined assessment time (e.g., 2 hours)
Global evaluation of treatment	Patient assessment of overall treatment effect (pain, symptom relief, adverse events) using a Likert or other validated scale at a defined assessment time (e.g., 2 hours)

Review of Evidence

Randomized Controlled Trials

Use of REN for the treatment of migraine has been assessed in 2 RCTs (Yarnitsky et al, 2017[17] and 2019[18]) comparing an active REN device (Nerivio Migra) with a sham device in patients with an acute migraine attack due to episodic migraine (Table 2).

A pilot crossover trial conducted by Yarnitsky et al (2017) included data from 71 (of 86 randomized) patients who received active or sham REN.(17) All patients were given an identical REN device that was preprogrammed to deliver in random order 4 active treatment sessions ranging from 80 to 120 hertz (Hz), corresponding to pulse widths of 50 to 200 milliseconds, and 1 sham session of 0.1 hertz (45 millisecond pulse width). Both active and sham treatments were programmed for a duration of 20 minutes each. Most patients were women (80%) in their mid-40s (mean age: 46 years), with a mean of 5 migraine attacks per month with a mean pain intensity of 8.8, corresponding to severe pain. Race and/or ethnicity were not reported. In the trial, treatment with active REN was more frequently associated with reduction in, and freedom from, migraine pain than sham REN at 2-hour follow-up (Table 3). When the device was programmed to deliver an active treatment session, it was most effective at reducing pain when used within 20 minutes of migraine onset. Treatment response to active REN diminished over time of initiation following migraine onset, and no active REN participants reported complete pain relief if the device was initiated more than 1 hour from onset. No adverse events were reported, though patients were more likely to rate their treatment perception of the active REN sessions as painful (11%) or unpleasant (28%) compared with sham REN sessions (1% painful; 13% unpleasant). Other outcomes were not reported in this study. Study limitations appear in Tables 4 and 5.

A second, larger (N=252) RCT was conducted by Yarnitsky et al in 2019 (Table 2).(18) The mean age of study participants was 43 years, 81% were female. Most participants were of White race (88%); 7% were Black and less than 1% were Asian. Time since migraine diagnosis was not reported; participants experienced a mean of 7 migraine days per month. Seventy-one percent of participants managed migraines with the use of acute medication, but important details about type and dosage were not provided. At baseline, 50% of participants reported that light sensitivity was their most bothersome symptom apart from migraine pain, followed by nausea (27%) and sound sensitivity (19%). After a 2 to 4-week run-in during which study participants kept a headache diary, participants were randomized to 4 to 6 weeks of at-home active or sham REN. The frequency was 100 to 120 Hz for the active device and less than 0.1 Hz for the sham device. The pulse width was 400 microseconds for the active device and ranged from 40 to 550 microseconds for the sham device, with the intent of mimicking a similar sensation as that delivered by the active device. At the time of randomization, participants were instructed on how to determine their optimal REN intensity, but this was unclearly defined as a threshold that was "perceptible not painful" (e.g., no specific measure of intensity was described) and no data on the actual intensities used during the study were reported. Participants were instructed to treat their migraine with the REN device as soon as possible following migraine onset, and no later than within 1 hour of onset. Participants who initiated device use more than 1 hour following onset were excluded from the outcome analyses. Study results are summarized in Table 3. Patients treated with active REN were more likely to report freedom from pain and pain relief at 2-hour follow-up, and sustained freedom from pain and pain relief at 48-hour follow-up compared with the sham REN group. There was no statistical between-group difference in the proportions of patients reporting freedom from their most bothersome symptom (MBS) at 2-hour follow-up, but a greater

proportion of active REN patients reported MBS relief at 2 hours relative to sham REN. Device-related adverse events were reported in 5% of active REN and 2% of sham REN participants ($p=.49$). At the conclusion of the study, participants were asked whether they believed they had received active or sham treatment as a measure of blinding. Half as many active participants correctly identified their device as did sham participants (23% in the active group vs. 50% in the sham group), although statistical analyses determined the treatment outcome differences between groups were not affected by participants perceived treatment group. Relevance and methodological limitations of the study are detailed in Tables 4 and 5. Notable limitations include an unclearly defined intended use population, a non-empirically determined optimum treatment regimen, and no assessment of functional or quality of life outcomes.

A post-hoc analysis of the Yarnitsky 2019 RCT retrospectively compared the effectiveness of acute migraine treatment with the Nerivio device with usual care (i.e., pharmacologic acute migraine management) used during the 2- to 4-week run-in phase of the trial.⁽¹⁹⁾ Pharmacologic treatment used during the run-in phase consisted of NSAIDs, acetaminophen (alone, or in combination with aspirin and caffeine) or triptans. In analysis of a subset of 99 trial participants, the rate of freedom from pain was similar for Nerivio (37.4% [37/99]) and usual care (26.3% [26/99]; $p=.099$) at 2-hour follow-up. Results were similar for achievement of pain relief (66.7% [66/99] vs. 52.5% [52/99]; $p=.034$). Randomized controlled trials directly comparing REN with pharmacologic management are needed to confirm these pain findings and to compare the effect of REN versus pharmacologic management on other outcomes.

Table 2. Summary of Key RCT Characteristics

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Yarnitsky et al (2017)	Israel	1	2016-2016	Adults (18 to 75 years) with ICHD-3 migraine 2 to 8 days/month with no preventive medication use 2 months prior to enrollment	n=86 Active REN device; 4/5 preprogrammed treatment sessions	NA; crossover trial Sham REN device; 1/5 preprogrammed treatment sessions
Yarnitsky et al (2019)	US, Israel	12	2017-2018	Adults (18 to 75 years) with ICHD-3 migraine 2 to 8 days/month but <12 days/month, with no or stable preventive medication use 2 months prior to enrollment	n=126 Active REN (Nerivio) device	n=126 Sham REN device

ICHD: International Classification of Headache Disorders; RCT: randomized controlled trial; REN: remote electrical neuromodulation.

Table 3. Summary of Key RCT Results

Study	Pain Free ¹ , 2 hours	Pain Relief ² , 2 hours	Sustained Pain Free, 48 hours	Sustained Pain Relief, 48 hours	MBS Free, 2 hours	MBS Relief ³ , 2 hours
Yarnitsky et al (2017)						
Active REN	44.1% (19/43)	76.7% (33/43)	NR	NR	NR	NR
Sham REN	5.9% (1/17)	23.5% (4/17)	NR	NR	NR	NR
p value	.005	.005	NR	NR	NR	NR

Yarnitsky et al (2019)						
Active REN	37.4% (37/99)	66.7% (66/99)	20.7% (18/87)	39.1% (34/87)	40.7% (33/81)	46.3% (44/95)
Sham REN	18.4% (19/103)	38.8% (40/103)	7.9% (7/89)	16.9% (15/89)	36.4% (32/88)	22.2% (22/99)
p value	.003	<.001	.014	.001	.55	.001

MBS: most bothersome symptom; NR: not reported; RCT: randomized controlled trial; REN: remote electrical neuromodulation.

¹ Change in headache severity from mild, moderate, or severe at baseline, to none.

² Change in headache severity from moderate, or severe at baseline, to none or mild, or a reduction in headache severity from mild to none.

³ Subjective (undefined) relief of most bothersome symptom.

Table 4. Study Relevance Limitations

Study	Population^a	Intervention^b	Comparator^c	Outcomes^d	Duration of Follow-up^e
Yarnitsky et al (2017)	1, 2 Intended use population is unclear (e.g., treatment naïve, those with contraindications to medication, or those who have failed pharmacologic treatment); time since migraine diagnosis and details about current migraine management regimen not reported	4 Predicate device not commercially marketed	2 Comparison versus an acute treatment with established efficacy would be preferred	1, 5 Functional and quality of life outcome measures not addressed	
Yarnitsky et al (2019)	1, 2 Intended use population is unclear (e.g., treatment naïve, those with contraindications to medication, or those who have failed pharmacologic treatment); time since migraine diagnosis and details about current migraine management regimen not reported	1, 5 Details about the mean timing of device initiation and mean, recommended or optimal device intensity (in mA) were not reported; a clinically relevant device intensity threshold has not been established; mean duration of treatments not reported	1, 2 Details and subgroup analysis on the effect of preventive medication use in 29% of active and 37% of sham participants were not reported; comparison versus an acute treatment with established efficacy would be preferred	1, 5 Functional and quality of life outcome measures not addressed	

mA: milliamperes.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5. Other.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 5. Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Yarnitsky et al (2017)	3 Method of allocation to active or sham treatment session not reported			1 No data reported for 17% (15/86) of enrolled participants	1 This was a pilot study; no sample size rationale or power calculations were reported	
Yarnitsky et al (2019)		4 23% of REN vs. 50% of sham participants correctly identified their treatment allocation; however, ad-hoc statistical analyses determined between-group treatment differences were not affected by perceived treatment group		1 19% (49/252) of randomized participants not accounted for in analysis described as modified intention to treat (ITT); ITT and per protocol efficacy analyses not reported; worst case scenario sensitivity analysis not performed for ITT population; sham response in preventive medication users not reported in Rapaport (2019)		

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4.

Inadequate control for selection bias; 5. Other.

^b Blinding key: 1. Participants or study staff not blinded; 2. Outcome assessors not blinded; 3. Outcome assessed by treating physician; 4. Other.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication; 4. Other.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials); 7. Other.

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference; 4. Other.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated; 5. Other.

Avoiding medication overuse has been postulated as a potential benefit of REN treatment of acute migraine. Marmura et al (2020)(20) reported the results of an observational 8-week open-label extension study following the double-blind phase of the Yarnitsky 2019 trial. The Marmura study compared within-subject data (N=117) from the trial run-in phase with data from the open-label phase, finding that a higher proportion of patients avoided medication use during the open-label phase (when the REN device was available for use; 89.7%) than in the run-in phase (when the REN device was not available for use; 15.4%). Although these results suggest that use of the REN device could result in less medication use and therefore reduce the risk of medication overuse, confirmatory studies designed to directly assess the role of REN in populations at risk of medication overuse are needed.

Nonrandomized Studies

Numerous nonrandomized, uncontrolled studies have been conducted examining the effectiveness of REN with the Nerivio device for acute migraine.(21-26) The most relevant studies are discussed below.

Three single-arm, open-label clinical trials of the Nerivio device were used to inform US Food and Drug Administration (FDA) approval for use in patients other than those with acute migraine due to episodic migraine (Table 6). This includes 2 studies (23,25) in patients with chronic migraine and 1 study (23) in adolescents. In the 2 studies (24,25) of patients with chronic migraine, the mean age was 42 and 44 years and was 15 years in the study of adolescents.(23) In all 3 studies most participants were female (60% to 83%) and of White race (86% to 100%). In the study by Hershey et al (2021)(23) conducted in adolescents, patients with episodic and chronic migraine were eligible for study inclusion. The studies reported on the effectiveness of the Nerivio device for acute migraine at 2 and 24 hours; study results are summarized in Table 7. The Nerivio device was associated with improvements in pain, symptoms, and function in all 3 studies. Adverse events related to the Nerivio device occurred in 1.0 to 2.0% of study participants across the 3 studies; no serious adverse events were reported in any of the studies. Results from these studies are limited due to their open-label study design, lack of control groups, and variable follow-up.

Table 6. Summary of Key Nonrandomized Clinical Trial Characteristics

Study	Country	Dates	Participants	Treatment	Follow-Up
Nierenburg et al 2020	US, Israel	2019-2020	N=42 adults (18 to 75 years) with ICHD-3 chronic migraine	REN (Nerivio)	24 hours
Grosberg et al 2021	US	2019-2020	N=126 adults (18 to 75 years) with ICHD-3 chronic migraine	REN (Nerivio)	24 hours
Hershey et al 2021	US	2019-2020	N=45 adolescents (12 to 17 years) with ICHD-3 migraine ≥3 attacks/month	REN (Nerivio)	24 hours

ICHHD: International Classification of Headache Disorders; REN: remote electrical neuromodulation.

Table 7. Summary of Key Nonrandomized Clinical Trial Results

Study	Pain Free, 2 hours	Pain Relief, 2 hours	Sustained Pain Free, 24 hours	Sustained Pain Relief, 24 hours	Symptom free, 2 hours	Functional improvement, 2 hours	Return to normal function, 2 hours
Nierenburg et al 2020	N=38	N=38	N=20	N=32	N=31	N=35	N=35
Proportion (n/N)	26.3% (10/38) ¹	73.7% (28/38) ¹	45.0% (9/20) ¹	84.4% (27/32) ¹	Nausea/vomiting: 58.3% (14/24) Photophobia: 35.5% (11/31) Phonophobia: 40.0% (10/25)	45.7% (16/35)	28.6% (10/35)
Grosberg et al 2021	N=99	N=99	NR	N=54	N=82	N=40	NR
Proportion (n/N)	19.2% (19/99) ²	54.5% (54/99) ³	NR	53.7% (29/54)	Nausea/vomiting: 40.8% (20/49) Photophobia: 36.6% (30/82) Phonophobia: 39.7% (129/73)	47.5% (19/40)	NR
Hershey et al 2021	N=39	N=39	N=11	N=22	N=31	N=33	NR

Proportion (n/N)	35.9% (14/39) ²	71.8% (28/39) ³	90.9% (10/11)	90.9% (20/22)	Nausea/vomiting: 54.5% (12/22)	69.7% (23/33)	NR
					Photophobia: 41.9% (13/31)		
					Phonophobia: 40.0% (10/25)		

NR: not reported.

¹ Pain free and pain relief for at least 50% of treated attacks.

² Change in headache severity from mild, moderate, or severe at baseline to none.

³ Change in headache severity from moderate or severe at baseline to none or mild; or a reduction in headache severity from mild to none.

A post-hoc analysis of the Hershey et al (2021) study, conducted in adolescents, compared the effect of Nerivio use (during the study phase) versus medication use (during the run-in phase) based on within-subject data.(22) Thirty-five adolescents who used medication during the 4-week run-in phase and who had Nerivio use data from the study phase were included in the post-hoc analysis. Nerivio users were more likely to report freedom from pain than medication users (p=.004) but there was no difference between Nerivio and medication in the proportions of patients who achieved pain relief (p=.225). Studies designed to directly compare the Nerivio device with medication are needed to adequately assess comparative effectiveness.

A real-world study (Ailani et al, 2021) sponsored by the Nerivio manufacturer collected data from 23,151 treatments from 5,805 Nerivio users between October 2019 and May 2021.(21) This study is unique in including data on use of the Nerivio device as monotherapy and in combination with medications. Nerivio users reported use of medications (over-the-counter, triptans, or other medications) in addition to the Nerivio device for about one-third of the treatment sessions. For use of Nerivio as monotherapy at 2-hour follow-up, the proportion of patients with freedom from pain, pain relief, return to normal function, and functional disability improvement was 20.3%, 55.6%, 24.9%, and 51.2%, respectively. When the Nerivio device was used in conjunction with medication, proportions ranged from 10.1% to 15.5% for freedom from pain, 38.5% to 51.3% for pain relief, 11.0% to 19.7% for return to normal function, and 39.8% to 49.6% for functional disability improvement, depending on the drug class used.

Pediatric Experience

The FDA 510(k) summary for the 2024 expanded approval of acute use of Nerivio in pediatric patients (ages 8-11) was based on a retrospective real-world analysis of 293 children, aged 6-11 at their first use of the device.(14) Median patient age was 11 (73.7% female). Safety data were primarily collected via self-reported customer service complaints, which indicated no adverse events. Effectiveness was assessed in individuals who completed voluntary pre- and post-treatment surveys. Available efficacy data was only available in 18 participants, of which consistent headache relief was reported by 72.2% (13/18), consistent freedom from headache by 83.3% (15/18), and consistent functional disability freedom by 38.9% (7/18). Additional controlled studies are required to confirm efficacy in this population. Individuals in this population used REN as a standalone treatment, with over-the-counter medications, and with prescribed headache medications, in 45.4%, 34.4%, and 20.9% of treatments, respectively.(28)

In 2024, Hershey et al published the results of a survey of 332 students aged 7-17 (80.4% female).(29) After being prescribed REN, the percentage of students reporting having their

headaches treated at school increased by 11.5%. The most common reasons given for preferring REN treatment at school are the ability to avoid going to the nurse's office (42.5%) and the ability to treat discreetly (39.2%). Barriers to treatment included concerns about standing out (42.2%), and permission to use a smartphone in class to control the REN device (22.9%).

Coverage with Evidence Development Study

In 2025, Synowiec and coworkers published the outcomes of a real-world postmarketing coverage with evidence development study conducted in partnership with Highmark Inc.(30) Members aged 12-75 years who were diagnosed with migraine were prescribed REN. Eligibility criteria included: (1) at least 1 standard-of-care acute migraine therapy had failed for 1 or more of the following reasons: contraindication, lack of sufficient efficacy, or intolerability of adverse effects; (2) were at risk of drug-drug interaction with other medications; (3) were a pregnant woman, woman trying to conceive, or breastfeeding woman; (4) had chronic migraine and were at risk of or diagnosed with medication overuse headache (MOH); or (5) were younger than 18 years. Efficacy was assessed by change in Migraine Disability Assessment (MIDAS) score from baseline to 3 months of treatment and by prospective pain and disability reports 2 hours post treatment. Device utilization was assessed through prescription fulfillment rates. A total of 381 patients (mean age, 40.5; 91.1% female) participated in the study. Change in MIDAS score was calculated for all participants who answered the questionnaire twice (n=116), indicating a statistically and clinically meaningful improvement of -12.1 point (p =.014). Pain relief, pain freedom, functional disability relief, and functional disability freedom were reported in 77.8%, 33.3%, 70.6%, and 50.0% of participants, respectively Three minor adverse events were reported and patients used a mean of 4.0 (SD, 3.1) devices annually. Generalizability of outcomes is limited due to lack of a control group and high degree of missing data (70%).

Section Summary: Acute Migraine due to Episodic or Chronic Migraine

Evidence from 2 small RCTs found REN with the Nerivio device was more effective than a sham device for measures of pain and symptom relief at 2-hours post-treatment. Patients treated with the Nerivio device were also more likely than those treated with a sham device to report 48-hour freedom from pain and pain relief based on 1 RCT. The remaining evidence from post-hoc and nonrandomized studies suggests that REN with the Nerivio device may provide improvements in acute pain and symptomatology. There is sufficient evidence to determine that the technology may result in an improvement in net health outcomes.

PREVENTION OF MIGRAINE

Clinical Context and Therapy Purpose

The purpose of REN as preventive therapy in individuals who have acute migraine attacks due to episodic or chronic migraine is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following therapies are currently being used to prevent acute migraine due to episodic or chronic migraine: medical management or no treatment. A number of medications are used as prevention for migraine. For most adults with episodic migraines who may benefit from preventive therapy, initial therapy with an antiepileptic drug (divalproex sodium, sodium valproate, topiramate) or beta-blockers (metoprolol, propranolol, timolol) is recommended. Frovatriptan may be beneficial as initial therapy for prevention of menstrually associated

migraine. Antidepressants (amitriptyline, venlafaxine), alternative beta-blockers (atenolol, nadolol), and additional triptans (naratriptan, zolmitriptan for menstrually associated migraine prevention) may be considered if initial therapy is unsuccessful. For preventive treatment of pediatric migraine, many children and adolescents who received placebo in clinical trials improved and most preventive medications were not superior to placebo. Possibly effective preventive treatment options for children and adolescents may include amitriptyline, topiramate, or propranolol.

The general outcomes of interest are: symptoms, functional outcomes, quality of life, and treatment-related morbidity. Specific important health outcomes include reduction of future attack frequency, severity, and duration, improved responsiveness to acute treatments, improved function and reduced disability, and prevention of progression of episodic migraine to chronic migraine.

Follow-up over several days to months is needed to monitor for preventive treatment effects.

Review of Evidence

Randomized Controlled Trials

Use of REN for the prevention of migraine has been assessed in 1 double-blind, multicenter RCT by Tepper et al (2023), comparing an active REN device (Nerivio) used every other day with a sham device in adult patients with at least a 6-month history of headaches that meet the International Classification of Headache Disorders, third edition (ICHD-3) and 6 to 24 headache days per 28-day period in the past 3 months.⁽³¹⁾ Included participants either did not use preventive medicine or were on a stable dose of a single migraine preventive medication during the 2 months before enrollment and throughout the study. Prior to initiation of REN, all patients participated in a 4-week baseline phase, where they were instructed to continue their regular medications when needed, and document daily reports, regardless of if they had a headache that day or not, to rate symptoms using a 4-point scale. Symptoms that were collected included pain, functional disability, presence or absence of nausea and/or vomiting, photophobia, and phonophobia, and acute medication usage.

To be eligible for the intervention phase, individuals had to have had 6 to 24 headache days during the 28-day baseline period, with at least 4 headache days fulfilling ICHD-3 criteria for migraine and had at least 80% compliance on completing their daily record of symptoms. The intervention phase was 8 weeks long and included participants were randomized 1:1 to active REN or sham REN. The active and sham devices were visually identical, so staff and participants were blinded to their randomized group. Participants were directed to complete a full 45-minute treatment with REN every other day and to complete a daily diary. If acute treatment was needed, participants were instructed to use their usual acute treatments. The primary outcome was the mean change in number of migraine days per month in the 4-week baseline phase compared to the last 4 weeks of treatment phase (weeks 9 through 12). Overall, patients treated with the active REN device had statistically fewer migraine days during the intervention period compared to baseline compared to those treated with sham. This was also demonstrated in subanalyses based on episodic or chronic migraines. Of the participants, 40.8% used a preventive medication in combination with REN. Half of the medication users were on first-line preventive medications (e.g., amitriptyline, topiramate), while the other half were on second line agents (e.g., anti-calcitonin gene-related peptide monoclonal antibodies, onabotulinumtoxin A, gepants). There were 2 non-device-related

serious adverse events in the REN arm. There was a single device-related adverse event in the sham group and no device-related adverse events in the active group. There were no differences in quality of life questionnaires or Headache Impact Tests, a tool used to capture the impact of headache on functional health and well-being, between groups at any time period. These results are limited by the 8-week duration, shorter than the recommended 12-week duration by the International Headache Society guidelines for neuromodulation devices and lack of medical history reporting previous preventive medications used by participants. Tables 8 and 9 describe the key characteristics and results of the RCT. No significant difference between REN and sham groups was noted for mean change in the Headache Impact Test Short Form (HIT-6) or Migraine Specific Quality of Life Questionnaire - Function Domain (MSQ). Tables 10 and 11 describe notable limitations.

Table 8. Summary of Key RCT Characteristics

Study	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Tepper et al (2023)	US	15	2021-2022	Adults (18 to 75 years) with ICHD-3 migraine at least 4 days/month in baseline period with no preventive medication use or stable medication use 2 months prior to enrollment; 85% female, mean age of 41.7 years; and ratio of episodic to chronic patients was 47.6%: 52.4%.	n=128 (ITT); 95 (mITT) Active REN device, use every other day	n=120 (ITT); 84 (mITT) Sham REN device

ICHHD: International Classification of Headache Disorders; ITT: intention to treat; mITT: modified intention to treat; RCT: randomized controlled trial; REN: remote electrical neuromodulation.

Table 9. Summary of Key RCT Results

Study	Overall mean change in migraine days/month ¹	Mean change in migraine days/month: Episodic subgroup ¹	Mean change in migraine days/month: Chronic subgroup ¹	Mean change in moderate/severe headache days	Mean change in number of headache days	Percentage of patients achieving at least 50% reduction from baseline in headache days
Tepper et al (2023)						
n	n=95 active REN; n=84 sham REN	n=45 active REN; n=42 sham REN	n=50 active REN; n=42 sham REN	n=95 active REN; n=84 sham REN	n=95 active REN; n=84 sham REN	n=95 active REN; n=84 sham REN
Active REN	-4.0±4.0	-3.2±3.4	-4.7±4.4	-3.8±3.9	-4.5±4.1	26.3%
Sham REN	-1.3±4.0	-1.0±3.6	-1.6±4.4	-2.2±3.6	-1.8±4.6	11.9%
Difference versus sham (95% CI); p value	-2.7 (-3.9 to -1.5); <.001	2.3 (NR); .003	3.0 (NR); .001	-1.6 (-2.7 to -0.5); .005	-2.7 (-3.9 to -1.5); <.001	NR; NR; .015

CI: confidence interval; NR: not reported; RCT: randomized controlled trial; REN: remote electrical neuromodulation.

¹ Change in migraine days from baseline (weeks 1 through 4) compared to last 4 weeks (weeks 9 through 12)

Table 10. Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
Tepper et al (2023)	1, 2 Intended use population is unclear (e.g., treatment naïve, those with contraindications to medication, or those who have failed pharmacologic treatment); time since migraine diagnosis and details about current migraine management regimen and previously failed treatments not reported	1 A clinically relevant device intensity threshold has not been established	2 Comparison versus specific pharmacologic preventive treatments with established efficacy would be preferred if attempting to establish first-line use		3. 8-week duration is less than the recommended 12-week duration by IHS guidelines for neuromodulation devices

IHS: International Headache Society.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5. Other.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 11. Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Tepper et al (2023)				2 LOCF imputation methodology for full ITT set not prespecified and generally does not meet MCAR assumptions; worst case sensitivity analysis not reported; 28% of data missing in mITT analysis		

ITT: intention to treat; LOCF: last observation carried forward; MCAR: missing completely at random; mITT: modified ITT.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias; 5. Other.

^b Blinding key: 1. Participants or study staff not blinded; 2. Outcome assessors not blinded; 3. Outcome assessed by treating physician; 4. Other.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication; 4. Other.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials); 7. Other.

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference; 4. Other.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated; 5. Other.

Nonrandomized Studies

Prospective, real-world data collected and analyzed by the manufacturer on the use of Nerivio in adolescents was summarized in the FDA approval packet for the indication of Nerivio in

migraine prevention in adolescents and adults.(13) The data were collected from adolescents who used the device for acute migraine treatment, but use was equivalent to the suggested preventive use (10 times per month or higher). Prospective data were collected through the Nerivio app between January 2021 and November 2022. Eligible adolescent patients used Nerivio on at least 10 days in their first 28-day month of using the device and used the device on at least 3 days in each of the 2 subsequent months. The goal of analysis was to assess the mean reduction in migraine headache days from the first month of use to the second and third month of use. In total, 61 patients (mean age, 15.7 ± 1.3 years, 87% female) were eligible for analysis. Investigators found significant month-to-month reduction in migraine headache days from 15 days (standard error [SE], 0.6) in month 1, to 10.6 days (SE, 0.8) in month 2 ($p < .0001$), and 8.7 days (SE, 0.7) in month 3 ($p < .0001$), demonstrating substantial reduction from baseline during months 2 and 3 of device use. Data limitations include a lack of comparator and no description of medications or alternative interventions patients were additionally using.

A prospective, real-world evidence analysis investigated whether the use of Nerivio in adolescents who have frequently utilized the REN wearable device had reduced mean monthly migraine treatment days (MMTD) compared to baseline.(32) Patients ($N=83$) were 15.9 ± 1.3 years of age (mean $\pm$ SD) and were evaluated for a 3 month period. There was a statistically significant monthly reduction in MMTD (a reduction of $3.6 [\pm 4.8]$ MMTD) from the first month to the second month of consecutive use ($p < .001$). In the third month of treatment, there was a further reduction of $1.6 (\pm 4.1)$ MMTD ($p < .001$), for a cumulative total reduction of $5.2 (\pm 4.8)$ MMTD throughout the first 3 months of consecutive treatment.

The FDA 510(k) summary for the expanded approval of preventive use of Nerivio in pediatric patients (ages 8-11) was based on a retrospective real-world analysis of 293 children, aged 6-11.(14) Preventive use of the device was assumed by analyzing patients whose frequency of use in month one was suggestive of preventative treatments. No specific data on proportion of patients in whom preventive use was assumed or efficacy outcomes in the assumed preventive use population were reported in the 510(k) summary. Well-defined, controlled studies are required to confirm benefit in this population.

Section Summary: Prevention of Migraine

Evidence from a small RCT found REN with the Nerivio device was more effective than a sham device for decreasing migraine days per month, regardless of episodic or chronic subgroup, when used every other day for 8 weeks. Patients treated with the Nerivio device were also more likely than those treated with sham to have reduced moderate to severe headache days, reduced headache days in general, and at least a 50% reduction from their baseline in overall headache days. Approximately half of patients included in this study were also taking preventive pharmacologic therapy. There were no differences in quality of life or functional health patient-reported outcomes between groups at any time point. Prospective, observational data in 2 real world evidence studies using the device for acute treatment of migraine demonstrated a significant reduction in migraine headache days from baseline to months 2 and 3 with device use in adolescent patients. Based on the existing evidence, it is unclear how Nerivio would fit into the current migraine prevention pathway, although it could provide benefit for those who do not receive adequate benefit from pharmacologic first- or second-line therapies, or who may have a contraindication to pharmacologic therapies. The specific intended use and associated empirically-documented recommended regimen(s) based on test results must be specified in order to adequately evaluate net health benefit.

Supplemental Information

Clinical Input

BCBSA questioned if the use of remote electrical neuromodulation (i.e., Nerivio) would provide clinically meaningful improvements in the overall net health outcomes for individuals 8 years of age and older, who experience episodic or chronic migraines, with or without aura, and are treated for acute migraine or migraine prophylaxis. Clinical input was received (2025) from 4 respondents, including 1 physician-level respondent identified by the American Academy of Neurology (AAN), and 3 physician-level responses identified by various academic medical centers. In addition, an informal response was provided by the American Academy of Pediatrics Section on Neurology (AAP). The majority of respondents supported the use of remote electrical neuromodulation (REN) for migraines and indicated it is consistent with accepted medical practice. Respondents noted that all individuals with migraines may benefit from a nonpharmacologic therapy as either a stand-alone option or as an adjunctive to pharmacology; Particularly among those who (1) have failed other options, (2) have contraindications or an intolerance to alternatives, (3) are at risk for or have a history of medication overuse, or (4) are at risk of drug-drug interactions. Additionally, clinicians support first-line use of REN for acute or preventive treatment, particularly in adolescent and pediatric populations where there are: (1) limited alternatives or (2) suitable side effect profiles. Respondents emphasize that the discreet nature of the REN device is ideal for use by children and adolescents in the school setting.

American Academy of Neurology/American Headache Society

A 2012 joint guideline by the American Academy of Neurology (AAN) and the American Headache Society (AHS) on pharmacologic treatment for episodic migraine prevention in adults was published prior to the approval of Nerivio in the US and did not address the use of remote electrical neuromodulation (REN) or other nonpharmacologic treatments.(7) Similarly, 2019 joint guidelines issued by AAN and AHS on the treatment of acute migraine (33) and prevention of migraine (8) in children and adolescents did not address the use of REN or other nonpharmacologic treatments.

American Headache Society

In 2021, the American Headache Society (AHS) issued guidance on the integration of new migraine treatments, including REN, into clinical practice.(4) The AHS addressed the use of neuromodulatory devices as a group that included electrical trigeminal nerve stimulation, noninvasive vagus nerve stimulation, single-pulse transcranial magnetic stimulation and REN; no guidance specific to REN use was issued.

- The AHS determined that initiation of a neuromodulatory device is appropriate when all of the following criteria are met:
 - Prescribed/recommended by a licensed clinician.
 - Patient is at least 18 years of age (the guidance noted that 3 devices, including REN, are approved for use in patients age 12 to 17 years)
 - Diagnosis of International Classification of Headache Disorders (ICHD) migraine with aura, migraine without aura, or chronic migraine
 - Either of the following:
 - Contraindications to or inability to tolerate triptans.

- Inadequate response to 2 or more oral triptans, as determined by EITHER of the following:
 - Validated acute treatment patient-reported outcome questionnaire (Migraine Treatment Optimization Questionnaire, Patient Perception of Migraine Questionnaire-Revised, Functional Impairment Scale, Patient Global Impression of Change)
 - Clinician attestation.

Department of Veterans Affairs/Department of Defense

The U.S Department of Veterans Affairs/Department of Defense (VA/DoD) 2023 guidelines for the management of headache state that "there is insufficient evidence to recommend for or against any form of neuromodulation for the treatment and/or prevention of migraine"; examples of neuromodulation treatments mentioned include remote electrical neurostimulation.(34)

Ongoing and Unpublished Clinical Trials

Some currently ongoing trials that might influence this review are listed in Table 12.

Table 12. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing			
<i>Unpublished</i>			
NCT05940870 ^a	A Prospective, Open-label, Post-marketing Observational Study Assessing the Safety and Efficacy of Nerivio for Migraine Prevention in Real-world Environment	300	Aug 2024 (completed)

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

Regulatory Status

In May 2019, Nerivio Migra™ (Theranica Bio-Electronics Ltd.) was granted a de novo classification by the FDA (class II, special controls, product code: QGT).(10) This new classification applied to this device and substantially equivalent devices of this generic type. Nerivio Migra was initially cleared for treatment of acute migraine in adults who do not have chronic migraine.

In October, 2020, Nerivio was cleared for marketing by the FDA through the 510(k) process (K201824). FDA determined that this device was substantially equivalent to Nerivio Migra for use in adults.(11) The device name changed to just "Nerivio" and the exclusion of chronic migraine patients was removed. The Nerivio device can provide more treatments than the predicate Nerivio Migra (12 treatments vs. 8 treatments) and has a longer shelf life (24 months vs. 9 months). In January, 2021, the Nerivio device was cleared for use in patients aged 12 to 17 years.(12) In February 2023, Nerivio's indication was expanded to include preventive treatment of migraine with or without aura in individuals 12 years and age or older and was cleared for marketing through the 510(k) process (K223169).(13) In May 2025, the Nerivio and rechargeable Nerivio Infinity devices were cleared for marketing (K241756) with an expanded

indication for acute and/or preventive treatment of migraine with or without aura in patients 8 years and older.(14)

Government Regulations

National:

There is no national coverage determination.

Local:

There is no local coverage determination.

(The above Medicare information is current as of the review date for this policy. However, the coverage issues and policies maintained by the Centers for Medicare & Medicare Services [CMS, formerly HCFA] are updated and/or revised periodically. Therefore, the most current CMS information may not be contained in this document. For the most current information, the reader should contact an official Medicare source.)

Related Policies

- Oxygen Therapy for Cluster Headache
 - Sphenopalatine Ganglion Block for Headache
 - Surgical Deactivation of Headache Trigger Sites
 - Transcutaneous Electrical Modulation Pain Reprocessing (Scrambler Therapy)
 - Vagus Nerve Stimulation
 - Transcutaneous Electrical Nerve Stimulation (e.g., Cefaly®, eTNS, Axon) and Transcutaneous Afferent Patterned Stimulation (e.g., Cala ONE™, Cala TRIO™, CALA kIQ,)
-

References

1. VanderPluym JH, Halker Singh RB, Urtecho M, et al. Acute Treatments for Episodic Migraine in Adults: A Systematic Review and Meta-analysis. JAMA. Jun 15 2021; 325(23): 2357-2369. PMID 34128998
2. Burch R, Rizzoli P, Loder E. The prevalence and impact of migraine and severe headache in the United States: Updated age, sex, and socioeconomic-specific estimates from government health surveys. Headache. Jan 2021; 61(1): 60-68. PMID 33349955
3. Singh RBH, VanderPluym JH, Morrow AS, et al. Acute Treatments for Episodic Migraine. Rockville (MD): Agency for Healthcare Research and Quality (US); December 2020. Accessed April 5, 2022.
4. Ailani J, Burch RC, Robbins MS. The American Headache Society Consensus Statement: Update on integrating new migraine treatments into clinical practice. Headache. Jul 2021; 61(7): 1021-1039. PMID 34160823
5. Burch RC, Loder S, Loder E, et al. The prevalence and burden of migraine and severe headache in the United States: updated statistics from government health surveillance studies. Headache. Jan 2015; 55(1): 21-34. PMID 25600719
6. Silberstein SD. Practice parameter: evidence-based guidelines for migraine headache (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology. Neurology. Sep 26 2000; 55(6): 754-62. PMID 10993991

7. Silberstein SD, Holland S, Freitag F, et al. Evidence-based guideline update: pharmacologic treatment for episodic migraine prevention in adults: report of the Quality Standards Subcommittee of the American Academy of Neurology and the American Headache Society. *Neurology*. Apr 24 2012; 78(17): 1337-45. PMID 22529202
8. Oskoui M, Pringsheim T, Billingshurst L, et al. Practice guideline update summary: Pharmacologic treatment for pediatric migraine prevention: Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology and the American Headache Society. *Neurology*. Sep 10 2019; 93(11): 500-509. PMID 31413170
9. Nierenburg H, Stark-Inbar A. Nerivio (R) remote electrical neuromodulation for acute treatment of chronic migraine. *Pain Manag*. Apr 2022; 12(3): 267-281. PMID 34538078
10. U.S. Food and Drug Administration. De Novo Classification Request for Nerivio Migma. https://www.accessdata.fda.gov/cdrh_docs/reviews/DEN180059.pdf. Accessed August 9, 2025.
11. U.S. Food and Drug Administration. 501(k) Summary: Theranica Bio-Electronics LTDs Nerivio. https://www.accessdata.fda.gov/cdrh_docs/pdf18/DEN180059.pdf. Accessed August 9, 2025.
12. U.S. Food and Drug Administration. 510(k) Summary: Nerivio Approval in Adolescents. https://www.accessdata.fda.gov/cdrh_docs/pdf20/K203181.pdf. Accessed August 8, 2025.
13. U.S. Food and Drug Administration. 510(k) Summary (K223169): Nerivio Approval for Preventative Treatment. https://www.accessdata.fda.gov/cdrh_docs/pdf22/K223169.pdf. Accessed August 5, 2024.
14. Food and Drug Administration (FDA). 510(k) Summary. Theranica Bio-Electronics LTD.'s Nerivio Infinity and Nerivio (K241756). October 8, 2024. https://www.accessdata.fda.gov/cdrh_docs/pdf24/K241756.pdf. Accessed August 18, 2025.
15. Tassorelli C, Diener HC, Silberstein SD, et al. Guidelines of the International Headache Society for clinical trials with neuromodulation devices for the treatment of migraine. *Cephalalgia*. Oct 2021; 41(11-12): 1135-1151. PMID 33990161
16. Diener HC, Tassorelli C, Dodick DW, et al. Guidelines of the International Headache Society for controlled trials of acute treatment of migraine attacks in adults: Fourth edition. *Cephalalgia*. May 2019; 39(6): 687-710. PMID 30806518
17. Yarnitsky D, Volokh L, Ironi A, et al. Nonpainful remote electrical stimulation alleviates episodic migraine pain. *Neurology*. Mar 28 2017; 88(13): 1250-1255. PMID 28251920
18. Yarnitsky D, Dodick DW, Grosberg BM, et al. Remote Electrical Neuromodulation (REN) Relieves Acute Migraine: A Randomized, Double-Blind, Placebo-Controlled, Multicenter Trial. *Headache*. Sep 2019; 59(8): 1240-1252. PMID 31074005
19. Rapoport AM, Bonner JH, Lin T, et al. Remote electrical neuromodulation (REN) in the acute treatment of migraine: a comparison with usual care and acute migraine medications. *J Headache Pain*. Jul 22 2019; 20(1): 83. PMID 31331265
20. Marmura MJ, Lin T, Harris D, et al. Incorporating Remote Electrical Neuromodulation (REN) Into Usual Care Reduces Acute Migraine Medication Use: An Open-Label Extension Study. *Front Neurol*. 2020; 11: 226. PMID 32318014
21. Ailani J, Rabany L, Tamir S, et al. Real-World Analysis of Remote Electrical Neuromodulation (REN) for the Acute Treatment of Migraine. *Front Pain Res (Lausanne)*. 2021; 2: 753736. PMID 35295483

22. Hershey AD, Irwin S, Rabany L, et al. Comparison of Remote Electrical Neuromodulation and Standard-Care Medications for Acute Treatment of Migraine in Adolescents: A Post Hoc Analysis. *Pain Med.* Apr 08 2022; 23(4): 815-820. PMID 34185084
23. Hershey AD, Lin T, Gruper Y, et al. Remote electrical neuromodulation for acute treatment of migraine in adolescents. *Headache.* Feb 2021; 61(2): 310-317. PMID 33349920
24. Nierenburg H, Vieira JR, Lev N, et al. Remote Electrical Neuromodulation for the Acute Treatment of Migraine in Patients with Chronic Migraine: An Open-Label Pilot Study. *Pain Ther.* Dec 2020; 9(2): 531-543. PMID 32648205
25. Tepper SJ, Lin T, Montal T, et al. Real-world Experience with Remote Electrical Neuromodulation in the Acute Treatment of Migraine. *Pain Med.* Dec 25 2020; 21(12): 3522-3529. PMID 32935848
26. Grosberg B, Rabany L, Lin T, et al. Safety and efficacy of remote electrical neuromodulation for the acute treatment of chronic migraine: an open-label study. *Pain Rep.* Nov-Dec 2021; 6(4): e966. PMID 34667919
27. Nierenburg H, Rabany L, Lin T, et al. Remote Electrical Neuromodulation (REN) for the Acute Treatment of Menstrual Migraine: a Retrospective Survey Study of Effectiveness and Tolerability. *Pain Ther.* Dec 2021; 10(2): 1245-1253. PMID 34138449
28. Werner K, Gerson T, Stark-Inbar A, et al. Acute treatment of migraine in children aged 6-11: Real-world analysis of remote electrical neuromodulation (REN). *Annals of the Child Neurology Society.* May 21,2024; 2(2): 135-145. <https://doi.org/10.1002/cns3.20073>.
29. Hershey AD, Shmueli S, Stark-Inbar A, et al. Patterns, Barriers, and Preferences of Treating Migraine Within the School Setting: A Survey Study of Students. *Children (Basel).* Oct 25 2024; 11(11). PMID39594861
30. Synowiec A, Stark-Inbar A, Santamaria DD, et al. Coverage with evidence development study shows benefits in patients with migraine treated with remote electrical neuromodulation. *Am J Manag Care.* Apr 08 2025. PMID 40834199
31. Tepper SJ, Rabany L, Cowan RP, et al. Remote electrical neuromodulation for migraine prevention: A double-blind, randomized, placebo-controlled clinical trial. *Headache.* Mar 2023; 63(3): 377-389. PMID 36704988
32. Monteith TS, Stark-Inbar A, Shmueli S, et al. Remote electrical neuromodulation (REN) wearable device for adolescents with migraine: a real-world study of high-frequency abortive treatment suggests preventive effects. *Front Pain Res (Lausanne).* 2023; 4: 1247313. PMID 38028429.
33. Oskoui M, Pringsheim T, Holler-Managan Y, et al. Practice guideline update summary: Acute treatment of migraine in children and adolescents: Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology and the American Headache Society. *Headache.* Sep 2019; 59(8): 1158-1173. PMID 31529481
34. Management of Headache Work Group. VA/DoD Clinical Practice Guideline. Washington, DC. September 2023. <https://www.healthquality.va.gov/guidelines/pain/headache/VA-DoD-CPG-Headache-Full-CPG.pdf>. Accessed August 13, 2024.
35. Hayes, Inc. Evolving Evidence Review. Nerivio (Theranica Bio-Electronics Ltd.) for treatment of acute migraine episodes. <https://evidence.hayesinc.com>. Published July 23, 2021. Accessed January 23, 2025.

The articles reviewed in this research include those obtained in an Internet based literature search for relevant medical references through March 11, 2026, the date the research was completed.

Joint BCBSM/BCN Medical Policy History

Policy Effective Date	BCBSM Signature Date	BCN Signature Date	Comments
11/1/22	8/16/22		Joint policy established
11/1/23	8/15/23		Routine maintenance (slp) Vendor Managed: Northwood
7/1/24	4/16/24		• Routine maintenance (slp) • Vendor managed: Northwood • Review added for prevention of migraine based on expansion of FDA-approved indications – EI • MPS clarified to include acute, chronic, episodic, and prevention of migraines • Code update: K1023 deleted and replaced with A4540 (nomenclature remains same)
7/1/25	4/15/25		• Routine maintenance (slp) • Vendor managed: Northwood
7/1/26	4/21/26		• Routine maintenance (slp/fr) • Vendor managed: Northwood • Policy stance changed to EST with FDA approved indications, based on SME input.

Next Review Date: 2nd Qtr, 2027

BLUE CARE NETWORK BENEFIT COVERAGE
POLICY: REMOTE ELECTRICAL NEUROMODULATION FOR MIGRAINES

I. Coverage Determination:

Commercial HMO (includes Self-Funded groups unless otherwise specified)	Not Covered
BCNA (Medicare Advantage)	Refer to the Medicare information under the Government Regulations section of this policy.
BCN65 (Medicare Complementary)	Coinsurance covered if primary Medicare covers the service.

II. Administrative Guidelines:

- The member's contract must be active at the time the service is rendered.
- Coverage is based on each member's certificate and is not guaranteed. Please consult the individual member's certificate for details. Additional information regarding coverage or benefits may also be obtained through customer or provider inquiry services at BCN.
- The service must be authorized by the member's PCP except for Self-Referral Option (SRO) members seeking Tier 2 coverage.
- Services must be performed by a BCN-contracted provider, if available, except for Self-Referral Option (SRO) members seeking Tier 2 coverage.
- Payment is based on BCN payment rules, individual certificate and certificate riders.
- Appropriate copayments will apply. Refer to certificate and applicable riders for detailed information.
- CPT - HCPCS codes are used for descriptive purposes only and are not a guarantee of coverage.
- Duplicate (back-up) equipment is not a covered benefit.