
Medical Policy



Nonprofit corporations and independent licensees
of the Blue Cross and Blue Shield Association

Joint Medical Policies are a source for BCBSM and BCN medical policy information only. These documents are not to be used to determine benefits or reimbursement. When Centers for Medicare and Medicaid (CMS) coverage rules are not fully developed, this medical policy may be used by BCBSM or BCN Medicare Advantage plans 42 CFR § 422.101 (b)(6). Please reference the appropriate certificate or contract for benefit information. This policy may be updated and is therefore subject to change.

***Current Policy Effective Date: 11/1/26**
(See policy history boxes for previous effective dates)

Title: Nontraditional Biomarkers in Risk Assessment and Management of Cardiovascular Disease

Description

Various lipid and non-lipid biomarkers have been proposed as potential risk markers for cardiovascular disease (CVD). Biomarkers discussed in this policy include apolipoprotein B (apo B), apolipoprotein A1 (apo A1), apolipoprotein E (apo E), B-type natriuretic peptide, cystatin C, fibrinogen, high-density lipoprotein (HDL) subclass, leptin, low-density lipoprotein (LDL) subclass, lipoprotein (a), high sensitivity C-reactive protein (hs-CRP), and genetic biomarkers. These biomarkers have been studied as alternatives or additions to standard lipid panels for risk stratification in CVD or as treatment targets for lipid-lowering therapy.

Medical Policy Statement

Measurement of apolipoprotein B is established when criteria are met.

Measurement of high sensitivity C-reactive protein (hs-CRP) is established when criteria are met.

Measurement of lipoprotein(a) is established when criteria are met.

Testing of other nontraditional biomarkers, including but not limited to apolipoprotein A1, apolipoprotein E or APOE genotypes, brain natriuretic peptide (BNP), cystatin C, fibrinogen, leptin, LDL subclass, and HDL subclass is experimental/investigational. There is insufficient evidence to determine that the technology results in an improvement in net health outcomes.

Inclusionary and Exclusionary Guidelines

Inclusions:

Apolipoprotein B (apo B)

Measurement of apolipoprotein B in individuals receiving lipid lowering therapy who have achieved low-density lipoprotein cholesterol and/or non-high-density lipoprotein cholesterol treatment goals to guide further therapeutic intensification, when one or more of the following is present:

- Established atherosclerotic cardiovascular disease; **OR**
- Cardiovascular-kidney-metabolic syndrome: **OR**
- Type 2 diabetes; **OR**
- Elevated triglycerides

High-sensitivity C-reactive protein (hs-CRP)

High-sensitivity C-reactive protein (hs-CRP) testing for cardiovascular risk assessment in adults without known atherosclerotic cardiovascular disease to guide decisions regarding therapeutic initiation when **all** the following are met:

- estimated 10-year atherosclerotic cardiovascular disease borderline risk*: AND
- Standard cardiovascular risk assessment has been completed and the decision regarding initiation of lipid-lowering therapy remains uncertain: AND
- Testing is performed when the individual is clinically stable and not during acute infection or other acute inflammatory illness and identifiable secondary causes of CRP elevation have been considered

*Several tools are available to calculate 10-year risk of atherosclerotic cardiovascular disease (ASCVD).

The following are examples:

ACC/AHA: https://tools.acc.org/ldl/ascvd_risk_estimator/index.html#!/calculate/estimator/

ClinCalc.com: <https://clincalc.com/Cardiology/ASCVD/PooledCohort.aspx>

Lipoprotein(a) (Lp[a])

- A single measurement of Lipoprotein (a) in adults for ASCVD risk assessment.

Exclusions:

- Measurement of apolipoprotein B, high-sensitivity C-reactive protein, and lipoprotein(a) is excluded for all other indications
- Laboratory testing of other nontraditional biomarkers to assess cardiovascular risk, including but not limited to apolipoprotein AI, apolipoprotein E or *APOE* genotypes, brain natriuretic peptide (BNP), cystatin C, fibrinogen, leptin, LDL subclass, and HDL subclass

CPT/HCPCS Level II Codes

Established codes:

81401	81405	81406	82172	83695
84999	86141			

Other codes (investigational, not medically necessary, etc.):

82397	82610	82664	83520	83700	83701
83704	83718	83719	83721	83722	83880
84181	85384	85385	0052U		

Notes:

- *Code(s) may not be covered by all contracts or certificates. Please consult customer or provider inquiry resources at BCBSM or BCN to verify coverage.*
- *Individual policy criteria determine the coverage status of the CPT/HCPCS code(s) on this policy. Codes listed in this policy may have different coverage positions (such as established or experimental/investigational) in other medical policies.*

Background

LOW-DENSITY LIPOPROTEINS AND CARDIOVASCULAR DISEASE

Low-density lipoproteins (LDLs) have been identified as the major atherogenic lipoproteins and have long been identified by the National Cholesterol Education Project as the primary target of cholesterol-lowering therapy. An LDL particle consists of a surface coat composed of phospholipids, free cholesterol, and apolipoproteins surrounding an inner lipid core composed of cholesterol ester and triglycerides. Traditional lipid risk factors such as LDL-cholesterol (LDL-C), while predictive on a population basis, are weaker markers of risk on an individual basis. Only a minority of individuals with elevated LDL and cholesterol levels will develop clinical disease, and up to 50% of cases of coronary artery disease (CAD) occur in those with "normal" levels of total cholesterol and LDL-C. Thus, there is considerable potential to improve the accuracy of current cardiovascular risk prediction models. To assess CVD risk, especially when LDL alone does not fully explain risk (e.g., in metabolic syndrome, diabetes mellitus, or residual risk despite statin use), other lipid biomarkers are useful.

Lipid Markers

Apolipoprotein B

Apolipoprotein (Apo) B is the major protein moiety of all lipoproteins except for high-density lipoprotein (HDL). The most abundant form of apo B, large B or B₁₀₀, constitutes the apo B found in LDL and very-low density LDL. Because LDL and very-low density LDL each contain 1 molecule of apo B, the measurement of apo B reflects the total number of these atherogenic particles, 90% of which are LDL. Because LDL particles can vary in size and in cholesterol content, for a given concentration of LDL-C, there can be a wide variety in size and numbers of LDL particles. Thus, it has been postulated that apo B is a better measure of the atherogenic potential of serum LDL than the LDL concentration.

Apolipoprotein AI

High-density lipoprotein contains 2 associated apolipoproteins: (i.e., apo AI, apo AII). High-density lipoprotein particles can also be classified by whether they contain apo AI only or they contain both apo AI and apo-AII. Because all HDL particles contain apo AI, this lipid marker can be used as an approximation for HDL number, similar to the way apo B has been proposed as an approximation of the LDL number.

Direct measurement of apo AI concentrations has been proposed as more accurate than the traditional use of HDL level in the evaluation of cardioprotective, or “good,” cholesterol. In addition, the ratio of apo B/apo AI has been proposed as a superior measure of the ratio of proatherogenic (i.e., “bad”) cholesterol to anti-atherogenic (i.e., “good”) cholesterol.

Apolipoprotein E

Apolipoprotein E is the primary apolipoprotein found in very-low density LDLs and chylomicrons. Apolipoprotein E is the primary binding protein for LDL receptors in the liver and is thought to play an important role in lipid metabolism. *APOE* gene is polymorphic, consisting of 3 epsilon alleles (e2, e3, and e4) that code for 3 protein isoforms, known as E2, E3, and E4, which differ from one another by 1 amino acid. These molecules mediate lipid metabolism through their different interactions with LDL receptors. The genotype of apo E alleles can be assessed by gene amplification techniques, or the *APOE* phenotype can be assessed by measuring plasma levels of apo E.

It has been proposed that various *APOE* genotypes are more atherogenic than others and that *APOE* measurement may provide information on risk of CAD beyond traditional risk factor measurement. It has also been proposed that the *APOE* genotype may be useful in the selection of specific components of lipid-lowering therapy, such as drug selection. In the major lipid-lowering intervention trials, including trials of statin therapy, there is considerable variability in response to therapy that cannot be explained by factors such as compliance. The *APOE* genotype may be a factor that determines an individual's degree of response to interventions such as statin therapy.

High-Density Lipoprotein Subclass

High-density lipoprotein particles exhibit considerable heterogeneity, and it has been proposed that various subclasses of HDL may have a greater role in protection from atherosclerosis. Particles of HDL can be characterized based on size or density and/or on apolipoprotein composition. Using size or density, HDL can be classified into HDL₂, the larger, less dense particles that may have the greatest degree of cardioprotection, and HDL₃, which are smaller, denser particles.

An alternative to measuring the concentration of subclasses of HDL (e.g., HDL₂ and HDL₃) is direct measurement of HDL particle size and/or number. Particle size can be measured by nuclear magnetic resonance (NMR) spectroscopy or by gradient-gel electrophoresis. High-density lipoprotein particle numbers can be measured by NMR spectroscopy. Several commercial labs offer these measurements of HDL particle size and number. Measurement of apo AI has used HDL particle number as a surrogate, based on the premise that each HDL particle contains a single apo AI molecule.

Low-Density Lipoprotein Subclass

Two main subclass patterns of LDL, called A and B, have been described. In subclass pattern A, particles have a diameter larger than 25 nm and are less dense, while in subclass pattern B, particles have a diameter less than 25 nm and a higher density. Subclass pattern B is a commonly inherited disorder associated with a more atherogenic lipoprotein profile, also termed “atherogenic dyslipidemia.” In addition to small, dense LDL, this pattern includes elevated levels of triglycerides, elevated levels of apo B, and low levels of HDL. This lipid profile is commonly seen in type 2 diabetes and is one component of the “metabolic syndrome,” defined by the Third Report of the Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III; ATP III) to also include high normal blood

pressure, insulin resistance, increased levels of inflammatory markers such as C-reactive protein, and a prothrombotic state. The presence of the metabolic syndrome is considered by Adult Treatment Panel III to be a substantial risk-enhancing factor for CAD.

Low-density lipoprotein size has also been proposed as a potentially useful measure of treatment response. Lipid-lowering treatment decreases total LDL and may also induce a shift in the type of LDL, from smaller, dense particles to larger particles. It has been proposed that this shift in lipid profile may be beneficial in reducing risk for CAD independent of the total LDL level. Also, some drugs may cause a greater shift in lipid profile than others. Niacin and/or fibrates may cause a greater shift from small to large LDL size than statins. Therefore, measurement of LDL size may potentially play a role in drug selection or may be useful in deciding to use a combination of drugs rather than a statin alone.

In addition to the size of LDL particles, interest has been shown in assessing the concentration of LDL particles as a distinct cardiac risk factor. For example, the commonly performed test for LDL-C is not a direct measure of LDL but, chosen for its convenience, measures the amount of cholesterol incorporated into LDL particles. Because LDL particles carry much of the cholesterol in the bloodstream, the concentration of cholesterol in LDL correlates reasonably well with the number of LDL particles when examined in large populations. However, for an individual patient, the LDL level may not reflect the number of particles due to varying levels of cholesterol in different sized particles. It is proposed that the discrepancy between the number of LDL particles and the serum level of LDL represents a significant source of unrecognized atherogenic risk. The size and number of particles are interrelated. For example, all LDL particles can invade the arterial wall and initiate atherosclerosis. However, small, dense particles are thought to be more atherogenic than larger particles. Therefore, for patients with elevated numbers of LDL particles, cardiac risk may be further enhanced when the particles are smaller versus larger.

Lipoprotein (a)

Lipoprotein (a) (Lp[a]) is a lipid-rich particle similar to LDL. The major apolipoprotein associated with LDL is Apo B; in Lp(a), however, there is an additional apo A covalently linked to apo B. The apo A molecule is structurally similar to plasminogen, suggesting that Lp(a) may contribute to the thrombotic and atherogenic basis of CVD. Levels of Lp(a) are relatively stable in individuals over time but vary up to 1,000-fold between individuals, presumably on a genetic basis. The similarity between Lp(a) and fibrinogen has stimulated intense interest in Lp(a) as a link between atherosclerosis and thrombosis. In addition, approximately 20% of patients with CAD have elevated Lp(a) levels. Therefore, it has been proposed that levels of Lp(a) may be an independent risk factor for CAD.

Other non-lipid markers have been identified as having an association with CVD, including high sensitivity C-reactive protein, B-type natriuretic peptide, cystatin C, fibrinogen, and leptin. These biomarkers may have a predictive role in identifying CVD risk or in targeting therapy.

Non-Lipid Markers

B-type or Brain Natriuretic Peptide

Brain natriuretic peptide (BNP, also called B-type natriuretic peptide) is an amino acid polypeptide secreted primarily by the ventricles of the heart when the pressure to the cardiac muscles increases or there is myocardial ischemia. Elevations in BNP levels reflect deterioration in cardiac loading levels and may predict adverse events. Brain natriuretic peptide has been studied as a biomarker for managing heart failure and predicting cardiovascular and heart failure risk.

Cystatin C

Cystatin C is a small serine protease inhibitor protein secreted from all functional cells in the body. It has primarily been used as a biomarker of kidney function. Cystatin C has also been studied to determine whether it may serve as a biomarker for predicting cardiovascular risk. Cystatin C is encoded by the *CST3* gene.

Fibrinogen

Fibrinogen is a circulating clotting factor and precursor of fibrin. It is important in platelet aggregation and a determinant of blood viscosity. Fibrinogen levels have been shown to be associated with future risk of CVD and all-cause mortality.

High-Sensitivity C-Reactive Protein (hs-CRP)

High sensitivity C-reactive protein (hs-CRP) is produced in response to inflammation in the body. It is a systemic marker of inflammation that has been extensively studied and identified as an independent risk factor for coronary artery disease (CAD). A high sensitivity CRP assay can detect low levels of chronic inflammation associated with heart disease.

Leptin

Leptin is a protein secreted by fat cells that has been found to be elevated in heart disease. Leptin has been studied to determine if it has any relation to the development of CVD.

Summary of Evidence

For individuals who are asymptomatic with risk of cardiovascular disease (CVD) who receive nontraditional cardiac biomarker testing (e.g., apolipoprotein B [apo B], apolipoprotein AI [apo AI], apolipoprotein E [apo E], B-type natriuretic peptide, cystatin C, fibrinogen, leptin, subclasses of low-density lipoprotein [LDL], high-density lipoprotein [HDL], and lipoprotein [a]), the evidence includes systematic reviews, meta-analyses, and large, prospective cohort studies. Relevant outcomes are overall survival, other test performance measures, change in disease status, morbid events, and medication use. The evidence from cohort studies and meta-analyses of these studies has suggested that some of these markers are associated with increased cardiovascular risk and may provide incremental accuracy in risk prediction. The evidence is sufficient to determine that apo B, hs-CRP and Lp(a) testing results in an improvement in the net health outcome.

For individuals with hyperlipidemia managed with lipid-lowering therapy who receive nontraditional cardiac biomarker testing (e.g., apo B, apo AI, apo E, apo E, HDL subclass, LDL subclass, Lp[a], BNP, cystatin C, fibrinogen, leptin), the evidence includes analyses of the intervention arm(s) of lipid-lowering medication trials. Relevant outcomes are overall survival,

change in disease status, morbid events, and medication use. In particular, apo B, apo AI, and apo E have been evaluated as markers of lipid-lowering treatment success, and evidence from the intervention arms of several RCTs has suggested that these markers are associated with treatment success. However, there is no direct evidence that using markers other than LDL and HDL as lipid-lowering treatment targets leads to improved health outcomes. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Rationale

Evidence reviews assess the clinical evidence to determine whether the use of a technology improves the net health outcome. Broadly defined, health outcomes are length of life, quality of life, and ability to function including benefits and harms. Every clinical condition has specific outcomes that are important to individuals and to managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms. The following is a summary of the key literature to date.

NONTRADITIONAL BIOMARKERS

A large body of literature has accumulated on the utility of nontraditional lipid risk factors in the prediction of future cardiac events. The evidence reviewed for this policy consists of systematic reviews, meta-analyses and large, prospective cohort studies that have evaluated the association between these lipid markers and cardiovascular outcomes. A smaller amount of literature is available on the utility of these markers as a measure of treatment response. Data on treatment responses are taken from randomized controlled trials (RCTs) that use 1 or more novel lipid markers as a target of lipid-lowering therapy.

The Adult Treatment Panel III (ATP III) guidelines noted that, to determine their clinical significance, the emerging risk factors should be evaluated against the following criteria:(1)

- Significant predictive power that is independent of other major risk factors
- A relatively high prevalence in the population (justifying routine measurement in risk assessment)
- Laboratory or clinical measurement must be widely available, well standardized, inexpensive, have accepted population reference values, and be relatively stable biologically

It is preferable, but not necessary, that modification of the risk factor in clinical trials will have shown reduction in risk.

ASYMPTOMATIC INDIVIDUALS WITH RISK OF CARDIOVASCULAR DISEASE

Clinical Context and Test Purpose

The purpose of nontraditional cardiac biomarker testing in individuals who are asymptomatic with risk of CVD is to inform a decision whether nontraditional cardiac biomarker testing improves CVD diagnosis and treatment decisions.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

APOLIPOPROTEIN B

A publication from a consensus conference of the American Diabetes Association and the American College of Cardiology Foundation (2008) includes specific recommendations for incorporating apo B testing into clinical care for high-risk patients. It is recommended that for patients with metabolic syndrome who are being treated with statins, both LDL-C and apo B should be used as treatment targets, with an apo B target of less than 90 mg/dL. This consensus statement also commented on the use of LDL particle number in patients with cardiometabolic risk and the limitations of the clinical utility of NMR measurement of LDL particle number or size, including lack of widespread availability. Also mentioned was the need for more independent data confirming the accuracy of the method and whether its predictive power is consistent across various patient populations.(2)

The American Association for Clinical Chemistry (2013) conducted a review of 25 clinical studies containing 85 outcomes for which both biomarkers apolipoprotein B (apo B) and LDL particle number (LDL-P) were determined. In 21 of 25 (84.0%) studies, both apo B and LDL-P were significant for at least 1 outcome. Neither was significant for any outcome in only 1 study (4.0%). In 50 of 85 comparisons (58.8%), both apo B and LDL-P had statistically significant associations with the clinical outcome, whereas in 17 comparisons (20.0%) neither was significantly associated with the outcome. In 18 comparisons (21.1%) there was discordance between apo B and LDL-P. The AACC concluded “in most studies, both apo B and LDL-P were comparable in association with clinical outcomes. The biomarkers were nearly equivalent in their ability to assess risk for CVD and both have consistently been shown to be stronger risk factors than LDL-C. We support the adoption of apo B and/or LDL-P as indicators of atherogenic particle numbers into CVD risk screening and treatment guidelines. Currently, in the opinion of this Working Group on Best Practices, apo B appears to be the preferable biomarker for guideline adoption because of its availability, scalability, standardization, and relatively low cost.”(3)

The American Association of Clinical Endocrinologists (as of 2020), the American Association of Clinical Endocrinology (AACE) and the American College of Endocrinology (ACE) (2017) published clinical practice guidelines for the diagnosis and treatment of dyslipidemia and prevention of atherosclerosis. The AACE states “when the triglyceride concentration is greater than 150 mg/dL or the HDL-C concentration is less than 40 mg/dL, the apo B or the apo B to apo AI ratio may be particularly useful in assessing residual risk in patients at risk for CAD (even when LDL-C levels are controlled); this includes patients with established CAD, type 2 diabetes, or the insulin resistance syndrome who are at high risk for CAD. AACE therefore recommends apo B testing in such patients.” The guidelines support the use of apolipoprotein B or LDL particle number measurements to refine efforts to achieve effective LDL-C lowering, provide screening recommendations for persons of different ages, and identify special issues for pediatric patients. AACE recommends the goals set by the American College of Cardiology and the American Diabetes Association that optimal apo B levels for patients at risk of CAD, including those with diabetes, are less than 90 mg/dL, while patients with established CAD or diabetes plus 1 or more additional risk factor should have an apo B goal of less than 80 mg/dL.(4)

The National Lipid Association (2021) published a scientific statement for lipid measurements in the management of cardiovascular disease. Their recommendations (evidence rating “B” or higher) regarding apolipoprotein B are as follows:(5)

- Apolipoprotein B measurement may be reasonable for initial evaluation (Strength: IIb. Evidence: B-NR)
- Apolipoprotein B measurement is reasonable on lipid therapy (Strength: IIa. Evidence: BNR)
- Apolipoprotein B measurement is recommended to facilitate diagnosis of Familial Dysbetalipoproteinemia and Familial Combined Hyperlipidemia (Strength: IIb. Evidence: B-NR)

The Emerging Risk Factors Collaboration (2012) published a patient-level meta-analysis of 37 prospective cohort studies enrolling 154,544 individuals.(6) Risk prediction was examined for a variety of traditional and non-traditional lipid markers. For apolipoprotein B (apo B), evidence from 26 studies on 139,581 individuals reported that apo B was an independent risk factor for cardiovascular events, with an adjusted hazard ratio of 1.24 (95% confidence interval [CI], 1.19 to 1.29).

The Canadian Cardiovascular Society updated guidelines (2021) recommend that for any individual with triglycerides greater than 1.5 mmol/L, either non-high-density lipoprotein cholesterol (non-HDL-C) or apolipoprotein B are the preferred lipid parameter for screening, rather than low-density lipoprotein cholesterol.(7)

Guidelines from the British Columbia Medical Services Commission (2008, updated 2014) state that those with initial risk assessments that reflect elevated lipids may be followed up with a non-HDL-C or apolipoprotein B assessment in 4-6 months.(8)

A cohort study (2005) of 15,632 participants from the Women’s Health Initiative provided similar information in women.(9) In this analysis, the hazard ratio for developing coronary heart disease in the highest versus the lowest quintiles was greater for apo B (2.50; 95% CI, 1.68 to 3.72) compared to LDL-C (1.62; 95% CI, 1.17 to 2.25), after adjustment for traditional cardiovascular risk factors.

Guidelines from the European Society of Cardiology and European Atherosclerosis Society (2019) states that apolipoprotein B (apo B) was a more accurate marker of cardiovascular risk than low-density lipoprotein cholesterol (LDL-C) and non-high-density lipoprotein cholesterol.(10)

The Copenhagen City Heart Study (2007) prospectively evaluated a cohort of 9231 asymptomatic persons from the Danish general population followed for 8 years.(11) Subjects with total apo B levels in the top one-third (top tertile) had a significantly increased relative risk of cardiovascular events than patients in the lowest one-third, after controlling for LDL-C and other traditional cardiovascular risk factors. This study also compared the discriminatory ability of apo B with that of traditional lipid measures, by using the area under the curve (AUC) for classifying cardiovascular events. Total apo B levels had a slightly higher AUC (0.58) than LDL-C (0.57).

Robinson et al (2012) published results of a Bayesian random-effects meta-analysis of RCTs to compare the effectiveness of lowering apo B versus LDL-C and non-HDL-C for reducing CVD, CHD, and stroke risk.(12) Selected for analysis were 131,134 patients from 25 RCTs

including 12 trials on statins, 5 on niacin, 4 on fibrates, 1 on simvastatin plus ezetimibe, 1 on aggressive versus standard LDL and blood pressure targets, and 1 on ileal bypass surgery. In the analysis of all trials, each apo B decrease of 10 mg/dL resulted in a 6% decrease in major CVD risk and a 9% decrease in CHD risk prediction.

The INTERHEART study compared apo B:apo A-1 with other cholesterol ratios as markers for the risk of acute myocardial infarction in all the major ethnic groups of the world. In this study, 12,461 patients with acute MI from the world's major regions and ethnic groups were compared with 14,637 age- and sex-matched controls to assess the contributions of various cardiovascular risk factors. Investigators obtained non-fasting blood samples from 9,345 cases and 12,120 controls and measured cholesterol fractions and apolipoproteins to determine their respective predictive values. Ratios were stronger predictors of MI than were individual components, and apolipoproteins were better predictors than their cholesterol counterparts. The apo B:apo A-1 ratio was the strongest predictor, with a population-attributable risk of 54%, compared with risks of 37% for LDL/HDL and 32% for total cholesterol/HDL. A 1-standard-deviation increase in apo B:apo A-1 was associated with an odds ratio of 1.59 for MI, compared with 1.17 for an equivalent increase in total cholesterol/HDL. The results were similar for both sexes and across all ethnic groups and ages.(13)

Sniderman et al (2012) reported on 9345 acute myocardial infarction (MI) patients who were compared with 12,120 controls in the standardized case-control INTERHEART study. The authors reported discordance in the levels of cholesterol contained in apo B and non-HDL-C. Apo B was found to be more accurate than non-HDL-C as a marker for cardiovascular risk.(14)

Sniderman et al (2011) performed a meta-analysis of all published reports containing apo B, LDL-C, and non-HDL-C cardiovascular relative risk ratios. Fifteen independent published analyses identified for this meta-analysis provided a total of 233,455 subjects and 22,950 events. The objectives were "to determine the overall balance of the evidence comparing the standardized relative risk ratios of all 3 markers and, if possible, to identify any factors associated with the variance among the studies." The researchers concluded that whether analyzed individually or head to head, apo B was the most potent marker of risk, LDL-C was the least, and non-HDL-C was intermediate. The study indicates that apo B is superior to LDL-C and non-HDL-C as a predictor of cardiovascular risk.(15)

The 2000 Air Force/Texas Coronary Atherosclerosis Prevention Study (AFCAPS/TexCAPS) investigated lipid parameters among 6605 men and women with average LDL-C and low HDL-C levels who were randomized to lovastatin or placebo.(16) This study reported that apo AI levels and the apo B/apo AI ratio were strong predictors of CAD events.

Kastelein et al (2008) combined data from 2 RCTs, the Treating to New Targets (TNT) and Incremental Decrease in End Points through Aggressive Lipid Lowering (IDEAL) trials, to compare the relationship between response to lipids, apo B levels, and other lipid measures.(17) The analysis included 18,889 patients with established coronary disease randomized to low- or high-dose statin treatment. In pairwise comparisons, the on treatment apo B level was a significant predictor of cardiovascular events (HR, 1.24; 95% CI, 1.13 to 1.36; $p < .001$), while LDL level was not. Similarly, the ratio of apo B/apo AI was a significant predictor of events (HR, 1.24; 95% CI, 1.17 to 1.32), while the total cholesterol/HDL-C was not. In another publication that reported on the TNT study (2012), the on treatment apo B level was also a significant predictor of future events (adjusted HR, 1.19; 95% CI, 1.11 to 1.28).(18)

Section Summary: Apolipoprotein B

The evidence has suggested that apo B provides independent information on risk assessment for CVD and that apo B may be superior to LDL-C in predicting cardiovascular risk. Numerous large prospective cohort studies and nested case-control studies have compared these measures, and most have concluded that apo B is a better predictor of cardiac risk than LDL-C.

LDL-C is not the best indicator of the risk associated with LDL, as risk correlates more closely with the number of circulating atherogenic particles than with the quantity of cholesterol transported by LDL particles. In individuals whose LDL particles contain less cholesterol than normal, the LDL-C concentration may underestimate the number of LDL particles. In these individuals, the apo B concentration will more accurately reflect the number of LDL particles and LDL-related CVD risk. Likewise, in individuals whose LDL particles contain more cholesterol than normal, the LDL-C concentration will overestimate the number of LDL particles. In these patients, the apo B concentration will more accurately reflect the number of LDL particles than will the LDL-C concentration.(4)

As a marker of response to cholesterol-lowering treatment, apo B may be more accurate than LDL-C and may provide a better measure of the adequacy of anti-lipid therapy than does LDL-C. Post hoc analyses of RCTs of statin treatment have reported that on-treatment levels of apo B are more highly correlated with clinical outcomes than standard lipid measures.

APOLIPOPROTEIN AI

Systematic Review

In the Emerging Risk Factors Collaboration meta-analysis (2012) described above, apo AI was also examined as an independent risk factor.(6) For apo AI, evidence from 26 studies (n=139,581) reported that apo AI was an independent risk factor for reduced cardiovascular risk. However, as with apo B, when apo AI was substituted for traditional lipids, there was no improvement in risk prediction. In fact, there was a slight worsening in the predictive ability, evidenced by a -0.0028 decrease in the C statistic ($p < .001$) and a -1.08% decrease in the net reclassification improvement ($p < .01$).

Observational Studies

Clarke et al (2007) published a prospective cohort study of 7044 elderly men enrolled in the Whitehall Cardiovascular Cohort from England.(19) Measurements of apolipoprotein levels were performed on 5344 of these men, and they were followed up for a mean of 6.8 years. The authors reported that the apo B/apo AI ratio was a significant independent predictor (Table 1) with similar predictive ability as the TC/HDL ratio (HR, 1.57; 95% CI, 1.32 to 1.86).

Ridker et al (2007), compared the predictive ability of apo AI and the apo B/apo AI ratio with standard lipid measurements.(20) Both ratios had similar predictive ability to standard lipid measurements but were no better. The HR for future cardiovascular events was 1.75 (95% CI, 1.30 to 2.38) for apo AI compared with 2.32 (95% CI, 1.64 to 3.33) for HDL-C (Table 1). The HR for the apo B/apo AI ratio was 3.01 (95% CI, 2.01 to 4.50), compared with 3.18 (95% CI, 2.12 to 4.75) for the LDL-C/HDL-C ratio.

A nested case-control study (2007) performed within the larger European Prospective Investigation into Cancer and Nutrition-Norfolk cohort study, evaluated the predictive ability of the apo B/apo AI ratio in relation to traditional lipid measures in 25,663 patients.(21) The case-

control subgroup study enrolled 869 patients who had developed CAD during a mean follow-up of 6 years and 1511 control patients without CAD. The authors reported that the apo B/apo AI ratio was an independent predictor of cardiovascular events after controlling for traditional lipid risk factors and the Framingham risk score (Table 1). However, the authors also reported that this ratio was no better than the TC/HDL ratio in discriminating between cases (AUC 0.673) and controls (AUC, 0.670; $p=.38$).

Table 1. Results of Diagnostic Apolipoprotein AI Studies

Study	Study Type	N	Efficacy of Apolipoprotein AI in Determining CVD Risk	
			HR (95% CI)	OR (95% CI)
ERFC (2012) ⁶	Review of prospective cohorts	139,581	0.87 (0.84 to 0.90)	-
Clarke et al (2007) ¹⁹	Prospective cohort	7044	1.54 (1.27 to 1.87)	-
Ridker et al (2007) ²⁰	Prospective cohort	2966	2.32 (1.64 to 3.33)	-
van der Steeg et al (2007) ²¹	Case-control	25,663	-	1.85 (1.15 to 2.98)

CI: confidence interval; CVD: cardiovascular disease; ERFC: Emerging Risk Factors Collaboration; HR: hazard ratio; OR: odds ratio.

The Apolipoprotein Mortality Risk Study (2001) followed 175,000 Swedish men and women for 5.5 years and reported that decreased apo AI was an independent predictor of CAD events. (22) The Air Force/Texas Coronary Atherosclerosis Prevention Study (AFCAPS/TexCAPS [2000]) investigated lipid parameters among 6605 men and women with average LDL-C and low HDL-C levels who were randomized to lovastatin or placebo. (16) This study reported that apo AI levels and the apo B/apo AI ratio were strong predictors of CAD events.

The Copenhagen City Heart Study (2007) was a prospective cohort study of 9231 asymptomatic persons from the Danish general population. (11) The apo B/apo AI ratio was reported as an independent predictor of cardiovascular events, with a hazard ratio similar to that for TC/HDL-C. This study also compared the discriminatory ability of the apo B/apo AI ratio with that of traditional lipid measures, using the AUC for classifying cardiovascular events. The apo B/apo AI ratio had a slightly higher AUC (0.59) than the TC/HDL-C ratio (0.58), but this difference was not statistically significant.

Section Summary: Apolipoprotein AI

The current evidence has generally indicated that measurement of apo AI, and the apo B/apo AI ratio are as good as, or better than currently used lipid measures such as LDL and HDL. Some experts have argued that the apo B/apo AI ratio is superior to the LDL/HDL ratio as a predictor of cardiovascular risk and should supplement or replace traditional lipid measures as both a risk marker and a treatment target. However, there is substantial uncertainty regarding the degree of improvement that these measures provide. The evidence suggests that any incremental improvement in predictive ability over traditional measures is likely to be small and of uncertain clinical significance.

APOLIPOPROTEIN E

A large body of research has established a correlation between lipid levels and the underlying *APOE* genotype. For example, in population studies, the presence of an apo e2 allele is associated with the lowest cholesterol levels and the apo e4 allele is associated with the highest levels.(16,17)

Systematic Reviews

A meta-analysis published by Bennet et al (2007) summarized the evidence from 147 studies on the association of *APOE* genotypes with lipid levels and cardiac risk.(23) Eighty-two studies with a total of 86,067 participants included data on the association of apo E and lipid levels, and 121 studies reported on the association with clinical outcomes. The authors estimated that patients with the apo e2 allele had LDL levels that were approximately 31% lower than those in patients with the apo e4 allele. Compared with patients with the apo e3 allele, patients with apo e2 had an approximately 20% lower risk for coronary events (odds ratio [OR], 0.80; 95% CI, 0.70 to 0.90). Patients with the apo e4 had an estimated 6% higher risk of coronary events, which was of marginal statistical significance (OR, 1.06; 95% CI, 0.99 to 1.13).

Sofat et al (2016) published a meta-analysis of 3 studies of circulating apo E and CVD events.(24) The method for selecting the studies was not described. The 3 studies included 9587 participants and 1413 CVD events. In a pooled analysis, there was no association between apo E and CVD events. The unadjusted odds ratio for CVD events for each standard deviation increase in apo E concentration was 1.02 (95% CI, 0.96 to 1.09). After adjustment for other cardiovascular risk factors, the odds ratio for CVD for each standard deviation increase in apo E concentration was 0.97 (95% CI, 0.82 to 1.15).

Observational Studies

Numerous studies have focused on the relation between genotype and physiologic markers of atherosclerotic disease. A number of small- to medium-sized cross-sectional and case-control studies have correlated apo E with surrogate outcomes such as cholesterol levels, markers of inflammation, or carotid intima-media thickness.(25,26,27,28,29,30) These studies have generally shown a relationship between apo E and these surrogate outcomes. Other studies have suggested that carriers of apo e4 are more likely to develop signs of atherosclerosis independent of total and LDL-cholesterol levels.(31,32,33,34)

Some larger observational studies have correlated *APOE* genotype with clinical disease. The Atherosclerosis Risk in Communities (ARIC) study (2001) followed up 12,000 middle-aged subjects free of CAD at baseline for 10 years.(35) This study reported that the e3/2 genotype was associated with carotid artery atherosclerosis after controlling for other atherosclerotic risk factors. Volcik et al (2006), also analyzing ARIC study data, reported that *APOE* polymorphisms were associated with LDL levels and carotid intima-media thickness but were not predictive of incident CAD.(36)

Section Summary: Apolipoprotein E

The evidence has suggested that *APOE* genotype may be associated with lipid levels and CAD but is probably not useful in providing additional clinically relevant information beyond established risk factors. Apo E is considered a relatively poor predictor of CAD, especially compared with other established and emerging clinical variables and does not explain a large percent of the inter-individual variation in total cholesterol (TC) and LDL levels. Moreover, apo E has not been incorporated into standardized cardiac risk assessment models and was not identified as an important “emerging risk factor” in the most recent ATP III recommendations.

HIGH-DENSITY LIPOPROTEIN PARTICLE SIZE AND CONCENTRATION

Systematic Review

Singh et al (2020) reported the results for a pooled analysis examining the association between HDL particle concentration and stroke and MI in patients without baseline atherosclerotic disease.(37) The analysis included 15,784 patients from 4 prospective cohort studies, which included the ARIC study. A significant inverse association was reported between HDL particle concentration and stroke and MI, when comparing patients with HDL particle concentration in the fourth quartile and the first quartile (HR, 0.64; 95% CI, 0.52 to 0.78). When comparing quartile 4 with quartile 1 with regard to the individual components of the primary endpoint, a significant reduction in both MI (HR, 0.63; 95%, 0.49 to 0.81) and stroke (HR, 0.66; 95% CI, 0.48 to 0.93) was reported. There was significant heterogeneity between studies with regard to patient ethnicity and geographic location. Sub-analysis by race revealed that the significant inverse association between HDL particle concentration and stroke and MI was not seen in black populations. When comparing quartile 4 with quartile 1 among black patients, HDL particle concentration did not have an inverse association with MI (HR, 1.22; 95% CI, 0.76 to 1.98). However, the heterogeneity and uneven distribution of patients may have contributed to subgroup analyses being underpowered and the possibility of type 2 error.

Randomized Controlled Trial

In the Justification for the Use of statins in Prevention: an Intervention Trial Evaluating Rosuvastatin (JUPITER RCT) (2013), 10,886 patients without CVD were randomized to rosuvastatin or placebo and followed for a median of 2 years.(38) Before randomization and 1 year after, levels of LDL-C, HDL-C, apo AI, and nuclear magnetic resonance (NMR)-measured HDL size and HDL particle numbers were evaluated. Statistically significant changes in the median and 25th and 75th percentile values of HDL measures between baseline and year 1 values occurred in the rosuvastatin and placebo groups for all levels ($p < .001$), except for apo AI and HDL particle size in the placebo group, which did not differ significantly ($p = .09$ and 0.74 , respectively). Changes in the rosuvastatin group were all statistically significant compared with placebo for LDL-C, HDL-C, apo AI, and HDL particle size and number (all $p < .001$). In the placebo group, inverse associations with CVD and HDL-C, apo AI, and HDL particle were reported. High-density lipoprotein particle number in the rosuvastatin group had a greater association with CVD (HR, 0.73; 95% CI, 0.57 to 0.93; $p = .01$) than HDL-C (HR, 0.82; 95% CI, 0.63 to 1.08; $p = .16$) or apo AI (HR, 0.86; 95% CI, 0.67 to 1.10; $p = .22$). This association remained after adjusting for HDL-C (HR, 0.72; 95% CI, 0.53 to 0.97; $p = .03$). Size of HDL was not significantly associated with CVD in risk factor-adjusted models.

Section Summary: High-Density Lipoprotein Particle Size and Concentration

One RCT and a pooled analysis have evaluated the association of HDL particle size and number as measured by NMR. While these studies found an association with HDL particle concentration (but not HDL size) and CVD, it is uncertain how NMR-measured HDL particle number would be used to change clinical management beyond the information provided by traditional lipid measures. It is also unclear whether the association between HDL particle concentration and cardiovascular events is seen in all patient populations.

LOW-DENSITY LIPOPROTEIN SUBCLASS AND LOW-DENSITY LIPOPROTEIN PARTICLE SIZE AND CONCENTRATION

Observational Studies

A nested case-control study (1996) from the Physician's Health Study, a prospective cohort study of approximately 15,000 men, investigated whether LDL particle size is an independent predictor of CAD risk, particularly compared to triglyceride levels.(39) The authors concluded that while LDL particle diameter was associated with risk of MI, this association was not present after adjustment for triglyceride level. Only the triglyceride level was independently significant.

The Quebec Cardiovascular Study evaluated the ability of "nontraditional" lipid risk factors, including LDL size, to predict subsequent CAD events in a prospective cohort of 2155 men followed for 5 years.(40,41) The presence of small LDL particles was associated with a 2.5-fold increased risk for ischemic heart disease after adjustment for traditional lipid values, indicating a level of risk similar to total LDL. This study also suggested an interaction in atherogenic risk between LDL size and apo B levels. In the presence of small LDL particles, elevated apo B levels were associated with a 6-fold increased risk of CAD, whereas when small LDL particles were not present, elevated apo B levels were associated with only a 2-fold increase in risk.

Tzou et al (2005) examined the clinical value of "advanced lipoprotein testing" in 311 randomly selected adults participating in the Bogalusa Heart Study.(42) Advanced lipoprotein testing consisted of subclass patterns of LDL (i.e., presence of large buoyant particles, intermediate particles, or small dense particles). These measurements were used to predict the presence of subclinical atherosclerosis, as measured ultrasonographically by carotid intimal-media thickness. In multivariate logistic regression models, substituting advanced lipoprotein testing for corresponding traditional lipoprotein values did not improve prediction of the highest quartile of carotid intimal-media thickness.

LOW-DENSITY LIPOPROTEIN PARTICLE SIZE AND CONCENTRATION MEASURED BY NUCLEAR MAGNETIC RESONANCE

Similar to small dense lipoprotein particles, several epidemiologic studies have shown that the lipoprotein particle size and concentration measured by nuclear magnetic resonance (NMR) are also associated with cardiac risk. For example, data derived from the Women's Health Study, Cardiovascular Health Study, and Pravastatin Limitation of Atherosclerosis in the Coronary Arteries (PLAC-1) trial have suggested that the number of LDL particles is an independent predictor of cardiac risk.(43,44,45) Translating these findings into clinical practice requires setting target values for lipoprotein numbers. Proposed target values have been derived from the same data set (i.e., Framingham study) used to set the ATP III target goals for LDL-C. For example, the ATP III targets for LDL-C correspond to the 20th, 50th, and 80th percentile values in the Framingham Offspring Study, depending on the number of risk factors present. Proposed target goals for lipoprotein numbers correspond to the same percentile values, and LDL particle concentrations corresponding to the 20th, 50th, and 80th percentile are 1100, 1400, and 1800 nmol/L, respectively.(46)

Systematic Review

Rosenson and Underberg (2013) conducted a systematic review of studies on lipid-lowering pharmacotherapies to evaluate changes in LDL particles pre- and post-treatment.(47) Reductions in mean LDL particles occurred in 34 of the 36 studies evaluated. Percentage

reductions of LDL particles in several statin studies were smaller than reductions in LDL-C. LDL particles and apo B changes were comparable. Reviewers suggested the differences in LDL particle reductions with different lipid-lowering therapies demonstrated potential areas of residual cardiovascular risk that could be addressed with LDL particle monitoring.

Observational Studies

Mora et al (2009) evaluated the predictive ability of LDL particle size and number measured by NMR in participants of the Women's Health Study, a prospective cohort study of 27,673 women followed over an 11-year period.(48) After controlling for nonlipid factors, LDL particle number was a significant predictor of incident CVD, with a hazard ratio of 2.51 (95% CI, 1.91 to 3.30) for the highest, compared with the lowest quintile. LDL particle size was similarly predictive of cardiovascular risk, with a hazard ratio of 0.64 (95% CI, 0.52 to 0.79). Compared with standard lipid measures and apolipoproteins, LDL particle size and number showed similar predictive ability but were not superior in predicting cardiovascular events.

Toth et al (2014) analyzed LDL-C and LDL particle levels and cardiovascular risk using commercial insurance and Medicare claims data on 15,569 high-risk patients from the HealthCore Integrated Research Database.(49) For each 100 nmol/L increase in LDL particle level, there was a 4% increase in risk of a CHD event (HR, 1.04; 95% CI, 1.02 to 1.05; $p < 0.001$). A comparative analysis, using 1:1 propensity score matching of 2094 patients from the LDL-C target cohort (LDL-C level < 100 mg/dL without a LDL particle level) and a LDL particle target cohort (LDL particle < 1000 nmol/L and LDL-C of any level) found a lower risk of CHD or stroke in patients who received LDL-C measurement and were presumed to have received more intensive lipid-lowering therapy (HR, 0.76; 95% CI, 0.61 to 0.96; at 12 months). A comparison of smaller LDL particle target groups at 24 ($n=1242$) and 36 ($n=705$) months showed similar reductions in CHD (HR=, 0.78; 95% CI, 0.62 to 0.97) and stroke (HR, 0.75; 95% CI, 0.58 to 0.97).

Section Summary: Low-Density Lipoprotein Subclass and Low-Density Lipoprotein Particle Size and Concentration

Small LDL size is a component of an atherogenic lipid profile; other components include increased triglycerides, increased apo B, and decreased HDL. Some studies have reported that LDL size is an independent risk factor for CAD, while others have reported that a shift in LDL size may be a useful marker of treatment response.

A relatively small number of studies have evaluated the predictive ability of LDL particle size and number as measured by NMR. These studies do not demonstrate that NMR-measured particle size and/or number offer predictive ability beyond that provided by traditional lipid measures. Measures by NMR have been proposed as indicators of residual cardiovascular risk in patients treated with statins who have met LDL goals, but there is no evidence that these measures improve health outcomes when used for this purpose.

LIPOPROTEIN (A)

Numerous prospective RCTs, cohort studies and systematic reviews have evaluated lipoprotein (a) [Lp(a)] as a cardiovascular risk factor. The following are representative prospective trials drawn from the relevant literature. Table 2 summarizes the results of diagnostic Lp(a) studies that assess the HR or OR of the efficacy of Lp(a) in determining CVD risk.

Systematic Review

The Emerging Risk Factors Collaboration (2012) published a patient-level meta-analysis assessing 37 prospective cohort studies enrolling 154,544 individuals.(6) Risk prediction was examined for a variety of traditional and nontraditional lipid markers. For Lp(a), evidence from 24 studies on 133,502 subjects reported that Lp(a) was an independent risk factor for reduced cardiovascular risk (Table 2). The addition of Lp(a) to traditional risk factors resulted in a small improvement in risk prediction, with a 0.002 increase in the C statistic. A reclassification analysis found no significant improvement in the net reclassification index (0.05%; 95% CI, -0.59 to 0.70).

Several meta-analyses have also examined the relation between Lp(a) levels and cardiovascular risk. Bennet et al (2008) synthesized the results of 31 prospective studies with at least 1 year of follow-up and that reported data on cardiovascular death and nonfatal MI. (50) The combined results revealed a significant positive relationship between Lp(a) and cardiovascular risk (Table 2). This analysis reported a moderately high degree of heterogeneity in selected studies ($I^2=43\%$), reflecting the fact that not all reported a significant positive association.

Smolders et al (2007) summarized evidence from observational studies on the relation between Lp(a) and stroke.(51) Five prospective cohort studies and 23 case-control studies were included in this meta-analysis. Results from prospective cohort studies showed that Lp(a) level added only incremental predictive information (combined RR for the highest one-third of Lp[a], 1.22; 95% CI, 1.04 to 1.43). Results from case-control studies showed an elevated Lp(a) level was associated with an increased risk of stroke (Table 2).

Randomized Controlled Trials

Several RCTs on lipid-lowering therapies have found Lp(a) is associated with residual cardiovascular risk. In a subgroup analysis of 7746 white patients from the JUPITER study (2014), median Lp(a) levels did not change in either group of patients randomized to treatment with rosuvastatin or placebo during a median 2-year follow-up.(52) Lp(a) was independently associated with a residual risk of CVD despite statin treatment. In the Atherothrombosis Intervention in Metabolic Syndrome with Low HDL/High Triglyceride and Impact on Global Health Outcomes study (2013), Lp(a) levels in 1440 patients at baseline and on simvastatin plus placebo or simvastatin plus extended-release niacin were significantly predictive of cardiovascular events (Table 2).(53)

Observational Studies

Kamstrup et al (2008) analyzed data from the Copenhagen City Heart Study, which followed 9330 subjects from the Copenhagen general population over 10 years.(54) This study reported on a graded increase in the risk of cardiac events with increasing Lp(a) levels. At extreme levels of Lp(a) above the 95th percentile, the aHR for MI was slightly higher for women than for men (Table 2). Tzoulaki et al (2007) reported on data from the Edinburgh Artery Study, a population cohort study that followed 1592 subjects for a mean of 17 years.(55) They reported that Lp(a) was an independent predictor of MI (Table 2).

Zakai et al (2007) evaluated 13 potential biomarkers for independent predictive ability compared with established risk factors, using data from 4510 subjects followed for 9 years in the Cardiovascular Health Study.(56) Lipoprotein (a) was 1 of 7 biomarkers that had incremental predictive ability above the established risk factors (Table 2).

Waldeyer et al (2017) analyzed data of 56,084 participants from Biomarkers for Cardiovascular Risk Assessment in Europe project, which followed 7 prospective population-based cohorts

across Europe, with a maximum follow-up of 24 years, to characterize the association of Lp(a) concentration with major coronary events, incident CVD, and total mortality.(57) The highest event rate of major coronary events and CVD was observed for Lp(a) levels at the 90th percentile or higher ($p<.001$ for major coronary events and CVD). Adjusting for age, sex, and cardiovascular risk factors, compared with Lp(a) levels in the lowest third in the 67th to 89th percentile, there were significant associations between Lp(a) levels and major coronary events (HR, 1.3; 95% CI, 1.15 to 1.46) and CVD (HR, 1.25; 95% CI, 1.12 to 1.39) (Table 2). For Lp(a) levels at the 90th percentile or higher, the aHR for the association between Lp(a) and major coronary events was 1.49 (95% CI, 1.29 to 1.73) and for the association between Lp(a) and CVD, it was 1.44 (95% CI, 1.25 to 1.65) compared with Lp(a) levels in the lowest third. There was no significant association between Lp(a) levels and total mortality.

Lee et al (2017) investigated whether elevated circulating Lp(a) level was a key determinant in predicting the incidence of major adverse cardiovascular events among the participants of the Dallas Health Study, a multiethnic prospective cohort with a median follow-up of 9.5 years (N=3419 patients).(58) Quartiles 4 of Lp(a) and oxidized phospholipid on apo B-100 were associated with HRs for time to major adverse cardiovascular events of 2.35 (95% CI, 1.50 to 3.69) and 1.89 (95%CI, 1.26 to 2.84), respectively, adjusting for age, sex, body mass index (BMI), diabetes, smoking, LDL, HDL-C, and triglycerides (Table 2). The addition of major apolipoprotein(a) isoform and 3 *LPA* single nucleotide variants prevalent among white, black, and Hispanic subjects in the model attenuated the risk, but significance was maintained for both Lp(a) and oxidized phospholipid on apo B-100.

Some researchers have hypothesized that there is a stronger relation between Lp(a) and stroke than CHD. Similar to the situation with cardiac disease, most prospective studies have indicated that Lp(a) level is an independent risk factor for stroke. In a prospective cohort study, Rigal et al (2007) reported that an elevated Lp(a) level was an independent predictor of ischemic stroke in men (Table 2).(59)

There also may be a link between Lp(a) level as a cardiovascular risk factor and hormone status in women. Suk Danik et al (2008) reported on the risk of a first cardiovascular event over a 10-year period in 27,736 women enrolled in the Women's Health Study.(60) After controlling for standard cardiovascular risk factors, Lp(a) levels were an independent predictor of risk in women not taking hormone replacement therapy (Table 2). However, for women who were taking hormone replacement therapy, Lp(a) levels were not a significant independent predictor of cardiovascular risk (HR, 1.13; 95% CI, 0.84 to 1.53; $p=.18$).

Table 2. Results of Diagnostic Lipoprotein(a) Studies

Study	Study Type	N	Efficacy of Lp(a) in Determining CVD Risk	
			HR (95% CI)	OR (95% CI)
ERFC (2012) ⁶	SR/MA	154,544	1.13 (1.09 to 1.18)	-
Khera et al (2014) ⁵²	RCT	7746	1.27 (1.01 to 1.59) $p=.04$	-
Albers et al (2013) ⁵³	RCT	1440	1.18-1.25	-

Kamstrup et al (2008) ⁵⁴	Post hoc analysis	9330	Men: 3.6 (1.7 to 7.7) Women: 3.7 (1.7 to 8.0)	-
Tzoulaki et al (2007) ⁵⁵	Prospective cohort	1592	1.49 (1.0 to 2.2)	-
Zakai et al (2007) ⁵⁶	Prospective cohort	4510	1.07 (1.0 to 1.12)	-
Waldeyer et al (2017) ⁵⁷	Post hoc analysis	56,084	1.3 (1.15 to 1.46)	-
Lee et al (2017) ⁵⁸	Prospective cohort	3419	2.35 (1.50 to 3.69)	-
Rigal et al (2007) ⁵⁹	Prospective cohort	100	-	Men: 3.55 (1.33 to 9.48) Women: 0.42 (0.12 to 1.26)
Suk Danik et al (2008) ⁶⁰	Prospective cohort	27,736	1.77 (1.36 to 2.30) p<.001	-
Bennet et al (2008) ⁵⁰	SR/MA	2047	-	1.45 (1.32 to 1.58)
Smolders et al (2007) ⁵¹	SR/MA of Observational	56,010	-	2.39 (1.57 to 3.63)

CI: confidence interval; CVD: cardiovascular disease; ERFC: Emerging Risk Factors Collaboration; HR: hazard ratio; MA: meta-analysis; Lp(a): lipoprotein(a); OR: odds ratio; RCT: randomized control trial; SR: systematic review.

Additional Studies

Beyond the studies describing the HR or OR for the efficacy of Lp(a) and CVD summarized in Table 2, additional key studies have examined the relation between Lp(a) and CVD risk, which are summarized below.

Additional Systematic Reviews

A systematic review by Genser et al (2011) included 67 prospective studies (N=181,683) that evaluated the risk of CVD associated with Lp(a).⁽⁶¹⁾ Pooled analysis was performed on 37 studies that reported the endpoints of cardiovascular events. When grouped by design and populations, the RRs for these studies, comparing the uppermost and lowest strata of Lp(a), ranged from 1.64 to 2.37. The RR for cardiovascular events was higher in patients with previous CVD than with patients without the previous disease. There were no significant associations found between Lp(a) levels, overall mortality, or stroke.

A patient-level meta-analysis (2009) of 36 prospective studies published between 1970 and 2009 included 126,634 participants.⁽⁶²⁾ Overall, the independent association between Lp(a) level and vascular disease was consistent across studies but modest in size. The combined RR, adjusted for age, sex, and traditional lipid risk factor, was 1.13 (95% CI, 1.09 to 1.18) for CHD and 1.10 (95% CI, 1.02 to 1.18) for ischemic stroke. There was no association between Lp(a) levels and mortality.

A systematic review and meta-analysis by Vazirian et al (2023) included 8 cross-sectional studies (n=18,668) and 4 cohort studies (n=15,355) to evaluate the association between risk of coronary artery calcification and elevated Lp(a) levels.⁽⁶³⁾ The pooled OR for coronary artery calcium scores among asymptomatic cardiovascular disease patients with elevated Lp(a) levels was 1.08 (95% CI, 1.02 to 1.13), with a substantial level of variability between studies

($I^2=90.6\%$; $p=.00$). Data from the 4 cohort studies reported a positive significant association between Lp(a) levels and coronary artery calcification, the combined OR for coronary artery calcification incidence was 1.58 (95% CI, 1.38 to 1.80), with no significant variability observed between the studies ($I^2=0.0\%$; $p=.483$).

Additional Randomized Controlled Trials

The Lipid Research Clinics Coronary Primary Prevention Trial (1994), one of the first large-scale RCTs of cholesterol-lowering therapy, measured initial Lp(a) levels and reported that Lp(a) was an independent risk factor for CAD when controlling for other lipid and non-lipid risk factors.(64)

The LIPID RCT (2013) randomized 7863 patients to pravastatin or placebo.(65) Patients were followed for a median of 6 years. Lipoprotein (a) concentrations did not change significantly at 1 year. Baseline Lp(a) concentration was associated with total CHD events ($p<.001$), total CVD events ($p=.002$), and coronary events ($p=.03$).

Additional Observational Studies

As part of the Framingham Offspring Study, Lp(a) levels were measured in 2191 asymptomatic men between the ages of 20 and 54 years.(66) After a mean follow-up of 15 years, there were 129 CHD events, including MI, coronary insufficiency, angina, or sudden cardiac death. Comparing the Lp(a) levels of these patients with the other participants, the authors concluded that elevated Lp(a) was an independent risk factor for the development of premature CHD (i.e., before age 55 years). The ARIC study (2001) evaluated the predictive ability of Lp(a) in 12,000 middle-aged subjects free of CAD at baseline who were followed for 10 years.(35) Lipoprotein (a) levels were significantly higher among patients who developed CAD than among those who did not, and Lp(a) levels were an independent predictor of CAD above traditional lipid measures.

In the ARIC prospective cohort study of 14,221 participants, elevated Lp(a) was a significant independent predictor of stroke in Black women (RR, 1.84; 95% CI, 1.05 to 3.07) and White women (RR, 2.42; 95% CI, 1.30 to 4.53) but not in Black men (RR, 1.72; 95% CI, 0.86 to 3.48) or white men (RR, 1.18; 95% CI, 0.47 to 2.90).(67)

Fogacci et al (2017) examined whether serum Lp(a) levels could predict long-term survival in 1215 adults with no CVD at enrollment and similar general cardiovascular risk profiles from Brisighella Heart Study cohort in Italy.(68) Subjects were stratified into a low ($n=865$), intermediate ($n=275$), and high ($n=75$) cardiovascular risk groups using an Italian-specific risk chart. Subjects at high and intermediate cardiovascular risk ages 56 to 69 years (regardless of sex) and women ages 40 to 55 years with a low cardiovascular risk profile who had lower Lp(a) levels showed statistically significant lower cardiovascular mortality ($p<.05$) and longer survival time ($p<.05$) during the 25-year follow-up. The authors constructed a receiver operating characteristic curve for each cardiovascular risk group using Lp(a) as a test variable and death as a state variable and identified serum Lp(a) as an independent long-term cardiovascular mortality prognostic indicator for subjects at high cardiovascular risk (AUC, 0.63; 95% CI, 0.50 to 0.76; $p=.049$) and for women at intermediate cardiovascular risk (AUC, 0.7; 95% CI, 0.52 to 0.79; $p=.034$).

Some studies, however, have failed to demonstrate such predictive ability. In the Physicians' Health Study (1993), initial Lp(a) levels in the 296 participants who subsequently experienced MI were compared with Lp(a) levels in matched controls who remained free from CAD.(69)

Authors found that the distribution of Lp(a) levels between the groups was identical. The European Concerted Action on Thrombosis and Disabilities study (2000), a trial of secondary prevention, evaluated Lp(a) as a risk factor for coronary events in 2800 patients with known angina pectoris.(70) In this study, Lp(a) levels did not differ significantly among patients who did and did not have subsequent events, suggesting that Lp(a) levels were not useful risk markers in this population.

Genetic studies have examined the association between various genetic loci and Lp(a) levels, and Mendelian randomization studies have examined whether Lp(a) level is likely to be causative for CAD. In a 2009 study, 3 separate loci were identified for increased Lp(a) levels.(71) Genetic variants identified at 2 of these loci that were independently associated with coronary disease (OR, 1.70; 95% CI, 1.49 to 1.95; OR, 1.92; 95% CI, 1.48 to 2.49). This finding strongly implies that elevated Lp(a) levels are causative of coronary disease, as opposed to simply being associated.

Lipoprotein(a) is recognized as a genetically determined, causal and prevalent risk factor for atherosclerotic cardiovascular disease. The American Heart Association performed an expert peer review of epidemiologic and genetic data in development of their scientific statement in 2022. Evidence revealed elevated levels of La(a) are prevalent in approximately 20% of the population. Observational and genetic data strongly supported an independent and causal relationship between high plasma concentrations of Lp(a) and increased risk of atherosclerotic cardiovascular disease-related events, such as myocardial infarction, stroke, and valvular stenosis. In the largest single study to date, a multiethnic population of 460,506 participants of the UK Biobank were examined. Concentrations of Lp(a) were variable across all racial subgroups and the associated risk were found to be similar. Awareness of the presence of elevated Lp(a) is important because it could inform clinical decision making. Presently the group states the evidence is in favor of performing targeted screening in selected individuals and to consider cascade screening when appropriate to identify those with high Lp(a). To lower overall ASCVD risk, it is recommended to target LDL-C with statin and adjunctive medications as initial therapy.(72,73)

Section Summary: Lipoprotein (a)

A large amount of epidemiologic evidence has determined that Lp(a) is an independent risk factor for cardiovascular disease. The overall degree of risk associated with Lp(a) levels appears to be modest, and the degree of risk may be mediated by other factors such as LDL levels and/or hormonal status.

B-TYPE OR BRAIN NATRIURETIC PEPTIDE

Observational Studies

The use of B-type or brain natriuretic peptide (BNP) levels for monitoring and managing established heart failure patients has been frequently studied and has demonstrated value. Studies on the use of BNP for determining cardiovascular risk in the asymptomatic population, however, are limited. In the Early Identification of Subclinical Atherosclerosis by Noninvasive Imaging Research study, Shaw et al (2009) evaluated BNP and coronary artery calcium levels in 2458 asymptomatic adults.(74) Levels of BNP ranging from 40 to 99.9 and 100 pg/m or higher had a 2.2 to 7.5 relative hazard for a cardiovascular event compared to BNP levels of less than 40 pg/ml ($p<.001$). Other large population cohort studies have shown a relationship between elevations in BNP levels and future risks of cardiovascular events or heart failure. Wu et al (2022) assessed the value of cardiac troponins and amino terminal B type cardiac

natriuretic peptide (NT-proBNP) in 2 different cohorts of asymptomatic patients (n=4102; n=2538).(75) Study investigators found that cardiac marker data correctly reclassified risk upwards in 6.7% of patients and downwards in 3.3% of patients; the overall C statistic for discrimination of the primary endpoint (composite of all first CV events) increased from 0.755 to 0.771 (+0.016, p=.01).In a cohort study (N=5067), Melander et al found adding C-reactive protein and BNP to a risk model of conventional factors increased the C-statistic for cardiovascular events by 0.007 (p=.04) and for coronary events by 0.009 (p=.08).(76) In a cohort study of 3346 patients without heart failure, Wang et al found BNP levels above the 80th percentile (20.0 pg/mL for men, 23.3 pg/mL for women) were associated with multivariable-adjusted hazard ratios of 1.62 for death (p=.02), 1.76 for a first major coronary event, (p=.03), 1.91 for atrial fibrillation (p=.02), 1.99 for stroke or transient ischemic attack (p=.02), and 3.07 for heart failure (p=.002).(77) However, any gains over use of conventional risk factors appear to be minimal.

Section Summary: B-Type or Brain Natriuretic Peptide

Levels of BNP appear to be associated with cardiovascular risks. However, no evidence was identified demonstrating that the use of BNP testing in clinical care improves outcomes.

CYSTATIN C

Ito et al (2011) evaluated the value of adding cystatin C to Framingham Risk Score variables to predict cardiovascular disease risk in 6653 adults without CVD from the Multi-Ethnic Study of Atherosclerosis.(78) Cardiovascular risk prediction did not improve with the addition of cystatin C to Framingham Risk Score variables. Lee et al (2010) conducted a meta-analysis of 14 studies (N=22,509) with predominantly high-cardiovascular-risk patients to evaluate the relation between elevated cystatin C levels and CVD risk.(79) Higher levels of cystatin C were associated with greater risk of CVD (RR=2.62; 95% CI, 2.05 to 3.37; p<0.001), CHD (RR, 1.72; 95% CI, 1.27 to 2.34; p<.001), and stroke (RR, 1.83; 95% CI, 1.12 to 3.00; p=.02) after adjusting for known cardiovascular risk factors. Luo et al (2015) reported on a meta-analysis of studies evaluating the relation between cystatin C and cardiovascular and all-cause mortality in the general population.(80) Reviewers included 9 prospective studies (N=39,854). Across the 6 studies reporting cardiovascular mortality-specific outcomes, the pooled aHR of cardiovascular mortality, comparing the highest and lowest cystatin C categories, was 2.74 (95% CI, 2.04 to 3.68, p=.021).

Section Summary: Cystatin C

Several meta-analyses have reported that higher levels of cystatin C are associated with higher cardiovascular risk and higher risk of cardiovascular death. In contrast, in a large cohort, cystatin C did not improve risk prediction of CVD. No evidence was identified demonstrating that the use of cystatin C testing in clinical care improves outcomes.

FIBRINOGEN

Systematic Reviews

Kengne et al (2013) evaluated data from 9 prospective, community-based cohorts from the British and Scottish general population-based health surveys.(81) In the analysis of 33,091 adults, 1006 of whom had diabetes, fibrinogen was positively associated with a higher risk of CVD by 34% (95% CI, 26% to 42%) and all-cause mortality by 30% (95% CI, 26% to 35%). The relation between cardiovascular mortality and higher fibrinogen produced HRs of 1.48

(95% CI, 1.21 to 1.81) in subjects with diabetes and 1.31 (95% CI, 1.23 to 1.39) in those without diabetes. The interaction between fibrinogen and CVD risk did not differ significantly between the diabetic and nondiabetic populations ($p=.47$). Despite improved predictive accuracy, the addition of fibrinogen to established risk factors was reported to not be clinically important.

Willeit et al (2016) reported results of a patient-level meta-analysis from 20 prospective studies to assess the association between a number of inflammatory markers (including fibrinogen) and atherosclerosis among patients without preexisting CVD.(82) Selected were prospective cohort studies from the PROG-IMT collaboration, which included participants from the general population and reported at least 2 visits with measurements of common carotid artery intima-media thickness as a marker of preclinical atherosclerosis, along with at least 1 inflammatory marker (high-sensitivity-CRP, leukocyte count, and/or fibrinogen). Overall, reviewers included 20 studies ($N=49,087$), of which 13 studies ($n=35,096$) reported fibrinogen levels. In a cross-sectional analysis, a 1 SD higher baseline fibrinogen level was associated with common carotid artery intima-media thickness (mean, 0.0073 mm; 95% CI, 0.0047 to 0.0097; $p<.001$). However, in longitudinal analysis, neither the baseline level of any of the inflammatory markers evaluated nor their progression was associated with progression of common carotid artery intima-media thickness.

Observational Studies

Other studies have found an association between fibrinogen and cardiovascular risk including the European Prospective Investigation into Cancer and Nutrition-Norfolk cohort study (83) and the Fibrinogen Studies Collaboration.(84,85) In a 2007 report from the Fibrinogen Studies Collaboration, it was noted that fibrinogen levels increased with age and were linked to established risk factors such as triglycerides, smoking and body mass index.(85)

Section Summary: Fibrinogen

Reports from a number of cohort studies suggest that fibrinogen levels are associated with cardiovascular risk. However, no evidence was identified demonstrating that the use of fibrinogen testing in clinical care improves outcomes.

LEPTIN

Systematic Reviews

Sattar et al (2009) reported on a prospective study of 5661 men and a systematic review of 7 prospective studies to evaluate the relationship between leptin and CVD.(86) Leptin levels in the top third had an odds for coronary heart disease of 1.25 (95% CI, 0.96 to 1.62) compared with the bottom third. After adjusting for body mass index (BMI), the odds decreased to 0.98 (95% CI, 0.72 to 1.34) suggesting an association of leptin with CVD disease is largely dependent on BMI.

Zeng et al (2014) conducted a meta-analysis of studies reporting the association between leptin levels and risk of CHD or stroke.(87) The meta-analysis included 8 nested case-control studies with 1980 patients and 11,567 controls. In a pooled analysis, leptin levels were significantly associated with pathogenic risk of CHD (OR, 1.90; 95% CI, 1.06 to 3.43; $p=.032$) and pathogenic risk of stroke (OR, 2.14; 95% CI, 1.48 to 3.08; $p<.001$).

Yang et al (2017) conducted a systematic review of case-control and cohort studies that assessed leptin concentration and CHD risk.(88) Thirteen epidemiologic studies totaling 4257

CVD patients and 26,710 controls were included. Adjusting for cardiovascular risk factors, there was no statistically significant association between leptin concentration and CHD risk (OR, 1.16; 95% CI, 0.97 to 1.40). The association did not change when analyses were restricted to high-quality studies (OR, 1.07; 95% CI, 0.96 to 1.19) for CHD. In a subgroup meta-analysis, a high leptin level was not independently associated with CHD in both females (OR, 1.03; 95% CI, 0.86 to 1.23) and male patients (OR, 1.09; 95% CI, 0.95 to 1.26).

Section Summary: Leptin

Two meta-analyses suggest that leptin levels are associated with CHD and stroke, although this association may depend on BMI. Another meta-analysis suggested no significant association between leptin concentration and CHD risk. No evidence was identified demonstrating that the use of leptin testing in clinical care improves outcomes.

Section Summary: Asymptomatic Individuals with Risk of Cardiovascular Disease

The evidence for asymptomatic individuals with risk of CVD who receive nontraditional cardiac biomarker testing includes systematic reviews, meta-analyses, and large, prospective cohort studies. The evidence from cohort studies and meta-analyses of these studies has suggested that some of these markers are associated with increased cardiovascular risk and may provide incremental accuracy in risk prediction. Several biomarkers have been identified as adding some incremental predictive value. However, it has not been established whether the incremental accuracy provides clinically important information beyond that of traditional lipid measures. Furthermore, no study has provided high-quality evidence that measurement of markers leads to changes in management that improve health outcomes.

INDIVIDUALS WITH HYPERLIPIDEMIA MANAGED WITH LIPID-LOWERING THERAPY

APOLIPOPROTEIN B

Systematic Reviews

A number of RCTs of statin therapy have examined the change in apo B on-treatment in relation to clinical CAD outcomes and assessed whether apo B predicted outcomes better than LDL-C.

Boekholdt et al (2012) published a patient-level meta-analysis of on-treatment levels of traditional and nontraditional lipids as a measure of residual risk.(89) Eight studies enrolling 62,154 participants were included. The aHR for each 1 SD increase in apo B was 1.14 (95% CI, 1.11 to 1.18), which did not differ significantly from LDL-C (aHR, 1.13; 95% CI, 1.10 to 1.17; $p=.21$). The aHR for HDL-C was 1.16 (95% CI, 1.12 to 1.19), which was significantly greater than LDL-C or apo B ($p=.002$). In a subsequent report from this meta-analysis, Boekholdt et al (2014) evaluated the LDL-C, non-HDL-C, and apo B levels of 38,153 patients allocated to the statin therapy groups.(90) Despite statin therapy, reductions in levels of LDL-C, non-HDL-C, and apo B from baseline to 1 year showed large interindividual variations.

Randomized Controlled Trials

Ballantyne et al (2013) reported on a post hoc analysis of 682 patients with acute coronary syndrome from the randomized, phase 3 Limiting Undertreatment of Lipids in Acute coronary syndrome with Rosuvastatin trial.(91) The Limiting Undertreatment of Lipids in Acute coronary syndrome with Rosuvastatin subgroup analysis examined apo B in relation to LDL-C and non-HDL-C under intensive statin therapy with rosuvastatin or atorvastatin. The treatment target level for apo B of 80 mg/dL correlated with an LDL-C level of 90 mg/dL and a non-HDL-C level

of 110 mg/dL at baseline and with an LDL-C of 74 mg/dL and a non-HDL-C of 92 mg/dL with statin therapy. Independent of triglyceride status, non-HDL-C was found to have a stronger correlation with apo B than with LDL-C and could be an adequate surrogate for apo B during statin therapy.

The AFCAPS/TexCAPS (2000) evaluated lipid parameters among 6605 men and women with average LDL-C and low HDL-C levels who were randomized to lovastatin or placebo.(16) Baseline LDL-C, HDL-C, and apo B levels were predictive of future coronary events. However, in the treatment group, post-treatment levels of LDL-C and HDL-C were not predictive of subsequent risk, while post-treatment apo B levels were.

In the Long-term Intervention with Pravastatin in Ischemic Disease trial (2002), the relation between on-treatment apo B levels and clinical outcomes was examined in 9140 patients randomized to pravastatin or placebo and followed for a mean of 6.1 years.(25) The aHR for apo B levels (2.10; 95% CI, 1.21 to 3.64, $p=.008$) was higher than that for LDL-C (1.20;95% CI, 1.00 to 1.45; $p=.05$). Also, the proportion of the treatment effect explained by on-treatment apo B levels (67%) was higher than that for LDL-C levels (52%).

Kastelein et al (2008) combined data from 2 RCTs, the Treating to New Targets (TNT) and Incremental Decrease in End Points through Aggressive Lipid Lowering trials, to compare the relation between response to lipids, apo B levels, and other lipid measures.(17) The analysis included 18,889 patients with established coronary disease randomized to low- or high-dose statin treatment. In pairwise comparisons, the on treatment apo B level was a significant predictor of cardiovascular events (HR=1.24; 95% CI, 1.13 to 1.36; $p<.001$), while LDL level was not. Similarly, the ratio of apo B/apo AI was a significant predictor of events (HR, 1.24; 95% CI, 1.17 to 1.32), while the TC/HDL-C ratio was not. In another publication that reported on the TNT study (2012), the on treatment apo B level was also a significant predictor of future events (aHR, 1.19; 95% CI, 1.11 to 1.28).(18) In this study, the known baseline variables performed well in discriminating future cases from non-cases, and the addition of apo B was not associated with additional risk.

Mora et al (2012) measured on-treatment lipid levels to assess the prediction of residual risk while on statin therapy.(92) Using data from the JUPITER trial, on-treatment levels of LDL-C, non-HDL-C, high-sensitivity CRP, apo B, and apo AI were used to predict subsequent cardiovascular events. The HRs for cardiovascular events were similar among the lipid measures, ranging from 1.22 to 1.31, with no significant differences between them. The residual risk declined overall with a decreasing level of LDL-C, with the lowest risk seen in subjects achieving an LDL-C level of less than 70 mg/dL.

Section Summary: Apolipoprotein B

As a marker of response to cholesterol-lowering treatment, apo B may be more accurate than LDL-C and may provide a better measure of the adequacy of antilipid therapy than LDL-C. Post hoc analyses of RCTs of statin treatment have reported that on-treatment levels of apo B are more highly correlated with clinical outcomes than standard lipid measures.

APOLIPOPROTEIN AI

Randomized Controlled Trials

A number of studies have evaluated the utility of the apo B/apo AI ratio as a marker of treatment response in RCTs of statin treatment. For example, in the Kastelein et al (2008)

study (described above), authors combined data from 2 RCTs, the TNT and Incremental Decrease in End Points Through Aggressive Lipid Lowering trials, to compare the relation between response to lipids, apo B/apo AI ratio, and other lipid measures.(17) The apo B/apo AI ratio was a significant predictor of events (HR, 1.24; 95% CI, 1.17 to 1.32) while the TC/HDL-C was not.

The Pravastatin or Atorvastatin Evaluation and Infection Therapy-Thrombolysis in MI (PROVE-IT TIMI) study (2009) randomized 4162 patients with an acute coronary syndrome to standard statin therapy or intensive statin therapy.(93) While the on-treatment apo B/apo AI ratio was a significant predictor of cardiac events (HR for each SD increment, 1.10; 95% CI, 1.01 to 1.20); it was not superior to LDL-C (HR, 1.20; 95% CI, 1.07 to 1.35) or the TC/HDL ratio (HR, 1.12; 95% CI, 1.01 to 1.24) as a predictor of cardiac events.

Preliminary studies of infusions of reconstituted apo AI have demonstrated plaque regression in a small number of patients with acute coronary syndrome.(94) Based on this research, there has been interest in developing synthetic apo AI mimetic proteins, and such agents are in the drug development stage. These types of agents would likely target patients with residual cardiac risk following maximal statin therapy, especially patients with low HDL levels.

Section Summary: Apolipoprotein AI

The use of apo AI and the apo B/apo AI ratio as a target of treatment response to statins may also be as good as or better than the traditional measure of LDL. However, to improve outcomes, clinicians must have the tools to translate this information into clinical practice. Such tools for linking apo AI to clinical decision making, both in risk assessment and treatment response, are currently not available. Apolipoprotein AI has not been incorporated into quantitative risk assessment models or treatment guidelines that can be used in clinical practice (e.g., the ATP III).(1) The ATP III practice guidelines continue to tie clinical decision making to conventional lipid measures, such as TC, LDL-C, and HDL-C. Therefore, it is not yet possible to conclude that these measures improve outcomes or that they should be adopted in routine clinical care. There is continued interest in developing new therapeutic agents that raise HDL, and apo AI mimetics are currently in development for this purpose.

APOLIPOPROTEIN E

Randomized Controlled Trials

Apolipoprotein E has been investigated as a predictor of response to therapy by examining apo E alleles in the intervention arm(s) of lipid-lowering trials. Some data have suggested that patients with an apo e4 allele may respond better to diet-modification strategies.(95,96) Other studies have suggested that response to statin therapy may vary by *APOE* genotype and that the e2 allele indicates greater responsiveness to statins.(95,97)

Chiodini et al (2007) examined differential response to statin therapy by *APOE* genotype in a reanalysis of data from the Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto-Prevenzione (GISSI-P) study.(98) The GISSI-P was an RCT comparing pravastatin with placebo in 3304 Italian patients with previous MI. Patients with the apo e4 allele treated with statins had a greater response to treatment as evidenced by lower overall mortality rates (1.85% vs 5.28%, respectively, $p=.023$), while there was no difference in mortality rates for patients not treated with statins (2.81% vs 3.67%, respectively, $p=.21$). This study corroborated results reported in previous studies but did not provide evidence that changes in treatment should be made as a result of *APOE* genotype.

Observational Studies

Other studies have evaluated *APOE* genetic status as a predictor of response to lipid-lowering therapy. Donnelly et al (2008) reported on 1383 patients treated with statins from the Genetics of Diabetes Audit and Research in Tayside, Scotland (Go-DARTS) database.(99) Researchers reported on final LDL levels and percentages of patients achieving target LDL by *APOE* genetic status. LDL levels following treatment were lower for patients who were homozygous for apo e2 (0.6 mmol/L), than for patients homozygous for apo e4 (1.7 mmol/L, $p<.001$). All patients who were homozygous for apo e2 reached their target LDL level, compared with 68% of patients homozygous for apo e4 ($p<.001$).

Vossen et al (2008) evaluated response to diet and statin therapy by apo E status in 981 patients with CAD who were enrolled in a cardiac rehabilitation program.(100) They reported that patients with an apo e4 allele were more responsive to diet and statin therapy than were patients with an apo e2 allele. The overall response to treatment was more dependent on baseline LDL levels than *APOE* genetic status, with 30% to 47% of the variation in response to treatment explained by baseline LDL, compared with only 1% of the variation explained by *APOE* status.

Section Summary: Apolipoprotein E

The evidence on the response to treatment indicates that *APOE* genotype may be a predictor of response to statins and may allow clinicians to better gauge a patient's chance of successful treatment, although not all studies have consistently reported this relation. At present, it is unclear how this type of information would change clinical management. Dietary modifications are a universal recommendation for those with elevated cholesterol or LDL levels, and statin drugs are the overwhelmingly preferred agents for lipid-lowering therapy. It is unlikely that a clinician would choose alternative therapies, even in the presence of an *APOE* phenotype that indicates a diminished response.

None of the available evidence has provided adequate data to establish that *APOE* genotype or phenotype improves outcomes when used in clinical care.

LOW-DENSITY LIPOPROTEIN SUBCLASS AND LOW-DENSITY LIPOPROTEIN PARTICLE SIZE AND CONCENTRATION

Patients with subclass pattern B have been reported to respond more favorably to diet therapy than those with subclass pattern A.(101) Subclass pattern B has also been shown to respond more favorably to gemfibrozil and niacin, with a shift from small, dense LDL particles to larger LDL particles. While statin drugs lower the overall concentration of LDL cholesterol, there is no shift to the larger LDL particles.

Randomized and Nonrandomized Controlled Trials

Superko et al (2005) reported that the response to gemfibrozil differed in patients who had LDL subclass A compared to those who had LDL subclass B.(102) There was a greater reduction in the small, LDL levels for patients with subclass B, but this did not correlate with clinical outcomes. Another study has reported that atorvastatin treatment led to an increase in mean LDL size, while pravastatin treatment led to a decrease in LDL size.(103)

Various studies have generally confirmed that small, dense LDL is impacted preferentially by fibrate treatment (104,105,106) and possibly also by statin therapy.(104,106) However, none

demonstrated that preferentially targeting small, dense LDL leads to improved outcomes, compared with standard LDL targets widely used in clinical care.

Several trials with angiographic outcomes have examined the change in LDL particle size in relation to the angiographic progression of CAD. The 1996 Stanford Coronary Risk Intervention Project trial studied the relation between small, dense LDL and the benefit of diet, counseling, and drug therapy in patients with CAD, as identified by initial coronary angiogram.(107) Patients with subclass pattern B showed a significantly greater reduction in CAD progression than those with subclass pattern A. The 1990 Familial Atherosclerosis Treatment Study randomized patients from families with premature CAD and elevated apo B levels.(108) Change in LDL particle size correlated significantly with angiographic progression of CAD in this study.

Fewer studies have evaluated clinical outcomes in relation to LDL particle size. In the 2001 Cholesterol and Recurrent Events trial, survivors of MI with normal cholesterol levels were randomized to lipid-lowering therapy or placebo.(109) A post hoc analysis from this trial failed to demonstrate a correlation between change in particle size and treatment benefit.

Section Summary: Low-Density Lipoprotein Subclass and Low-Density Lipoprotein Particle Size and Concentration

The direct clinical application of measuring small, dense lipoprotein particles is still unclear. To improve outcomes, clinicians must have tools to translate this information into clinical practice. Such tools for linking levels of small, dense LDL to clinical decision making are currently not available. Published data are inadequate to determine how such measurements should guide treatment decisions and whether these treatment decisions result in beneficial patient outcomes.

LIPOPROTEIN (A)

There is a lack of evidence to determine whether Lp(a) can be used as a target of treatment. Several randomized studies of lipid lowering therapy have included Lp(a) measurements as an intermediate outcome. While these studies have demonstrated that Lp(a) levels are reduced in patients receiving statin therapy, the data are inadequate to demonstrate how this laboratory test can be used to improve patient management.(110,111)

Section Summary: Lipoprotein (a)

There is limited evidence on the use of Lp(a) as a treatment target for patients with hyperlipidemia. The available evidence is insufficient related to impact on clinical outcomes.

HIGH SENSITIVITY C-REACTIVE PROTEIN (HS-CRP)

The most recent guidelines (2013) from the American College of Cardiology/American Heart Association (ACC/AHA) for the assessment of cardiovascular risk state “after quantitative risk assessment, a risk-based treatment decision is uncertain, assessment of 1 or more of the following—family history, hs-CRP, CAC score, or ABI—may be considered to inform treatment decision making.” The level of evidence is rated a “B”, indicating a moderate recommendation in which certainty is based on evidence that the net benefit is moderate to substantial, or there is high certainty that the net benefit is moderate.(112)

A Writing Group convened by the AHA and the Centers for Disease Control and Prevention (2003) endorsed the optional use of hs-CRP to identify persons without known cardiovascular disease who are at intermediate risk (10 to 20% risk of coronary heart disease over the next 10 years). For these patients, the results of hs-CRP testing may help guide considerations of further evaluation (e.g. imaging, exercise testing) or therapy (e.g., drug therapies with lipid-lowering, anti-platelet, or cardio-protective agents). High-sensitivity CRP testing is not necessary in high-risk patients who have a 10-year risk of greater than 20%, as these patients already qualify for intensive medical interventions. Individuals at low-risk (less than 10% per 10 years) will be unlikely to have a high-risk (greater than 20%) identified through hs-CRP testing. The Writing group recommended screening average risk (10-year risk less than 10 %) for hs-CRP for purposes of cardiovascular risk assessment.(113)

Additionally, the Writing Group recommended that hs-CRP be performed twice, optimally 2 weeks apart, fasting or non-fasting in metabolically stable patients. Patients with an average hs-CRP level greater than 3.0 mg/dL are considered to be at high relative risk of CHD. Patients with an average hs-CRP level less than 1 mg/L are at low relative risk, and patients with an hs-CRP level between 1.0 and 3.0 mg/L are at average relative risk. If hs-CRP level is greater than 10 mg/dL, the Writing Group recommended that testing should be repeated and the patient examined for sources of infections or inflammation. The Writing group recommended against the measurement of inflammatory markers other than hs-CRP (cytokines, other acute-phase reactants) for determination of coronary risk in addition to hs-CRP.

The AHA/CDC Writing Group Statement Summary:

- hsCRP is a global indicator of future vascular events in adults without any previous history of cardiovascular disease (CVD), with acceptable precision levels down to or below 0.3 mg/L
- hsCRP enhances risk assessment and therapeutic outcomes in primary CVD prevention
- hsCRP is particularly advantageous for assessing the risk in patients with:
 - Framingham 10-year risk scores of 10%-20% and/or
 - LDL levels of <160 mg/dL
- hsCRP acts as an independent marker for evaluating the possibility of recurrent cardiac events, such as myocardial infarction or restenosis, after percutaneous coronary intervention

Buckley et al (2009) conducted a systematic review and meta-analyses of epidemiologic studies to help the U.S. Preventive Services Task Force (USPSTF) determine whether CRP level should be incorporated into guidelines for coronary and cardiovascular risk assessment in primary care. It was concluded that there is strong evidence that CRP has predictive value for cardiovascular events independent of other risk factors and that moderate evidence suggests that may improve risk stratification for those at intermediate risk.(114)

High sensitivity CRP has been added to the European Society of Cardiology's guidelines for risk stratification of patients with acute coronary syndrome (2011). The guidelines state "There is solid evidence that even among patients with troponin-negative NSTEMI-ACS, elevated levels of hsCRP (>10 mg/L) are predictive of long-term mortality (>6 months up to 4 years). The FRISC study confirmed that elevated hsCRP levels are associated with increased mortality at the time of the index event and continuously increase over 4 years."(115)

Cook et al (2006) compared risk-prediction models that include or do not include hs-CRP. The models were applied to 15,048 Women's Health Study participants who were age 45 or older

and free of cardiovascular disease and cancer at baseline. During a mean follow-up of 10 years, 390 women developed CVD. For accurately predicting CVD events, hs-CRP was out-matched only by older age, current smoking, and high blood pressure among traditional Framingham variables. Non-diabetic women were classified according to their 10-year risk for CVD in a model without CRP. Adding CRP to the model substantially improved predictive accuracy for women with an initial 10-year CVD risk of at least 5%. The gain in accuracy was greatest among women who were initially classified in the 5% to 9.9% risk range: 21.3 % of those women were reclassified in a more accurate risk category when CRP was included in the risk prediction model (11.9% moved down a risk category (to less than 5 %) and 9.5 % moved up a risk category (to 10 % to 19.9%)). Accounting for the predictive value of older age, smoking, and high BP lessened the predictive contribution of CRP but still left CRP ahead of any cholesterol parameter (total, LDL, or HDL).(116)

Evidence from the JUPITER trial (2009) suggests that, for people choosing to start statin therapy, reduction in both LDL cholesterol and hs-CRP are indicators of successful treatment with statins. In an analysis of 15,548 initially healthy men and women participating in the JUPITER trial (87 % of full cohort), investigators prospectively assessed the effects of rosuvastatin versus placebo on rates of non-fatal myocardial infarction, non-fatal stroke, admission for unstable angina, arterial revascularization, or cardiovascular death during a maximum follow-up of 5 years (median of 1.9 years). Compared with placebo, participants allocated to rosuvastatin who achieved LDL cholesterol less than 1.8 mmol/L had a 55 % reduction in vascular events, and those achieving hs-CRP less than 2 mg/L a 62 % reduction. Although LDL cholesterol and hs-CRP reductions were only weakly correlated in individual patients (r values < 0.15), the investigators reported a 65 % reduction in vascular events in participants allocated to rosuvastatin who achieved both LDL cholesterol less than 1.8 mmol/L and hs-CRP less than 2 mg/L, versus a 33 % reduction in those who achieved 1 or neither target. In participants who achieved LDL cholesterol less than 1.8 mmol/L and hs-CRP less than 1 mg/L, the investigators found a 79 % reduction. The investigators reported that achieved hs-CRP concentrations were predictive of event rates irrespective of the lipid endpoint used, including the apolipoprotein B to apolipoprotein AI ratio.(117)

A meta-analysis found that hs-CRP concentration has continuous associations with the risk of coronary heart disease, ischemic stroke, and vascular mortality (Emerging Risk Factors Collaboration, 2010). Investigators assessed the associations of hs-CRP concentration with risk of vascular and non-vascular outcomes under different circumstances. Investigators meta-analyzed individual records of 160,309 people without a history of vascular disease from 54 long-term prospective studies. Within-study regression analyses were adjusted for within-person variation in risk factor levels. The investigators found that $\log(e)$ hs-CRP concentration was linearly associated with several conventional risk factors and inflammatory markers, and nearly log-linearly with the risk of ischemic vascular disease and non-vascular mortality. Risk ratios (RRs) for coronary heart disease per 1 standard deviation higher $\log(e)$ hs-CRP concentration (3 -fold higher) were 1.63 (95 % confidence interval (CI): 1.51 to 1.76) when initially adjusted for age and sex only, and 1.37 (1.27 to 1.48) when adjusted further for conventional risk factors; 1.44 (1.32 to 1.57) and 1.27 (1.15 to 1.40) for ischemic stroke; 1.71 (1.53 to 1.91) and 1.55 (1.37 to 1.76) for vascular mortality; and 1.55 (1.41 to 1.69) and 1.54 (1.40 to 1.68) for non-vascular mortality. The investigators noted that RRs were largely unchanged after exclusion of smokers or initial follow-up. After further adjustment for fibrinogen, the corresponding RRs were 1.23 (1.07 to 1.42) for coronary heart disease; 1.32 (1.18 to 1.49) for ischemic stroke; 1.34 (1.18 to 1.52) for vascular mortality; and 1.34 (1.20 to 1.50) for non-vascular mortality. The investigators concluded that hs-CRP concentration has

continuous associations with the risk of coronary heart disease, ischemic stroke, vascular mortality, and death from several cancers and lung disease that are each of broadly similar size. The investigators noted that the relevance of hs-CRP to such a range of disorders is unclear. The investigators found that associations with ischemic vascular disease depend considerably on conventional risk factors and other markers of inflammation.(118)

The 2009 Guidelines from the Canadian Cardiovascular Society state the measurement of hs-CRP is recommended in men older than 50 years and women older than 60 years of age who are at intermediate risk (10% to 19%) according to their Framingham risk score and who do not otherwise qualify for lipid-lowering therapy (i.e., if their LDL-C is less than 3.5 mmol/L). The guidelines explain that the rationale for measuring hs-CRP specifically in these individuals is that there is now class I evidence for the benefit of statin therapy in such individuals with hs-CRP greater than 2.0 mg/L. The guidelines found that data from the JUPITER study show that statin therapy reduces cardiovascular events (hazard ratio 0.56 [95% CI 0.46 to 0.69]; $P < 0.00001$). The guidelines note that because hs-CRP can be elevated during acute illness, clinical judgment should be exercised in the interpretation of any single measurement of hs-CRP. The 2012 Canadian Cardiovascular Society guidelines state that those individuals who meet JUPITER criteria (men > than 50 years and women > than 60 years of age and CRP greater than or equal to 2 mg/L and LDL greater than 3.5 mmol/L) could be considered for treatment based on the results of that study.(119)

An American Heart Association (AHA) statement on nontraditional risk factors and biomarkers in cardiovascular disease in youth states:(120) "There currently is no clinical role for measuring CRP routinely in children when assessing or considering therapy for CVD risk factors." The AHA statement explains that, although numerous studies suggest that CRP is elevated in children with higher CVD risk, correlates with the progression of atherosclerotic changes, and tracks, albeit weakly, over 21 years from childhood to adulthood independently of other metabolic and conventional cardiovascular risk factors, it is not yet clear whether high CRP levels during childhood and adolescence lead to an increased risk of CVD in later life. The AHA stated that lifestyle interventions have been shown to decrease CRP in children, and statins reduce CRP in adults. "However, minimal information is available on the effect of statins on CRP in children and youth and, importantly whether lowering CRP in children per se would modify preclinical disease or CVD outcomes."

The American Association of Clinical Endocrinologists (AACE) (2012) issued a 2b recommendation for the use of hs-CRP to stratify CVD risk in patients with a standard risk assessment that is borderline, or in those with an LDL-C concentration less than 130 mg/dL.(121)

Section Summary: hs-CRP

Several studies have shown that hs-CRP adds to the predictive capability of other established risk factors for CVD. The best evidence to date supports hs-CRP as an independent factor in predicting increased CVD risk. Although evidence for benefit or harm is not decidedly conclusive (Quality of Evidence IIb, Level C) for using hs-CRP as a marker in all patient groups as part of a global risk assessment, the AHA/CDC Writing Group endorses (at Evidence Level B) the use of hs-CRP as an adjunct to the major risk factors to assess and identify patients who may be at higher risk for CVD than estimated by major risk factors.

Supplemental Information

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

PRACTICE GUIDELINES AND POSITION STATEMENTS

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

National Heart, Lung, and Blood Institute

In 2001, the National Heart, Lung, and Blood Institute's National Cholesterol Education Program Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) issued a position statement.⁽¹⁾ Apo lipoprotein B (apo B), apolipoprotein AI (apo AI), lipid subclass, and lipoprotein(a) (Lp[a]) were listed as "emerging risk factors" for cardiovascular risk assessment, without specific recommendations for how these measures should be used in clinical practice. A 2004 update to these guidelines discussed the result of clinical trials of statin therapy.⁽¹²²⁾

In 2013, the Institute published a systematic evidence review on managing blood cholesterol in adults.⁽¹²³⁾ The review was used to develop joint guidelines by the American College of Cardiology (ACC) and American Heart Association (AHA) on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults (see below).⁽¹²⁴⁾

American College of Cardiology (ACC) / American Heart Association (AHA)

The March 2026 [ACC/AHA Guideline on the Management of Dyslipidemia](#) offers a comprehensive "one-stop shop" for addressing the evaluation, management and monitoring of individuals with dyslipidemias, including high blood cholesterol, hypertriglyceridemia, and elevated lipoprotein(a) (Lp[a]).

- Lp(a) should be measured at least once in adulthood. According to the guideline, lifestyle changes minimally affect Lp(a) levels, so repeat testing is generally not needed.
- Measuring Apolipoprotein B (apo B) may be used to assess any residual ASCVD risk and guide treatment among people with cardiovascular-kidney-metabolic syndrome, type 2 diabetes, high triglycerides or known cardiovascular disease who have reached their LDL-C and non-HDL-C goals. ApoB may be a more accurate risk marker than LDL-C in these groups of people.

American College of Cardiology

In 2023, the American College of Cardiology (ACC) published an update on lipoprotein(a) testing guidelines, which included the following:

- Lipoprotein(a) (Lp[a]) is an independent risk factor for atherosclerotic cardiovascular disease (CVD) and calcific valvular aortic stenosis.
- Lp(a) exhibits significant race/ethnic variations, with levels highest among persons of African ancestry.
- Lp(a) levels typically do not change after 5 years of age except during times of significant inflammation, liver disease, or kidney disease; hence, levels should be interpreted cautiously during these times.
- Current guidelines support once-in-a-lifetime measurement in most individuals with increased risk of atherosclerotic CVD.

- Emerging data appear to show a strong correlation with high-sensitivity C-reactive protein levels for predicting CVD risk. (125)

American Diabetes Association and American College of Cardiology Foundation

In 2008, a consensus statement from the American Diabetes Association and the ACC Foundation addressed lipoprotein management in patients with cardiometabolic risk.(127) This statement included specific recommendations for incorporating apo B testing into clinical care for high-risk patients and recommended that, for patients with metabolic syndrome being treated with statins, both LDL-C and apo B should be used as treatment targets, with an apo B target of less than 90 mg/dL, even if target LDL has been achieved.

This consensus statement also commented on the use of LDL particle number in patients with cardiometabolic risk and on the limitations of the clinical utility of nuclear magnetic resonance measurement of LDL particle number or size, including lack of widespread availability. The statement also noted that there is a need for more independent data confirming the accuracy of the method and whether its predictive power is consistent across various patient populations.

The American Diabetes Association 2024 Standards of Care do not discuss the use of specific novel biomarkers for cardiovascular disease and risk management.(128)

American Association of Clinical Endocrinologists and American College of Endocrinology

In 2017, the American Association of Clinical Endocrinologists (AACE, now the American Association of Clinical Endocrinology) and the American College of Endocrinology (ACE) published joint guidelines on the management of dyslipidemia and the prevention of cardiovascular diseases.(129) The guidelines recommended that, among patients with “triglyceride (TG) concentration of greater than 150 mg/dL or HDL-C concentration of less than 40 mg/dL, the apo B or the apo B to apo A1 ratio may be useful in assessing residual risk in individuals at risk for ASCVD (even when the LDL-C levels are controlled).”

In 2020, the AACE published an updated consensus statement on dyslipidemia and prevention of cardiovascular disease.(130) They recommended measurement of Lp(a) in several patient populations including those with ASCVD, those with a family history of premature ASCVD and/or increased Lp(a), and individuals with a 10-year ASCVD risk of 10% or greater. Recommendations also included consideration of apo B or LDL particle measurement “based on individual patient clinical circumstances.”

In 2022, the AACE published a guideline on comprehensive care plans in patients with diabetes.(131) In addition to treatment targets for LDL-C and non-HDL-C, the guideline defines target apo B levels of <90 mg/dL, <80 mg/dL, or <70 mg/dL for patients with high, very high, and extreme risk of ASCVD. Patients receiving statins should undergo monitoring for these parameters (including apo B) every 6 to 12 weeks, and monitoring frequency can decrease after targets are achieved.

National Lipid Association

In 2019, the National Lipid Association issued a scientific statement on the use of Lp(a), which notes that Lp(a) measurement “is reasonable” to refine risk assessment for ASCVD events in the following populations: patients with first-degree relatives with premature ASCVD (<55 years of age for men; <65 years of age for women), patients with premature ASCVD without

traditional risk factors, patients with severe hypercholesterolemia (LDL-C $\geq$ 190 mg/dL) or familial hypercholesterolemia, and patients with very-high risk of ASCVD that may be candidates for proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor therapy.(132) Additionally Lp(a) "may be reasonable" to measure in patients with the following: intermediate (7.5 to 19.9%) or borderline (5 to 7.4%) ASCVD risk when statin initiation is uncertain for primary prevention, inadequate response to LDL-C lowering therapy despite adherence, family history of elevated Lp(a), calcific valvular aortic stenosis, or recurrent or progressive ASCVD despite lipid-lowering therapy.

In 2021, the National Lipid Association issued a scientific statement on lipid measurements in cardiovascular disease including information on apo B, small dense LDL, and Lp(a).(133) The authors refer to the 2019 statement for information on Lp(a), and they recommend that measurements of apo B and small dense LDL "may be reasonable at initial evaluation." Additionally, apo B measurement "is reasonable" for patients receiving lipid lowering therapy while small dense LDL measurement is "not recommended" for these patients.

U.S. PREVENTIVE SERVICES TASK FORCE RECOMMENDATIONS

The U.S. Preventive Services Task Force (2009) issued recommendations on the use of nontraditional risk factors for the assessment of coronary heart disease (CHD).(134) The Task Force included lipoprotein (a) in its summary statement: "The evidence is insufficient to assess the balance of benefits and harms of using the nontraditional risk factors discussed in this statement to screen asymptomatic men and women with no history of CHD to prevent CHD events."

The recommendation was updated in 2018 and came to the same conclusion: evidence is insufficient to assess the benefits and harms of novel testing methods to diagnose CVD. However, the nontraditional risk factors included in this recommendation were different than those in this evidence review.(135)

Regulatory Status

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests (LDTs) must meet the general regulatory standards of the Clinical Improvement Act of 1988 (CLIA). Lipid and non-lipid biomarker tests are available under the auspices of the CLIA. Laboratories that offer laboratory-developed tests must be licensed by the CLIA for high-complexity testing. To date, the U.S. Food and Drug Administration has chosen not to require any regulatory review of these tests.

Government Regulations

National:

No NCD specifically addresses nontraditional biomarkers for cardiovascular disease.

NCD 190.23 covers lipid panel testing for symptomatic patients for evaluating atherosclerotic CV disease, to monitor the progress of patients on anti-lipid dietary management and pharmacologic therapy for various lipid disorders.

Please refer to the following for coverage of lipid testing:

National Coverage Determination (NCD) for Lipid Testing (190.23)

Effective Date of this Version 1/1/05, Implementation Date 3/11/05

Local:

Wisconsin Physician Services Local Coverage Determination

LCD Title: MoIDX: Biomarkers in Cardiovascular Risk Assessment (L36523)

Original Effective Date: For services performed on or after 06/16/2016

Revision Effective Date: For services performed on or after 02/26/2026

(The following are selected excerpts from the LCD, and not the entire LCD)

Under preventative services, Medicare Part B covers the basic lipid panel (total cholesterol, high density lipoprotein-cholesterol (HDL-C), triglycerides, and low-density lipoprotein-cholesterol (LDL-C) for cardiovascular (CV) disease screening, every 5 years when ordered by a doctor.

This policy denies coverage for **all CV risk assessment panels**, except the basic lipid panel, for symptomatic (with signs and symptoms) patients with suspected or documented CV disease because panel testing is not specific to a given patient's lipid abnormality or disease.

High-sensitivity C-reactive protein (hs-CRP)

1. Patient has intermediate CV risk (10-20% risk of CVD per 10 years using the Framingham point score);and
2. Patient has LDL-C between 100-130 mg/dL; and
3. Patient has two or more CHD major risk factors, including
 - Age (Men > 50 years; Women > 60 years)
 - Current cigarette smoking
 - Family history of premature CHD (CHD in male first degree relative <55; CHD in female first degree relative <65 years of age)
 - Hypertension (Systolic > 140 mm Hg, or on anti-hypertensive medication)
 - Low HDL-C (<40 mg/dL)

The use of hs-CRP testing to evaluate the effects of treatment or to motivate patients to improve lifestyle behaviors are not considered medically reasonable and necessary, and therefore not covered by Medicare.

Lipoprotein(a) (Lp(a))

Lp(a) testing may be indicated in select patients, particularly intermediate risk patients, to assist physicians with the use of preventive therapies. Routine testing is not covered by Medicare.

There are fees found on the 2025 2nd Quarter CMS Clinical Laboratory Fee Schedule for codes 82172, 83695 and 86141. An assigned fee is not a guarantee of payment.

(The above Medicare information is current as of the review date for this policy. However, the coverage issues and policies maintained by the Centers for Medicare & Medicaid Services [CMS, formerly HCFA] are updated and/or revised periodically. Therefore, the most current CMS information may not be contained in this document. For the most current information, the reader should contact an official Medicare source.)

Related Policies

- Cardiovascular Risk Panels
 - Genetic Testing – Gene Expression Testing in the Evaluation of Patients with Stable Ischemic Heart Disease (Retired 2020)
 - Measurement of Lipoprotein-associated Phospholipase A₂ (Lp-PLA₂) and Secretory Type II Phospholipase A₂ (sPLA₂-IIA) in the Assessment of Cardiovascular Risk
 - Myeloperoxidase (MPO) Immunoassay for Cardiac Disease Risk
-

References

1. Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). JAMA. May 16 2001;285(19):2486-2497. PMID 11368702
2. Brunzell JD, Davidson M, Furberg CD, et al. Lipoprotein management in patients with cardiometabolic risk: consensus statement from the American Diabetes Association and the American College of Cardiology Foundation. Diabetes Care. Apr 2008;31(4):811-822. PMID 18375431
3. Cole, Thomas G. et al. Association of Apolipoprotein B and Nuclear Magnetic Resonance Spectroscopy-Derived LDL Particle Number with Outcomes in 25 Clinical Studies: Assessment by the AACC Lipoprotein and Vascular Diseases Division Working Group on Best Practices. Clinical Chemistry, 59 (5) p.752-70, May 2013.
4. Jellinger PS, Handelsman Y, Rosenblit PD, et al. American Association of Clinical Endocrinologists and American College of Endocrinology guidelines for management of dyslipidemia and prevention of cardiovascular disease. Endocr Pract. Apr 2017;23(Suppl 2):1-87. PMID 28437620
5. Wilson PW, Jacobson TA, Martin SS, et al. Lipid measurements in the management of cardiovascular diseases: Practical recommendations a scientific statement from the national lipid association writing group. J Clin Lipidol. Published online: September 24, 2021 [https://www.lipidjournal.com/article/S1933-2874\(21\)00245-2/fulltext](https://www.lipidjournal.com/article/S1933-2874(21)00245-2/fulltext) Accessed August 3, 2026.
6. Di Angelantonio E, Gao P, Pennells L, et al. Emerging Risk Factors Collaboration: Lipid-related markers and cardiovascular disease prediction. JAMA. Jun 20 2012;307(23):2499-2506. PMID 22797450
7. Pearson GJ, Thanassoulis G, Anderson TJ, et al. 2021 Canadian Cardiovascular Society Guidelines for the Management of Dyslipidemia for the Prevention of Cardiovascular Disease in Adults. Volume 37, Issue 8, P1129-1150, August 2021. [https://onlinecjc.ca/article/S0828-282X\(21\)00165-3/fulltext](https://onlinecjc.ca/article/S0828-282X(21)00165-3/fulltext)
8. Medical Services Commission. Cardiovascular disease-primary prevention. Victoria, BC: British Columbia Medical Services Commission; March 15, 2008, revised 2014. <https://www2.gov.bc.ca/assets/gov/health/about-bc-s-health-care-system/bc-guidelines/cvd.pdf> Accessed August 3, 2026.
9. Ridker PM, Rifai N, Cook NR, et al. Non-HDL cholesterol, apolipoproteins A-I and B100, standard lipid measures, lipid ratios, and CRP as risk factors for cardiovascular disease in women. JAMA. Jul 20 2005;294(3):326-333. PMID 16030277
10. Mach R, Baigent C, Catapano AL, et al. 2019 ESC/EAS Guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk: The Task Force for the management of dyslipidaemias of the European Society of Cardiology (ESC) and

European Atherosclerosis Society (EAS), *European Heart Journal*, Volume 41, Issue 1, 1 January 2020, Pages 111–188, <https://doi.org/10.1093/eurheartj/ehz455>

11. Benn M, Nordestgaard BG, Jensen GB, et al. Improving prediction of ischemic cardiovascular disease in the general population using apolipoprotein B: the Copenhagen City Heart Study. *Arterioscler Thromb Vasc Biol*. Mar 2007;27(3):661-670. PMID 17170368
12. Robinson JG, Wang S, Jacobson TA. Meta-analysis of comparison of effectiveness of lowering apolipoprotein B versus low-density lipoprotein cholesterol and nonhigh-density lipoprotein cholesterol for cardiovascular risk reduction in randomized trials. *Am J Cardiol*. Nov 15 2012;110(10):1468-1476. PMID 22906895
13. McQueen, Matthew J. et al. Lipids, lipoproteins, and apolipoproteins as risk markers of myocardial infarction in 52 countries (the INTERHEART study): a case-control study. *Lancet* 2008; 372: 224–33.
14. Sniderman AD, Islam S, Yusuf S, et al. Discordance analysis of apolipoprotein B and non-high density lipoprotein cholesterol as markers of cardiovascular risk in the INTERHEART study. *Atherosclerosis*. Dec 2012;225(2):444-449. PMID 23068583
15. Sniderman, Allan et al. A Meta-Analysis of Low-Density Lipoprotein Cholesterol, Non-High-Density Lipoprotein Cholesterol, and Apolipoprotein B as Markers of Cardiovascular Risk. *Circ Cardiovasc Qual Outcomes*, 2011;4:337-345; originally published online April 12, 2011.
16. Gotto AM, Jr., Whitney E, Stein EA, et al. Relation between baseline and on-treatment lipid parameters and first acute major coronary events in the Air Force/Texas Coronary Atherosclerosis Prevention Study (AFCAPS/TexCAPS). *Circulation*. Feb 8 2000;101(5):477-484. PMID 10662743
17. Kastelein JJ, van der Steeg WA, Holme I, et al. Lipids, apolipoproteins, and their ratios in relation to cardiovascular events with statin treatment. *Circulation*. Jun 10 2008;117(23):3002-3009. PMID 18519851
18. Mora S, Wenger NK, Demicco DA, et al. Determinants of residual risk in secondary prevention patients treated with high- versus low-dose statin therapy: the Treating to New Targets (TNT) study. *Circulation*. Apr 24 2012;125(16):1979-1987. PMID 22461416
19. Clarke R, Emberson JR, Parish S, et al. Cholesterol fractions and apolipoproteins as risk factors for heart disease mortality in older men. *Arch Intern Med*. Jul 9 2007;167(13):1373-1378. PMID 17620530
20. Ridker PM, Buring JE, Rifai N, et al. Development and validation of improved algorithms for the assessment of global cardiovascular risk in women: the Reynolds Risk Score. *JAMA*. Feb 14 2007;297(6):611-619. PMID 17299196
21. van der Steeg WA, Boekholdt SM, Stein EA, et al. Role of the apolipoprotein B-apolipoprotein A-I ratio in cardiovascular risk assessment: a case-control analysis in EPIC-Norfolk. *Ann Intern Med*. May 1 2007;146(9):640-648. PMID 17470832
22. Walldius G, Jungner I, Holme I, et al. High apolipoprotein B, low apolipoprotein A-I, and improvement in the prediction of fatal myocardial infarction (AMORIS study): a prospective study. *Lancet*. Dec 15 2001;358(9298): 2026-33. PMID 11755609
23. Bennet AM, Di Angelantonio E, Ye Z, et al. Association of apolipoprotein E genotypes with lipid levels and coronary risk. *JAMA*. Sep 19 2007;298(11):1300-1311. PMID 17878422
24. Sofat R, Cooper JA, Kumari M, et al. Circulating apolipoprotein E concentration and cardiovascular disease risk: meta-analysis of results from three studies. *PLoS Med*. Oct 2016;13(10):e1002146. PMID 27755538

25. Koch W, Hoppmann P, Schomig A, et al. Apolipoprotein E gene epsilon2 /epsilon3/epsilon4 polymorphism and myocardial infarction: case-control study in a large population sample. *Int J Cardiol.* Mar 28 2008; 125(1): 116-7. PMID 17433475
26. Kulminski AM, Ukraintseva SV, Arbeev KG, et al. Health-protective and adverse effects of the apolipoprotein E epsilon2 allele in older men. *J Am Geriatr Soc.* Mar 2008; 56(3): 478-83. PMID 18179501
27. Schmitz F, Mevissen V, Krantz C, et al. Robust association of the APOE epsilon4 allele with premature myocardial infarction especially in patients without hypercholesterolaemia: the Aachen study. *Eur J Clin Invest.* Feb 2007;37(2): 106-8. PMID 17217375
28. Vaisi-Raygani A, Rahimi Z, Nomani H, et al. The presence of apolipoprotein epsilon4 and epsilon2 alleles augments the risk of coronary artery disease in type 2 diabetic patients. *Clin Biochem.* Oct 2007; 40(15): 1150-6. PMID 17689519
29. Ciftdoğan DY, Coskun S, Ulman C, et al. The association of apolipoprotein E polymorphism and lipid levels in children with a family history of premature coronary artery disease. *J Clin Lipidol.* Jan-Feb 2012; 6(1): 81-7. PMID 22264578
30. Vasunilashorn S, Gleib DA, Lan CY, et al. Apolipoprotein E is associated with blood lipids and inflammation in Taiwanese older adults. *Atherosclerosis.* Nov 2011; 219(1): 349-54. PMID 21840004
31. de Andrade M, Thandi I, Brown S, et al. Relationship of the apolipoprotein E polymorphism with carotid artery atherosclerosis. *Am J Hum Genet.* Jun 1995; 56(6): 1379-90. PMID 7762561
32. Eichner JE, Kuller LH, Orchard TJ, et al. Relation of apolipoprotein E phenotype to myocardial infarction and mortality from coronary artery disease. *Am J Cardiol.* Jan 15 1993; 71(2): 160-5. PMID 8421977
33. Wilson PW, Myers RH, Larson MG, et al. Apolipoprotein E alleles, dyslipidemia, and coronary heart disease. The Framingham Offspring Study. *JAMA.* Dec 07 1994; 272(21): 1666-71. PMID 7966894
34. Wilson PW, Schaefer EJ, Larson MG, et al. Apolipoprotein E alleles and risk of coronary disease. A meta-analysis. *Arterioscler Thromb Vasc Biol.* Oct 1996; 16(10): 1250-5. PMID 8857921
35. Sharrett AR, Ballantyne CM, Coady SA, et al. Coronary heart disease prediction from lipoprotein cholesterol levels, triglycerides, lipoprotein(a), apolipoproteins A-I and B, and HDL density subfractions: The Atherosclerosis Risk in Communities (ARIC) Study. *Circulation.* Sep 4 2001;104(10):1108-1113. PMID 11535564
36. Volcik KA, Barkley RA, Hutchinson RG, et al. Apolipoprotein E polymorphisms predict low density lipoprotein cholesterol levels and carotid artery wall thickness but not incident coronary heart disease in 12,491 ARIC study participants. *Am J Epidemiol.* Aug 15 2006;164(4):342-348. PMID 16760224
37. Singh K, Chandra A, Sperry T, et al. Associations Between High-Density Lipoprotein Particles and Ischemic Events by Vascular Domain, Sex, and Ethnicity: A Pooled Cohort Analysis. *Circulation.* Aug 18 2020; 142(7): 657-669. PMID 32804568
38. Mora S, Glynn RJ, Ridker PM. High-density lipoprotein cholesterol, size, particle number, and residual vascular risk after potent statin therapy. *Circulation.* Sep 10 2013;128(11): 1189-1197. PMID 24002795
39. Stampfer MJ, Krauss RM, Ma J, et al. A prospective study of triglyceride level, low-density lipoprotein particle diameter, and risk of myocardial infarction. *JAMA.* Sep 18 1996;276(11):882-888. PMID 8782637

40. Lamarche B, Moorjani S, Lupien PJ, et al. Apolipoprotein A-I and B levels and the risk of ischemic heart disease during a five-year follow-up of men in the Québec cardiovascular study. *Circulation*. Aug 01 1996; 94(3): 273-8. PMID 8759066
41. Lamarche B, Tchernof A, Moorjani S, et al. Small, dense low-density lipoprotein particles as a predictor of the risk of ischemic heart disease in men. Prospective results from the Quebec Cardiovascular Study. *Circulation*. Jan 7 1997;95(1):69-75. PMID 8994419
42. Tzou WS, Douglas PS, Srinivasan SR, et al. Advanced lipoprotein testing does not improve identification of subclinical atherosclerosis in young adults: the Bogalusa Heart Study. *Ann Intern Med*. May 3 2005;142(9):742-750. PMID 15867406
43. Blake GJ, Otvos JD, Rifai N, et al. Low-density lipoprotein particle concentration and size as determined by nuclear magnetic resonance spectroscopy as predictors of cardiovascular disease in women. *Circulation*. Oct 8 2002;106(15):1930-1937. PMID 12370215
44. Kuller L, Arnold A, Tracy R, et al. Nuclear magnetic resonance spectroscopy of lipoproteins and risk of coronary heart disease in the Cardiovascular Health Study. *Arterioscler Thromb Vasc Biol*. Jul 1 2002;22(7):1175-1180. PMID 12117734
45. Rosenson RS, Otvos JD, Freedman DS. Relations of lipoprotein subclass levels and low-density lipoprotein size to progression of coronary artery disease in the Pravastatin Limitation of Atherosclerosis in the Coronary Arteries (PLAC-I) trial. *Am J Cardiol*. Jul 15 2002;90(2):89-94. PMID 12106834
46. Otvos JD, Jeyarajah EJ, Cromwell WC. Measurement issues related to lipoprotein heterogeneity. *Am J Cardiol*. Oct 17 2002;90(8A):22i-29i. PMID 12419478
47. Rosenson RS, Underberg JA. Systematic review: Evaluating the effect of lipid-lowering therapy on lipoprotein and lipid values. *Cardiovasc Drugs Ther*. Jul 28 2013;27(5):465-479. PMID 23893306
48. Mora S, Otvos JD, Rifai N, et al. Lipoprotein particle profiles by nuclear magnetic resonance compared with standard lipids and apolipoproteins in predicting incident cardiovascular disease in women. *Circulation*. Feb 24 2009;119(7):931-939. PMID 19204302
49. Toth PP, Grabner M, Punekar RS, et al. Cardiovascular risk in patients achieving low-density lipoprotein cholesterol and particle targets. *Atherosclerosis*. Aug 2014;235(2):585-591. PMID 24956532
50. Bennet A, Di Angelantonio E, Erqou S, et al. Lipoprotein(a) levels and risk of future coronary heart disease: large-scale prospective data. *Arch Intern Med*. Mar 24 2008; 168(6): 598-608. PMID 18362252
51. Smolders B, Lemmens R, Thijs V. Lipoprotein (a) and stroke: a meta-analysis of observational studies. *Stroke*. Jun 2007; 38(6): 1959-66. PMID 17478739
52. Khera AV, Everett BM, Caulfield MP, et al. Lipoprotein(a) concentrations, rosuvastatin therapy, and residual vascular risk: an analysis from the JUPITER Trial (Justification for the Use of Statins in Prevention: an Intervention Trial Evaluating Rosuvastatin). *Circulation*. Feb 11 2014; 129(6): 635-42. PMID 24243886
53. Albers JJ, Slee A, O'Brien KD, et al. Relationship of apolipoproteins A-1 and B, and lipoprotein(a) to cardiovascular outcomes: the AIM-HIGH trial (Atherothrombosis Intervention in Metabolic Syndrome with Low HDL/High Triglyceride and Impact on Global Health Outcomes). *J Am Coll Cardiol*. Oct 22 2013; 62(17): 1575-9. PMID 23973688
54. Kamstrup PR, Benn M, Tybjaerg-Hansen A, et al. Extreme lipoprotein(a) levels and risk of myocardial infarction in the general population: the Copenhagen City Heart Study. *Circulation*. Jan 15 2008; 117(2): 176-84. PMID 18086931

55. Tzoulaki I, Murray GD, Lee AJ, et al. Relative value of inflammatory, hemostatic, and rheological factors for incident myocardial infarction and stroke: the Edinburgh Artery Study. *Circulation*. Apr 24 2007; 115(16): 2119-27. PMID 17404162
56. Zakai NA, Katz R, Jenny NS, et al. Inflammation and hemostasis biomarkers and cardiovascular risk in the elderly: the Cardiovascular Health Study. *J Thromb Haemost*. Jun 2007; 5(6): 1128-35. PMID 17388967
57. Waldeyer C, Makarova N, Zeller T, et al. Lipoprotein(a) and the risk of cardiovascular disease in the European population: results from the BiomarCaRE consortium. *Eur Heart J*. Aug 21 2017; 38(32): 2490-2498. PMID28449027
58. Lee SR, Prasad A, Choi YS, et al. LPA Gene, Ethnicity, and Cardiovascular Events. *Circulation*. Jan 17 2017; 135(3): 251-263. PMID 27831500
59. Rigal M, Ruidavets JB, Viguier A, et al. Lipoprotein (a) and risk of ischemic stroke in young adults. *J Neurol Sci*. Jan 15 2007; 252(1): 39-44. PMID 17113602
60. Suk Danik J, Rifai N, Buring JE, et al. Lipoprotein(a), hormone replacement therapy, and risk of future cardiovascular events. *J Am Coll Cardiol*. Jul 08 2008; 52(2): 124-31. PMID 18598891
61. Genser B, Dias KC, Siekmeier R, et al. Lipoprotein (a) and risk of cardiovascular disease--a systematic review and meta-analysis of prospective studies. *Clin Lab*. 2011; 57(3-4): 143-56. PMID 21500721
62. Erqou S, Kaptoge S, Perry PL, et al. Lipoprotein(a) concentration and the risk of coronary heart disease, stroke, and nonvascular mortality. *JAMA*. Jul 22 2009; 302(4): 412-23. PMID 19622820
63. Vazirian F, Sadeghi M, Kelesidis T, et al. Predictive value of lipoprotein(a) in coronary artery calcification among asymptomatic cardiovascular disease subjects: A systematic review and meta-analysis. *NutrMetab Cardiovasc Dis*. Nov 2023; 33(11): 2055-2066. PMID 37567791
64. Schaefer EJ, Lamon-Fava S, Jenner JL, et al. Lipoprotein(a) levels and risk of coronary heart disease in men. The lipid Research Clinics Coronary Primary Prevention Trial. *JAMA*. Apr 06 1994; 271(13): 999-1003. PMID 8139085
65. Nestel PJ, Barnes EH, Tonkin AM, et al. Plasma lipoprotein(a) concentration predicts future coronary and cardiovascular events in patients with stable coronary heart disease. *Arterioscler Thromb Vasc Biol*. Dec 2013; 33(12): 2902-8. PMID 24092750
66. Bostom AG, Cupples LA, Jenner JL, et al. Elevated plasma lipoprotein(a) and coronary heart disease in men aged 55 years and younger. A prospective study. *JAMA*. Aug 21 1996; 276(7): 544-8. PMID 8709403
67. Ohira T, Schreiner PJ, Morrisett JD, et al. Lipoprotein(a) and incident ischemic stroke: the Atherosclerosis Risk in Communities (ARIC) study. *Stroke*. Jun 2006; 37(6): 1407-12. PMID 16675734
68. Fogacci F, Cicero AF, D'Addato S, et al. Serum lipoprotein(a) level as long-term predictor of cardiovascular mortality in a large sample of subjects in primary cardiovascular prevention: data from the Brisighella Heart Study. *Eur J Intern Med*. Jan 2017; 37: 49-55. PMID 27553697
69. Ridker PM, Hennekens CH, Stampfer MJ. A prospective study of lipoprotein(a) and the risk of myocardial infarction. *JAMA*. Nov 10 1993; 270(18): 2195-9. PMID 8411602
70. Bolibar I, von Eckardstein A, Assmann G, et al. Short-term prognostic value of lipid measurements in patients with angina pectoris. The ECAT Angina Pectoris Study Group: European Concerted Action on Thrombosis and Disabilities. *Thromb Haemost*. Dec 2000; 84(6): 955-60. PMID 11154140

71. Clarke R, Peden JF, Hopewell JC, et al. Genetic variants associated with Lp(a) lipoprotein level and coronary disease. *N Engl J Med*. Dec 24 2009; 361(26): 2518-28. PMID 20032323
72. Reyes-Soffer G, Ginsberg HN, Berglund L, et al. Lipoprotein(a): A genetically determined, causal, and prevalent risk factor for atherosclerotic cardiovascular disease: a scientific statement from the American Heart Association. *Arterioscler Thromb Vasc Biol*. 2022;42:e:48-e60.
73. Patel AP, Wang M, Pirruccello JP, et al. Lp(a) (Lipoprotein[a]) concentrations and incident atherosclerotic cardiovascular disease. *Arterioscler Thromb Vasc Biol*. 2021;42:465-474.
74. Shaw LJ, Polk DM, Kahute TA, et al. Prognostic accuracy of B-natriuretic peptide measurements and coronary artery calcium in asymptomatic subjects (from the Early Identification of Subclinical Atherosclerosis by Noninvasive Imaging Research [EISNER] study). *Am J Cardiol*. Nov 01 2009; 104(9): 1245-50. PMID 19840570
75. Wu Z, Pilbrow AP, Liew OW, et al. Circulating cardiac biomarkers improve risk stratification for incident cardiovascular disease in community dwelling populations. *EBioMedicine*. Aug 2022; 82: 104170. PMID 35850010
76. Melander O, Newton-Cheh C, Almgren P, et al. Novel and conventional biomarkers for prediction of incident cardiovascular events in the community. *JAMA*. Jul 01 2009; 302(1): 49-57. PMID 19567439
77. Wang TJ, Larson MG, Levy D, et al. Plasma natriuretic peptide levels and the risk of cardiovascular events and death. *N Engl J Med*. Feb 12 2004; 350(7): 655-63. PMID 14960742
78. Ito H, Pacold IV, Durazo-Arvizu R, et al. The effect of including cystatin C or creatinine in a cardiovascular risk model for asymptomatic individuals: the multi-ethnic study of atherosclerosis. *Am J Epidemiol*. Oct 15 2011; 174(8):949-57. PMID 21880578
79. Lee M, Saver JL, Huang WH, et al. Impact of elevated cystatin C level on cardiovascular disease risk in predominantly high cardiovascular risk populations: a meta-analysis. *Circ Cardiovasc Qual Outcomes*. Nov 2010;3(6): 675-83. PMID 20923994
80. Luo J, Wang LP, Hu HF, et al. Cystatin C and cardiovascular or all-cause mortality risk in the general population: A meta-analysis. *Clin Chim Acta*. Oct 23 2015; 450: 39-45. PMID 26192218
81. Kengne AP, Czernichow S, Stamatakis E, et al. Fibrinogen and future cardiovascular disease in people with diabetes: aetiological associations and risk prediction using individual participant data from nine community-based prospective cohort studies. *Diab Vasc Dis Res*. Mar 2013; 10(2): 143-51. PMID 22786872
82. Willeit P, Thompson SG, Agewall S, et al. Inflammatory markers and extent and progression of early atherosclerosis: Meta-analysis of individual-participant-data from 20 prospective studies of the PROG-IMT collaboration. *Eur J Prev Cardiol*. Jan 2016; 23(2): 194-205. PMID 25416041
83. Ahmadi-Abhari S, Luben RN, Wareham NJ, et al. Seventeen year risk of all-cause and cause-specific mortality associated with C-reactive protein, fibrinogen and leukocyte count in men and women: the EPIC-Norfolk study. *Eur J Epidemiol*. Jul 2013; 28(7): 541-50. PMID 23821244
84. Danesh J, Lewington S, Thompson SG, et al. Plasma fibrinogen level and the risk of major cardiovascular diseases and nonvascular mortality: an individual participant meta-analysis. *JAMA*. Oct 12 2005; 294(14): 1799-809. PMID 16219884
85. Kaptoge S, White IR, Thompson SG, et al. Associations of plasma fibrinogen levels with established cardiovascular disease risk factors, inflammatory markers, and other characteristics: individual participant meta-analysis of 154,211 adults in 31 prospective

- studies: the fibrinogen studies collaboration. *Am J Epidemiol.* Oct 15 2007; 166(8): 867-79. PMID 17785713
86. Sattar N, Wannamethee G, Sarwar N, et al. Leptin and coronary heart disease: prospective study and systematic review. *J Am Coll Cardiol.* Jan 13 2009; 53(2): 167-75. PMID 19130985
 87. Zeng R, Xu CH, Xu YN, et al. Association of leptin levels with pathogenetic risk of coronary heart disease and stroke: a meta-analysis. *Arq Bras Endocrinol Metabol.* Nov 2014; 58(8): 817-23. PMID 25465603
 88. Yang H, Guo W, Li J, et al. Leptin concentration and risk of coronary heart disease and stroke: A systematic review and meta-analysis. *PLoS One.* 2017; 12(3): e0166360. PMID 28278178
 89. Boekholdt SM, Arsenault BJ, Mora S, et al. Association of LDL cholesterol, non-HDL cholesterol, and apolipoprotein B levels with risk of cardiovascular events among patients treated with statins: a meta-analysis. *JAMA.* Mar 28 2012; 307(12): 1302-9. PMID 22453571
 90. Boekholdt SM, Hovingh GK, Mora S, et al. Very low levels of atherogenic lipoproteins and the risk for cardiovascular events: a meta-analysis of statin trials. *J Am Coll Cardiol.* Aug 05 2014; 64(5): 485-94. PMID 25082583
 91. Ballantyne CM, Pitt B, Loscalzo J, et al. Alteration of relation of atherogenic lipoprotein cholesterol to apolipoprotein B by intensive statin therapy in patients with acute coronary syndrome (from the Limiting UNdertreatment of lipids in ACS With Rosuvastatin [LUNAR] Trial). *Am J Cardiol.* Feb 15 2013; 111(4): 506-9. PMID 23237107
 92. Mora S, Glynn RJ, Boekholdt SM, et al. On-treatment non-high-density lipoprotein cholesterol, apolipoprotein B, triglycerides, and lipid ratios in relation to residual vascular risk after treatment with potent statin therapy: JUPITER (justification for the use of statins in prevention: an intervention trial evaluating rosuvastatin). *J Am Coll Cardiol.* Apr 24 2012; 59(17): 1521-8. PMID 22516441
 93. Ray KK, Cannon CP, Cairns R, et al. Prognostic utility of apoB/AI, total cholesterol/HDL, non-HDL cholesterol, orhs-CRP as predictors of clinical risk in patients receiving statin therapy after acute coronary syndromes: results from PROVE IT-TIMI 22. *Arterioscler Thromb Vasc Biol.* Mar 2009; 29(3): 424-30. PMID 19122170
 94. Osei-Hwedieh DO, Amar M, Sviridov D, et al. Apolipoprotein mimetic peptides: Mechanisms of action as anti-atherogenic agents. *Pharmacol Ther.* Apr 2011; 130(1): 83-91. PMID 21172387
 95. Ordoas JM, Mooser V. The APOE locus and the pharmacogenetics of lipid response. *Curr Opin Lipidol.* Apr 2002; 13(2): 113-7. PMID 11891412
 96. Sarkkinen E, Korhonen M, Erkkila A, et al. Effect of apolipoprotein E polymorphism on serum lipid response to the separate modification of dietary fat and dietary cholesterol. *Am J Clin Nutr.* Dec 1998; 68(6): 1215-22. PMID 9846849
 97. Carmena R, Roederer G, Mailloux H, et al. The response to lovastatin treatment in patients with heterozygous familial hypercholesterolemia is modulated by apolipoprotein E polymorphism. *Metabolism.* Jul 1993; 42(7): 895-901. PMID 8345800
 98. Chiodini BD, Franzosi MG, Barlera S, et al. Apolipoprotein E polymorphisms influence effect of pravastatin on survival after myocardial infarction in a Mediterranean population: the GISSI-Prevenzione study. *Eur Heart J.* Aug 2007; 28(16): 1977-83. PMID 17567623
 99. Donnelly LA, Palmer CN, Whitley AL, et al. Apolipoprotein E genotypes are associated with lipid-lowering responses to statin treatment in diabetes: a Go-DARTS study. *Pharmacogenet Genomics.* Apr 2008; 18(4): 279-87. PMID 18334912

100. Vossen CY, Hoffmann MM, Hahmann H, et al. Effect of APOE genotype on lipid levels in patients with coronary heart disease during a 3-week inpatient rehabilitation program. *Clin Pharmacol Ther.* Aug 2008; 84(2): 222-7. PMID 18388879
101. Kwiterovich PO. Clinical relevance of the biochemical, metabolic, and genetic factors that influence low-density lipoprotein heterogeneity. *Am J Cardiol.* Oct 17 2002; 90(8A): 30i-47i. PMID 12419479
102. Superko HR, Berneis KK, Williams PT, et al. Gemfibrozil reduces small low-density lipoprotein more in normolipemic subjects classified as low-density lipoprotein pattern B compared with pattern A. *Am J Cardiol.* Nov01 2005; 96(9): 1266-72. PMID 16253595
103. Sirtori CR, Calabresi L, Pisciotta L, et al. Effect of statins on LDL particle size in patients with familial combined hyperlipidemia: a comparison between atorvastatin and pravastatin. *Nutr Metab Cardiovasc Dis.* Feb 2005; 15(1):47-55. PMID 15871851
104. Arca M, Montali A, Pigna G, et al. Comparison of atorvastatin versus fenofibrate in reaching lipid targets and influencing biomarkers of endothelial damage in patients with familial combined hyperlipidemia. *Metabolism.* Nov2007; 56(11): 1534-41. PMID 17950105
105. Rosenson RS, Wolff DA, Huskin AL, et al. Fenofibrate therapy ameliorates fasting and postprandial lipoproteinemia, oxidative stress, and the inflammatory response in subjects with hypertriglyceridemia and the metabolic syndrome. *Diabetes Care.* Aug 2007; 30(8): 1945-51. PMID 17483155
106. Tokuno A, Hirano T, Hayashi T, et al. The effects of statin and fibrate on lowering small dense LDL- cholesterol in hyperlipidemic patients with type 2 diabetes. *J Atheroscler Thromb.* Jun 2007; 14(3): 128-32. PMID 17587764
107. Miller BD, Alderman EL, Haskell WL, et al. Predominance of dense low-density lipoprotein particles predicts angiographic benefit of therapy in the Stanford Coronary Risk Intervention Project. *Circulation.* Nov 01 1996; 94(9):2146-53. PMID 8901665
108. Brown G, Albers JJ, Fisher LD, et al. Regression of coronary artery disease as a result of intensive lipid-lowering therapy in men with high levels of apolipoprotein B. *N Engl J Med.* Nov 08 1990; 323(19): 1289-98. PMID 2215615
109. Campos H, Moye LA, Glasser SP, et al. Low-density lipoprotein size, pravastatin treatment, and coronary events. *JAMA.* Sep 26 2001; 286(12): 1468-74. PMID 11572739
110. Bays HE, Dujovne CA, McGovern ME, et al. Comparison of once-daily, niacin extended-release/lovastatin with standard doses of atorvastatin and simvastatin (the ADVICOR Versus Other Cholesterol-Modulating Agents Trial Evaluation [ADVOCATE]). *Am J Cardiol.* Mar 15 2003; 91(6): 667-72. PMID 12633795
111. van Wissen S, Smilde TJ, Trip MD, et al. Long term statin treatment reduces lipoprotein(a) concentrations in heterozygous familial hypercholesterolaemia. *Heart.* Aug 2003; 89(8): 893-6. PMID 12860867
112. Goff, DC et al. 2013 ACC/AHA Guideline on the Assessment of Cardiovascular Risk: A Report of the American College of Cardiology /American Heart Association Task Force on Practice Guidelines. *Circulation*; November 12, 2013.
113. Pearson TA et al. Markers of inflammation and cardiovascular disease. Application to clinical and public health practice. A statement for healthcare professionals from the Centers for Disease Control and Prevention and the American Heart Association. *AHA/CDC Scientific Statement, Circulation*, 2003;107:499-511.
114. Buckley DI et al. Review C-reactive protein as a risk factor for coronary heart disease: a systematic review and meta-analyses for the U.S. Preventive Services Task Force. *Ann Intern Med.* 2009 Oct 6; 151(7):483-95.
115. Hamm, Christian W. et al. ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: The Task

Force for the management of acute coronary syndromes (ACS) in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC). *European Heart Journal* (2011) 32, 2999–3054.

116. Cook, NR et al. The effect of including C-reactive protein in cardiovascular risk prediction models for women. *Ann Intern Med*. 2006 Jul 4;145(1):21-9.
117. Ridker PM, Danielson E, Fonseca FA, et al. JUPITER Trial Study Group. Reduction in C-reactive protein and LDL cholesterol and cardiovascular event rates after initiation of rosuvastatin: A prospective study of the JUPITER trial. *Lancet*, 2009;373(9670):1175-1182.
118. Di Angelantonio, E et al. The Emerging Risk Factors Collaboration: C-reactive protein concentration and risk of coronary heart disease, stroke, and mortality: an individual participant meta-analysis. *Lancet*, 2010; 375: 132–40.
119. Genest J et al. 2009 Canadian Cardiovascular Society/Canadian guidelines for the diagnosis and treatment of dyslipidemia and prevention of cardiovascular disease in the adult - 2009 recommendations. *Can J Cardiol* 2009; 25:567-79.
120. Balagopal, PB et al. Nontraditional risk factors and biomarkers for cardiovascular disease: mechanistic, research, and clinical considerations for youth: a scientific statement from the American Heart Association. *Circulation*, 2011 Jun 14;123(23):2749-69.
121. Jellinger, Paul S. et al. AACE Task Force for the Management of Dyslipidemia and Prevention of Atherosclerosis Writing Committee. AACE Lipid and Atherosclerosis Guidelines, *Endor Pract*. 2012; 18, (suppl 1), retrieved June 2, 2021 from: <https://www.aace.com/files/lipid-guidelines.pdf>.
122. Grundy SM, Cleeman JI, Merz CN, et al. Implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III guidelines. *Circulation*. Jul 13 2004; 110(2): 227-39. PMID 15249516
123. National Heart Lung and Blood Institute. Managing Blood Cholesterol in Adults: Systematic Evidence Review From the Cholesterol Expert Panel, 2013. Bethesda, MD: National Heart, Lung, and Blood Institute; 2013. <https://www.nhlbi.nih.gov/sites/default/files/media/docs/cholesterol-in-adults.pdf>. Accessed August 3, 2026.
124. Stone NJ, Robinson JG, Lichtenstein AH, et al. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. Jul 01 2014; 63(25 Pt B): 2889-934. PMID24239923
125. Alebna PL, Mehta A. An Update on Lipoprotein(a): The Latest on Testing, Treatment, and Guideline Recommendations. *Expert Analysis, American College of Cardiology*. Sep 19, 2023. <https://www.acc.org/Latest-in-Cardiology/Articles/2023/09/19/10/54/An-Update-on-Lipoprotein-a>
126. Arnett DK, Blumenthal RS, Albert MA, et al. 2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation*. Sep 10 2019; 140(11): e596-e646. PMID 30879355
127. Brunzell JD, Davidson M, Furberg CD, et al. Lipoprotein management in patients with cardiometabolic risk: consensus statement from the American Diabetes Association and the American College of Cardiology Foundation. *Diabetes Care*. Apr 2008; 31(4): 811-22. PMID 18375431

128. ElSayed NA, Aleppo G, Bannuru RR, et al. 10. Cardiovascular Disease and Risk Management: Standards of Care in Diabetes-2024. Diabetes Care. Jan 01 2024; 47(Suppl 1): S179-S218. PMID 38078592
129. Jellinger PS, Handelsman Y, Rosenblit PD, et al. American Association of Clinical Endocrinologists and American College of Endocrinology Guidelines for Management of Dyslipidemia and Prevention of Cardiovascular Disease. Endocr Pract. Apr 2017; 23(Suppl 2):1-87. PMID 28437620
130. Handelsman Y, Jellinger PS, Guerin CK, et al. Consensus Statement by the American Association of Clinical Endocrinologists and American College of Endocrinology on the Management of Dyslipidemia and Prevention of Cardiovascular Disease Algorithm - 2020 Executive Summary. Endocr Pract. Oct 2020; 26(10): 1196-1224. PMID33471721
131. Blonde L, Umpierrez GE, Reddy SS, et al. American Association of Clinical Endocrinology Clinical Practice Guideline: Developing a Diabetes Mellitus Comprehensive Care Plan-2022 Update. Endocr Pract. Oct 2022; 28(10): 923-1049. PMID 35963508
132. Wilson DP, Jacobson TA, Jones PH, et al. Use of Lipoprotein(a) in clinical practice: A biomarker whose time has come. A scientific statement from the National Lipid Association. J Clin Lipidol. May 2019; 13(3): 374-392. PMID31147269
133. Wilson PW, Jacobson TA, Martin SS, et al. Lipid measurements in the management of cardiovascular diseases: Practical recommendations a scientific statement from the national lipid association writing group. J Clin Lipidol. Published online: September 24, 2021
134. Helfand M, Buckley DI, Freeman M, et al. Emerging risk factors for coronary heart disease: a summary of systematic reviews conducted for the U.S. Preventive Services Task Force. Ann Intern Med. Oct 06 2009; 151(7):496-507. PMID 19805772
135. Curry SJ, Krist AH, Owens DK, et al. Risk Assessment for Cardiovascular Disease With Nontraditional Risk Factors: US Preventive Services Task Force Recommendation Statement. JAMA. Jul 17 2018; 320(3): 272-280.PMID 29998297

The articles reviewed in this research include those obtained in an Internet based literature search for relevant medical references through August 3, 2026, the date the research was completed.

Joint BCBSM/BCN Medical Policy History

Policy Effective Date	BCBSM Signature Date	BCN Signature Date	Comments
3/1/12	10/11/11	11/9/11	Joint policy established. Consolidated previous policies, “Low Density Lipoprotein (LDL) Particle Size” and “High Density Lipoprotein Subclass Testing in the Diagnosis and Management of Cardiovascular Disease”. These two policies will be moved out of the active policies queue.
3/1/13	12/11/12	12/31/12	Routine maintenance. Added new code, 81401; Revised language in policy statement. Policy title changed from “Lipid Risk Factors in the Assessment and Management of Cardiovascular Disease” to “Novel Lipid Risk Factors in the Risk Assessment and Management of Cardiovascular Disease”.
7/1/14	4/8/14	4/15/14	Policy extensively rewritten and intent changed; added apo B and hs-CRP as established; title changed from “Novel Lipid Risk Factors in the Risk Assessment and Management of Cardiovascular Disease” to current title; references updated
11/1/15	8/18/15	9/16/15	Added description and rationale information on genetic biomarkers for CVD, FH (heterozygous and homozygous) Added code 81401, as this describes <i>APOB</i> molecular testing Added code 81405, as this describes <i>LDLR</i> molecular testing Added code 81406, used for <i>PCSK9</i> molecular testing Added 84999, may be used for <i>ARH</i> adaptor protein testing Added <i>LDLR</i> , <i>APOB</i> , <i>PCSK9</i> and <i>ARH</i> adaptor protein (<i>LDLRAP1</i>) to genetic tests established for FH. Added to MPS: “The safety and effectiveness of genetic testing of <i>LDLR</i> , <i>APOB</i> , <i>PCSK9</i> and <i>ARH</i> adaptor protein (<i>LDLRAP1</i>) have been established. It may be a useful

			diagnostic option for individuals with suspected homozygous familial hypercholesterolemia, in whom the diagnosis is uncertain.” Revised MPS to include all biomarkers noted in policy. Added “APOE genotyping” to exclusions and MPS (literature does not support) Added to Exclusions: “Laboratory testing of other novel biomarkers to assess cardiovascular risk, including but not limited to apolipoprotein AI, apolipoprotein E or APOE genotypes, LDL subclass, HDL subclass, lipoprotein[a], Cystatin C, brain natriuretic peptide (BNP), fibrinogen, and leptin.”
11/1/16	8/16/16	8/16/16	Routine maintenance Literature review Added code 82397, may be used for leptin testing Added code 82664, may be used for HDL subclass Added code 83520, may be used for leptin testing Added code 83700, classified for lipoprotein, blood; electrophoretic separation and quantitation Added code 83721, classified for Lipoprotein, direct measurement; LDL cholesterol Added code 84181, may be used for APOE phenotyping or genotyping Added code 85385, classified for Fibrinogen, antigen LCD update PULS cardiac testing added to MPS
11/1/17	8/15/17	8/15/17	Routine maintenance
11/1/18	8/21/18	8/21/18	• Routine maintenance • Removed homozygous familial hypercholesterolemia from policy • 0055U added to EI
1/1/19	10/16/18	10/16/18	• 0055U removed • 0052U added
1/1/20	10/15/19		• Routine maintenance • 83722 added per code update EI
11/1/20	8/18/20		• Routine maintenance
11/1/21	8/17/21		• Routine maintenance

11/1/22	8/16/22		• Routine maintenance • Lipoprotein(a) (83695) moved from E/I to EST • Apo B and hs-CRP criteria aligned with avalon policy • Ref added 135,136
11/1/23	8/15/23		• Routine maintenance • Vendor: Avalon (ky)
11/1/24	8/20/24		• Routine maintenance • Use of "novel" edited to "nontraditional" in policy statements; intent unchanged. • Title changed from Novel Biomarkers in Risk Assessment and Management of Cardiovascular Disease to Nontraditional Biomarkers in Risk Assessment and Management of Cardiovascular Disease. • Vendor: Avalon (ky)
11/1/25	8/19/25		• Routine maintenance • Formatting updated • Ref 63,125,128 added • PULS cardiac test removed • Vendor: Avalon (ls)
11/1/26	8/18/26		• Routine maintenance • Vendor: Avalon • Lp (a) and hs-CRP criteria updated (rh/fr)

Next Review Date: 3rd Qtr, 2027

BLUE CARE NETWORK BENEFIT COVERAGE
POLICY: NONTRADITIONAL BIOMARKERS IN RISK ASSESSMENT AND MANAGEMENT OF
CARDIOVASCULAR DISEASE

I. Coverage Determination:

Commercial HMO (includes Self-Funded groups unless otherwise specified)	Covered, policy criteria apply
BCNA (Medicare Advantage)	Refer to the Medicare information under the Government Regulations section of this policy.
BCN65 (Medicare Complementary)	Coinsurance covered if primary Medicare covers the service.

II. Administrative Guidelines:

- The member's contract must be active at the time the service is rendered.
- Coverage is based on each member's certificate and is not guaranteed. Please consult the individual member's certificate for details. Additional information regarding coverage or benefits may also be obtained through customer or provider inquiry services at BCN.
- The service must be authorized by the member's PCP except for Self-Referral Option (SRO) members seeking Tier 2 coverage.
- Services must be performed by a BCN-contracted provider, if available, except for Self-Referral Option (SRO) members seeking Tier 2 coverage.
- Payment is based on BCN payment rules, individual certificate and certificate riders.
- Appropriate copayments will apply. Refer to certificate and applicable riders for detailed information.
- CPT - HCPCS codes are used for descriptive purposes only and are not a guarantee of coverage.