
Medical Policy



Nonprofit corporations and independent licensees
of the Blue Cross and Blue Shield Association

Joint Medical Policies are a source for BCBSM and BCN medical policy information only. These documents are not to be used to determine benefits or reimbursement. When Centers for Medicare and Medicaid (CMS) coverage rules are not fully developed, this medical policy may be used by BCBSM or BCN Medicare Advantage plans 42 CFR § 422.101 (b)(6). Please refer to the appropriate certificate or contract for benefit information. This policy may be updated and is therefore subject to change.

***Current Policy Effective Date: 11/1/26**
(See policy history boxes for previous effective dates)

Title: Accelerated Breast Irradiation and Brachytherapy Boost after Breast-Conserving Surgery for EARLY-STAGE Breast Cancer

Description

Radiotherapy is the standard of care for patients with breast cancer undergoing breast-conserving surgery because it reduces recurrences and lengthens survival. A conventional radiotherapy regimen consists of approximately 25 treatments of 2 Gray (a measure of absorbed radiation dose) delivered over 5 to 6 weeks. Nonetheless, not all patients undergo radiotherapy following breast-conserving surgery; the duration and logistics of treatment may be barriers for some women. Accelerated radiotherapy approaches have been proposed to make the regimen less burdensome for patients with early-stage breast cancer at a low risk of recurrence. Accelerated (also called hypofractionated) whole-breast irradiation (AWBI) reduces the number of fractions and the duration of treatment to about 3 weeks. Accelerated partial-breast irradiation (APBI) targets a limited part of the breast in and close to the tumor cavity. By reducing the area irradiated, fewer treatments are needed, and the total treatment takes about 1 week.

Medical Policy Statement

Following breast-conserving surgery (BCS) for early-stage breast cancer and ductal carcinoma in situ (DCIS) accelerated (hypofractionated) whole breast irradiation (AWBI) with or without boost therapy, accelerated (hypofractionated) partial breast irradiation (APBI), and brachytherapy are established when criteria are met.

Intraoperative radiation therapy is experimental/investigational. There is insufficient evidence to determine that the technology results in an improvement in net health outcomes.

Noninvasive brachytherapy using Accuboot® is experimental/investigational. There is insufficient evidence to determine that the technology results in an improvement in net health outcomes.

Inclusionary and Exclusionary Guidelines

Inclusions:

Following breast-conserving surgery for early-stage breast cancer:

- Accelerated (hypofractionated) whole breast irradiation (AWBI) for stage 0 (DCIS), I, or II breast cancer. (See Table 1 below)
- AWBI (hypofractionated) followed by electrons or photons as a boost to the surgical bed for radiation treatment to the whole breast with or without treatment to the low axilla.
- AWBI (hypofractionated) followed by electrons or photons as a boost to the surgical bed in **ANY** of the following situations:
 - Regional lymph node radiation requiring a separate supraclavicular, axillary, and/or internal mammary node field,
 - Collagen vascular disease,
 - Breast augmentation,
 - Previous radiation to the breast or chest wall.
- Interstitial or balloon brachytherapy for patients undergoing initial treatment for stage I or II breast cancer when used as local boost irradiation in patients who are also treated with breast-conserving surgery and whole-breast external-beam radiotherapy.
- Accelerated partial breast irradiation (APBI)* for individuals with early-stage invasive breast cancer when **ALL** of the following criteria are met:
 - Age $\geq$ 40 years of age
 - Tumor less than or equal to 2 cm
 - ER-positive histology
 - Grade 1-2 disease
 - Negative surgical margins
 - Negative lymph nodes
 - No lymphovascular space invasion
 - No known germline pathogenic BRCA 1 or BRCA 2 variant
 -
- Accelerated partial breast irradiation (APBI)* for individuals with DCIS with **ALL** of the following criteria are met:
 - Low-to-intermediate grade
 - Age $\geq$ 40 years
 - Size $\leq$ 2 cm
 - Negative surgical margins
 - No known germline pathogenic BRCA 1 or BRCA 2 variant
 -

*APBI via 3-dimensional conformal radiation therapy (3-D CRT), Intensity modulated radiation therapy (IMRT) or multicatheter brachytherapy.

Table 1. Stages of Breast Cancer (American Cancer Society)

Stage	Category	Description
0	Tis, NO, MO	- This is ductal carcinoma in situ (DCIS), a pre-cancer of the breast. Many consider DCIS the earliest form of breast cancer. In DCIS, cancer cells are still within a duct and have not yet invaded deeper into the surrounding fatty breast tissue. - Paget disease of the nipple (without an underlying tumor mass) is also stage 0. - In all cases the cancer has not spread to lymph nodes or distant sites.
1A	T1, NO, MO	The tumor is 2 cm (about $\frac{3}{4}$ of an inch) or less across (T1) and has not spread to lymph nodes (NO) or distant sites (MO).
IIA	T0 or T1, N1 (but not N1mi), MO	- The tumor is 2 cm or less across (or is not found) (T1 or T0) and either: ✓ It has spread to 1 to 3 axillary lymph nodes, with the cancer in the lymph nodes larger than 2 mm across (N1a), or ✓ It has spread to 1 to 3 axillary lymph nodes and to internal mammary lymph nodes (found on sentinel lymph node biopsy) (N1c). - The cancer has not spread to distant sites (MO).
1B	T0 or T1, N1mi, MO	- The tumor is 2 cm or less across (or is not found) (T0 or T1) with micrometastases in 1 to 3 axillary lymph nodes (the cancer in the underarm lymph nodes is greater than 0.2 mm across and/or more than 200 cells but is not larger than 2 mm) (N1mi). - The cancer has not spread to distant sites (MO).
IIB	T2, N1, MO T3, NO, MO	- The tumor is larger than 2 cm but no larger than 5 cm; small groups of breast cancer cells—larger than 0.2 mm but not larger than 2 mm—are found in the lymph nodes, or - The tumor is larger than 2 cm but no larger than 5 cm; cancer has spread to 1 to 3 axillary lymph nodes near the breastbone, or - The tumor is larger than 5 cm but has not spread to the axillary lymph nodes.
IIIA	T0 to T2, N2, MO T3, N1 or N2, MO	- The tumor is not more than 5 cm across (or cannot be found). It has spread to 4 to 9 axillary lymph nodes, or it has enlarged the internal mammary lymph nodes. The cancer has not spread to distant sites - The tumor is larger than 5 cm across but does not grow into the chest wall or skin. It has spread to 1 to 9 axillary nodes, or to internal mammary nodes. The cancer has not spread to distant sites.
IIIB	T4, NO to N2, MO	- The tumor has grown into the chest wall or skin and one of the following applies: ✓ It has not spread to the lymph node ✓ It has spread to 1 to 3 axillary lymph nodes and/or tiny amounts of cancer are found in internal mammary lymph nodes on sentinel lymph node biopsy ✓ It has spread to 4 to 9 axillary lymph nodes, or it has enlarged the internal mammary lymph nodes. - The cancer has not spread to distant sites. - Inflammatory breast cancer is classified as T4d and is at least stage IIIB. If it has spread to many nearby lymph nodes it could be stage IIIC, and if it has spread to distant lymph nodes or organs it would be stage IV.
IIIC	Any T, N3, MO	- The tumor is any size (or cannot be found), and one of the following applies: ✓ Cancer has spread to 10 or more axillary lymph nodes. ✓ Cancer has spread to the lymph nodes under the collar bone. ✓ Cancer has spread to the lymph nodes above the collar bone. ✓ Cancer involves axillary lymph nodes and has enlarged the internal mammary lymph nodes.

		✓ Cancer has spread to 4 or more axillary lymph nodes, and tiny amounts of cancer are found in internal mammary lymph nodes on sentinel lymph node biopsy • The cancer has not spread to distant sites.
IV	Any T, any N, M1	• The cancer can be any size and may or may not have spread nearby lymph nodes. • It has spread to distant organs or to lymph nodes far from the breast. • The most common sites of spread are the bones, liver, brain, or lungs.

Exclusions:

- Accelerated whole breast irradiation for patients not meeting the above inclusions.
- Interstitial or balloon brachytherapy in all other situations not specified under the inclusions.
- Noninvasive brachytherapy using Accuboot® as part of any treatment for individuals who have undergone breast-conserving surgery.
- Local boost irradiation when combined with whole-breast radiotherapy but *without* surgical excision.
- Use of intraoperative radiation therapy.
- Use of electronic brachytherapy.

CPT/HCPCS Level II Codes (Note: The inclusion of a code in this list is not a guarantee of coverage. Please refer to the medical policy statement to determine the status of a given procedure)

Established codes:

19296	19297	19298	77261	77262	
77263	77280	77285	77290	77295	77316
77317	77318	77778			

Other codes (investigational, not medically necessary, etc.):

N/A	19294
-----	-------

Background

Radiotherapy is the standard care for patients with breast cancer undergoing breast-conserving surgery (BCS), as it reduces recurrences and lengthens survival. The conventional radiation therapy regimen consists of approximately 25 treatments of 2 Gray (Gy; a measure of absorbed radiation dose) delivered over 5 to 6 weeks. Nonetheless, not all patients undergo radiation therapy following BCS; the duration and logistics of treatment may be barriers for some women. Accelerated radiotherapy approaches have been proposed to make the regimen less burdensome for patients with early-stage breast cancer at low risk of recurrence:

- *Accelerated* (also called hypofractionated) whole-breast irradiation (AWBI) reduces the number of fractions and the duration of treatment to about 3 weeks.
- *Accelerated* partial-breast irradiation (APBI) irradiates a limited part of the breast in and close to the tumor cavity. By reducing the area irradiated, fewer treatments are needed, and the total treatment takes about 1 week. Options for the delivery of APBI include

brachytherapy (both interstitial brachytherapy and intracavitary brachytherapy), intraoperative radiation therapy (RT), or external beam radiation (conformal external beam radiation (EBR) or Intensity-modulated RT). Conformal external beam radiation is the most commonly used delivery system in the United States.

BREAST CANCER

Current estimates suggest that 321,910 new cases of breast cancer of any stage will occur in the United States in 2026. Based on adjusted data from 2019 to 2024, among women, the number of new cases is 132.5 per 100,000 women per year and the number of deaths 18.9 per 100,000 women per year.²

Breast Conservation Therapy

For patients diagnosed with stage I or II breast tumors, survival after breast-conservation therapy (BCT) is equivalent to survival after mastectomy. BCT is a multimodality treatment that consists of BCS to excise the tumor with adequate margins, followed by whole-breast external-beam radiotherapy (EBRT) administered as 5 daily fractions per week over 5 to 6 weeks. Local boost irradiation to the tumor bed often is added to whole-breast irradiation to provide a higher dose of radiation at the site where recurrence most frequently occurs. For some patients, BCT also includes axillary lymph node dissection, sentinel lymph node biopsy, or irradiation of the axilla. A number of randomized, controlled trials have demonstrated that the addition of radiotherapy after BCS reduces recurrences and mortality. In an expanded update of an individual-level meta-analysis, the Early Breast Cancer Trialists' Collaborative Group (EBCTCG) reported that radiotherapy halved the annual recurrence rate after 10 years for women with node-negative disease (n=7287), from 31.0% for those not receiving radiotherapy to 15.6% for those receiving it.³ It also reduced the 15-year risk of breast cancer death from 20.5% to 17.2% (p=0.005). For women with node-positive disease (n=1050), radiotherapy reduced the 1-year recurrence risk from 26.0% to 5.1%. Radiotherapy also reduced the 15-year risk of breast cancer death from 51.3% to 42.8% (p=0.01).

Consequently, radiotherapy is generally recommended following breast-conserving surgery. A potential exception is for older women at low risk of recurrence. For example, current National Comprehensive Cancer Network (NCCN) guidelines state that women ages 70 years or older may omit radiotherapy if they are hormone receptor-positive, *HER2*-negative, have T1 tumors, have clinically negative lymph nodes, and plan to take adjuvant endocrine therapy, or if they are 65 years or older with hormone receptor positive and *HER2*-negative tumors (≤ 3 cm) and have no lymph node metastases.⁴ However, the agreement is not universal.⁵

Controversy continues on the length of follow-up needed to determine whether APBI is equivalent to WBI.⁶ Because recurrences are relatively rare among low-risk early breast cancer patients, it may take considerable time for there to be enough recurrences to achieve sufficient power to compare rates across radiotherapy approaches. Additionally, radiation-induced adverse cardiovascular effects and radiation-induced non-breast cancers tend to occur 10 or more years after treatment.⁷⁻⁹ For accelerated whole-breast irradiation (AWBI), some 10-year data are available.

Currently, most patients diagnosed with stage I or II breast cancer are offered a choice between BCT and mastectomy, but BCT is selected less often than expected. Studies have shown that those living farthest from treatment facilities are least likely to select BCT instead of mastectomy and most likely to forgo radiotherapy after breast-conserving surgery,^{10,11,12} and

have recommended the use of multidisciplinary management strategies to eliminate known disparities in rural, minority, and uninsured populations.¹³

Approaches to Radiotherapy Following Breast-Conservation Treatment

The goals of cancer radiotherapy are to deliver a high dose of homogeneous radiation (i.e., all parts of the tumor cavity receive close to the targeted dose) to the tumor or tumor bed. Areas adjacent to the tumor may be given a lower dose of radiation (e.g., with WBI) to treat any unobserved cancerous lesions. Radiation outside the treatment area should be minimal or nonexistent. The goal is to target the tumor or adjacent areas at risk of harboring unseen cancer with an optimum dose while avoiding healthy tissues.

Table 2 outlines the major types of radiotherapy used after BCS. They differ by technique, instrumentation, dose delivery, and possible outcomes.

Table 2. Major Types of Radiotherapy Following Breast-Conserving Surgery^a

Radiation Type	Accelerated?	Whole or Partial Breast	EBRT or Brachytherapy	Treatment Duration
Conventional WBI	No	Whole	EBRT	5-6 wk
Accelerated WBI	Yes	Whole	EBRT	3 wk
Interstitial APBI ^b	Yes	Partial	Brachytherapy	1 wk
Balloon APBI ^c	Yes	Partial	Brachytherapy	1 wk
EBRT APBI ^d	Yes	Partial	EBRT	1 wk
Intraoperative APBI ^e	Yes	Partial	Not applicable	1 d

APBI: accelerated partial-breast irradiation; EBRT: external-beam radiotherapy; RCT: randomized controlled trial; WBI: whole-breast irradiation.

^a Noninvasive breast brachytherapy using AccuBoost has been described by the manufacturer as capable of delivering APBI, but no studies for this indication were found.

^b Interstitial brachytherapy entails placement of multiple hollow needles and catheters to guide placement of the radioactive material by a remote after-loading device. It is more difficult to perform than other types of brachytherapy and has a steep learning curve.

^c Balloon brachytherapy (e.g., MammoSite) entails inserting a balloon into the tumor bed, inflating the balloon, confirming its position radiographically, and then using a remote after-loader to irradiate the targeted area. Some brachytherapy systems combine aspects of interstitial and balloon brachytherapy.

^d External-beam APBI is delivered in the same way as conventional or accelerated whole-breast radiotherapy but to a smaller area. All three external-beam regimens can use 3-dimensional conformal radiotherapy or intensity-modulated radiotherapy.

^e Intraoperative APBI is performed during breast-conserving surgery with a single dose of radiation delivered to the exposed tumor bed.

Summary of Evidence

Accelerated Whole-Breast Irradiation

For individuals who have node-negative, early-stage breast cancer with clear surgical margins who receive accelerated whole-breast irradiation (AWBI) after breast-conserving surgery (BCS), the evidence includes randomized controlled trials (RCTs) and systematic reviews. Relevant outcomes are overall survival, disease-specific survival, change in disease status, and treatment-related morbidity. Two randomized noninferiority trials both reported 10-year follow-up data on local recurrence. Both trials found that local recurrence rates with AWBI were no worse than conventional whole-breast irradiation (WBI), when applying a noninferiority margin of 5%. Conclusions apply to patients meeting eligibility criteria of the RCTs trials, including having early-stage invasive breast cancer, clear surgical margins, and negative lymph nodes. In addition, consistent with national guidelines, these conclusions apply to

tumors more than 5 cm in diameter and women at least 50 years old. There is sufficient evidence to determine that the technology results in an improvement in net health outcomes.

Accelerated Partial-Breast Irradiation

For individuals who have early-stage breast cancer who receive interstitial brachytherapy, the evidence includes 1 completed RCT. Relevant outcomes are overall survival, disease-specific survival, change in disease status, and treatment-related morbidity. The RCT reported 5-year follow-up data and found that interstitial brachytherapy was noninferior to WBI for rates of local breast cancer recurrence, when applying a noninferiority margin of 3%. Ten-year follow-up data published in 2023 reported a nonsignificant 1.93% difference in the cumulative incidence of ipsilateral tumor recurrence. There is sufficient evidence to determine that the technology results in an improvement in net health outcomes.

For individuals who have early-stage breast cancer who receive intraoperative brachytherapy, the evidence includes systematic reviews and RCTs, and a large registry study. Relevant outcomes are overall survival, disease-specific survival, change in disease status, and treatment-related morbidity. Several RCTs have been published, but they have not demonstrated that outcomes after intraoperative brachytherapy are noninferior to WBI. Results of 2 RCTs (TARGIT-A, ELIOT) comparing intraoperative brachytherapy to WBI found higher rates of local recurrence with intraoperative brachytherapy than with WBI. There is insufficient evidence to determine that the technology results in an improvement in net health outcomes.

For individuals who have early-stage breast cancer who receive external-beam accelerated partial breast irradiation (APBI), the evidence includes a systematic review and RCTs. Relevant outcomes are OS, disease-specific survival, change in disease status, and treatment-related morbidity. A comparative-effectiveness review from the Agency for Healthcare Research and Quality reported with high-strength evidence that partial-breast irradiation (PBI) offers ipsilateral recurrence, overall survival, and cancer-free survival rates equivalent to whole-breast irradiation (WBI) through 10 years of follow-up, and its pre-specified subgroup analysis revealed no outcome differences between APBI and conventional PBI. Across 6 randomized trials with 5 to 20-year follow-up, cumulative ipsilateral breast-tumor recurrence with APBI ranged 2.7% to 9.6% versus 2.8% to 7.9% with WBI; hazard ratios for recurrence or disease-free survival met pre-specified non-inferiority margins in most studies, although the largest trial was found non-equivalent despite an absolute 10-year difference of <1 %. Overall survival and distant disease-free survival were not significantly different in observations from 5 to 20 years. Several studies have reported that APBI produces higher rates of acute skin reactions, adverse cosmesis, and fibrosis/telangiectasia; however, serious grade ≥3 toxicities remained uncommon and were comparable between APBI and WBI. There is sufficient evidence to determine that the technology results in an improvement in net health outcomes.

Local Boost Brachytherapy

For individuals who have early-stage breast cancer who receive local boost brachytherapy with WBI, the evidence includes nonrandomized studies and a systematic review. Relevant outcomes are overall survival, disease-specific survival, change in disease status, and treatment-related morbidity. Nonrandomized comparative studies have shown similar outcomes with brachytherapy local boost and with external-beam radiotherapy local boost. There is sufficient evidence to determine that the technology results in an improvement in net health outcomes.

Noninvasive Breast Brachytherapy

For individuals who have early-stage breast cancer who receive noninvasive breast brachytherapy, the evidence includes 1 retrospective comparative study. Relevant outcomes are overall survival, disease-specific survival, change in disease status, and treatment-related morbidity. The retrospective study was a matched comparison of noninvasive breast brachytherapy or electron-beam radiotherapy to provide boost radiation to the tumor bed. The study was subject to selection bias, relatively short follow-up, and use of a retrospective design. There is insufficient evidence to determine that the technology results in an improvement in net health outcomes.

Rationale

Evidence reviews assess clinical evidence to determine whether the use of a technology improves the net health outcome. Broadly defined, health outcomes are length of life, quality of life, and ability to function including benefits and harms. Every clinical condition has specific outcomes that are important to individuals and to managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms. The following is a summary of the key literature to date.

ACCELERATED WHOLE-BREAST IRRADIATION

Clinical Context and Therapy Purpose

The purpose of accelerated whole-breast irradiation (AWBI) after BCS in individuals who have node negative, early-stage breast cancer with clear surgical margins is to provide a treatment option that is an alternative to or an improvement on existing therapies.

Systematic Reviews

Systematic reviews of RCTs have compared AWBI (also referred to as accelerated whole-breast radiotherapy) with conventional 5-week WBI. A systematic review and meta-analysis by Valle et al (2017) included 13 trials (N=8189 patients) published prior to October 2014 that compared AWBI with standard fractionation.¹⁶ No differences were observed in local recurrence (7 trials; relative risk [RR], 0.97; 95% confidence interval [CI], 0.78 to 1.19), locoregional failure (8 trials; RR, 0.86; 95% CI, 0.63 to 1.16), or survival (4 trials; RR, 1.00; 95% CI, 0.85 to 1.17). There was less acute toxicity with AWBI (5 trials; RR, 0.36; 95% CI, 0.21 to 0.62), and no difference in late cosmesis (RR, 0.95; 95% CI, 0.81 to 1.12). The largest trials included in the meta-analysis were the Standardization of Breast Radiotherapy (START) A, START B, and NCIC (detailed below).¹⁷⁻¹⁹

Randomized Controlled Trials

Two of the RCTs included in the systematic review were noninferiority trials that directly compared a 5- week with a 3-week regimen (Bentzen et al, 2008; Whelan et al, 2002; Whelan et al (2010)).¹⁷⁻¹⁹ Both trials used noninferiority margins of 5 percentage points for local or locoregional recurrence in the accelerated group at 5 (1-sided $\alpha=0.025$ or 0.05)^{17,18} or 10 years (1-sided $\alpha=0.025$).¹⁹ Although the trials differed in specific fractionation schedules and

patient characteristics, they reported similar ipsilateral local recurrence rates (i.e., cancer recurrence in the same breast) across treatment arms.

The first RCT evaluating an accelerated whole-breast radiotherapy regimen (Standardizations of Breast Radiotherapy [START] B; 2008), from the U.K., included women with stage I, II, or III tumors (N=2215) who had clear tumor margins (≥ 1 mm).¹⁷ Approximately 75% of the women have negative lymph nodes, and approximately 42% had a radiation boost to the tumor bed. Randomization was stratified for hospital, type of surgery (8% underwent mastectomy), and plans for tumor bed boost. Systemic therapy, primarily tamoxifen, was used by some patients and appears to be fairly evenly distributed across treatment groups. Treatment arms compared a total dose of 40 Gy in 15 fractions over 3 weeks to 50 Gy in 25 fractions over 5 weeks. The primary efficacy outcome was locoregional relapse (relapse in ipsilateral breast or chest wall and or in ipsilateral axilla or supraclavicular fossa if previously irradiated) at 5 years. At median follow-up of 6.0 years (interquartile range [IQR], 5.0-6.2), estimated 5-year locoregional tumor relapse rate was 2.2% (95% confidence interval [CI], 1.3% to 3.1%) in the 40 Gy group and 3.3% (95% CI, 2.2% to 4.5%) in the 50 Gy group, for an absolute difference of -0.7% (95% CI, -1.7% to 0.9%). Hazard ratios (HRs) for 40 Gy accelerated whole-breast radiotherapy versus conventional whole-breast radiotherapy were not statistically significant (using the log-rank test) for local or local-regional relapses. There were statistically significant differences between the 2 treatment regimens for distant relapses and overall survival (OS), with relapses less frequent and survival longer for the 40 Gy accelerated whole-breast irradiation (AWBI) group. This unexpected difference between treatment arms began to appear at about 1 year; trial authors speculated that the difference may have been due to chance and may have changed with longer follow-up.

Subsequent publications provided additional results for both START trials (i.e., START A, which compared two 5-week whole breast radiotherapy regimens, and START B). Hopwood et al (2010) examined patient-reported breast, arm, and shoulder symptoms, as well as body image, over 5 years of follow-up.²⁰ There was no evidence that providing radiotherapy in fewer, larger fractions increased the incidence of these adverse events or adversely affected body image. Haviland et al (2013) reported 10-year relapse, survival, and adverse event outcomes (median follow-up, 9.9 years [IQR, 7.5-10.1 years]).²¹ Locoregional relapse did not differ significantly between the 2 treatment groups: 4.3% (95% CI, 3.2 to 5.9) for the AWBI group and 5.5% (95% CI, 4.2 to 7.2) for the standard whole-breast radiotherapy group (HR=0.77 [95% CI, 0.51 to 1.16]; p=0.21). However, breast shrinkage, telangiectasia, and breast edema were significantly less common in the AWBI group. These effects were assessed by physicians, photographic comparison with baseline, and patient report. Distant relapse (p=0.014), any breast-cancer-related event (local, regional, or distant relapse, breast cancer death, contralateral breast cancer; p=0.022), and all-cause mortality (p=0.042) were significantly less common in the AWBI group. These effects were assessed by a physician, photographic comparison with baseline, and patient report.

The second RCT assessing a 5- and 3-week radiotherapy regimen, from Canada, compared AWBI versus whole-breast irradiation (WBI) in women with lymph node-negative stage I, II, or III tumors.^{18,19} Treatment arms included a hypofractionated radiation group (n=622), who were treated with a total dose of 42.5 Gy in 16 fractions over 3 weeks, and a standard irradiation group (n=612), who were treated with 50 Gy in 25 fractions over 5 weeks without boost radiation. Five-year local recurrence-free survival was 97.2% in the accelerated arm and 96.8% in the conventional arm (absolute difference: 0.4% [95% CI, -1.5% to 2.4%]). Ten-year

local recurrence was 6.7% for the conventional arm and 6.7% for the conventional arm (absolute difference: -0.5% [95% CI, -2.5% to 3.5%]). At 5 or 10 years, local recurrence rates with AWBI were no worse than conventional WBI, when applying a noninferiority margin of 5%. In prespecified subgroup analyses, treatment effects were similar by age, tumor size, estrogen-receptor status, and chemotherapy use (48% had no systemic therapy).

An RCT by Shaitelman et al (2015), not included in the Valle et al (2017) systematic review, focused on acute and short-term toxicity for conventional versus accelerated whole-breast radiotherapy.²² This unblinded trial included 287 patients with stage 0 to III breast cancer treated with breast-conserving therapy who had negative tumor margins. Patients were randomized to conventional radiotherapy at 50 Gy in 25 fractions (n=149) or accelerated radiotherapy at 42 Gy in 16 fractions (n=138). The rate of grade 2 or higher acute toxic events was 47% in the accelerated radiotherapy group and 78% in the conventional radiotherapy group (p<0.001). A total of 271 (94%) of 287 patients were available for an assessment of 6-month toxic effects. There were no significant between-group differences in toxic effects at 6 months except that the rate fatigue (grade ≥2) was significantly lower in the accelerated radiotherapy group (0%) than in the conventional radiotherapy group (6%; p=0.01).

In 2020, Brunt et al published 10-year results of the FASTer radiotherapy for breast radiotherapy (FAST) trial.²³ This multicenter, phase III, RCT enrolled 915 women ≥50 years of age with low-risk invasive breast carcinoma who had undergone BCS with complete microscopic resection and randomly assigned them to 50 Gy in 25 fractions of 2 Gy, 30 Gy in 5 once weekly fractions of 6 Gy, or 28.5 Gy in 5 once weekly fractions of 5.7 Gy. At the time of this analysis, median follow-up was 9.9 years (interquartile range: 8.3 to 10.1 years). Results revealed that the odds ratios for any moderate/marked physician-assessed breast normal tissue effects (i.e., shrinkage, induration, telangiectasia, edema) were significantly higher for the 30 Gy versus 50 Gy group (2.12; 95% CI, 1.55 to 2.89; p<0.001), but not significantly different for the 28.5 Gy versus 50 Gy group (1.22; 95% CI, 0.87 to 1.72; p=0.248). Additionally, 11 ipsilateral breast cancer events (50 Gy: 3; 30 Gy: 4; 28.5 Gy: 4) and 96 deaths (50 Gy: 30; 30 Gy: 33; 28.5 Gy: 33) were reported at 10 years of follow-up. These results confirm that a 5-fraction schedule (28.5 Gy in 5 once weekly fractions) is radiobiologically equivalent to the standard 25-fraction schedule with regard to late normal tissue effects.

Brunt et al (2020) also published results from the multicenter, non-inferiority, randomized, FAST-Forward trial.²⁴ This study enrolled 4096 adults with invasive breast carcinoma following complete microscopic excision of the primary tumor by BCS or mastectomy who were randomly assigned to 3 groups of hypofractionated radiotherapy: 40 Gy in 15 fractions over 3 weeks, 27 Gy in 5 fractions over 1 week, or 26 Gy in 5 fractions over 1 week. At a median follow-up of 71.5 months (interquartile range: 71.3 to 71.7 months), ipsilateral breast tumor relapse occurred in a total of 79 patients (40 Gy: 31; 27 Gy: 27; 26 Gy: 21); the hazard ratio for 27 Gy versus 40 Gy was 0.86 (95% CI, 0.51 to 1.44) and for 26 Gy versus 40 Gy was 0.67 (95% CI, 0.38 to 1.16). The estimated cumulative incidence of ipsilateral breast tumor relapse up to 5 years was 2.1% (95% CI, 1.4 to 3.1) for 40 Gy; 1.7% (95% CI, 1.2 to 2.6) for 27 Gy; and 1.4% (95% CI, 0.9 to 2.2) for 26 Gy. Estimated absolute differences in this outcome were -0.3% (95% CI, -1.0 to 0.9) for 27 Gy versus 40 Gy and -0.7% (95% CI, -1.3 to 0.3) for 26 Gy versus 40 Gy. Moderate or marked physician-assessed normal tissue effects in the breast or chest wall were seen in 9.9% of 40 Gy patients, 15.4% of 27 Gy patients, and 11.9% of 26 Gy patients at 5 years; a significant difference between 40 and 27 Gy (p=0.0003) but not between 40 and 26 Gy (p=0.17) was observed. These results show that a 1-week

course of adjuvant breast radiotherapy delivered in 5 fractions is non-inferior to the standard 3-week schedule, with the 26 Gy dose level being similar to 40 Gy in terms of local tumor control and normal tissue effects for up to 5 years.

Observational Studies

Toxicity rates were also evaluated in a large 2014 retrospective study of patients with left-sided early-stage breast cancer published by Chan et al (2014, 2015).^{25,26} The study included 2706 patients who received conventional WBI (n=2221) or (n=485) AWBI. Cardiotoxic chemotherapy regimens were similar between groups. At a median follow-up of 14.2 years, there was no statistical difference in cardiac hospitalization or cardiac mortality, breast cancer mortality, or overall mortality. Results were similar for 2628 patients with right-sided tumors. This study was not designed to capture outcomes of moderate or mild cardiac toxicity.

Section Summary: Accelerated Whole-Breast Irradiation

The overall body of evidence on AWBI compared to conventional whole-breast irradiation has indicated that local recurrence rates with accelerated whole-breast radiotherapy are no worse than conventional WBI, when applying a noninferiority margin of 5%. Canadian and U.K. noninferiority trials have reported 10-year follow-up data. Thus, conclusions apply to patients meeting eligibility criteria of these trials, including having early-stage invasive breast cancer, clear surgical margins, and negative lymph nodes. In addition, consistent with national guidelines, these conclusions apply to tumors greater than 5 cm in diameter and women at least 50 years old. Based on 14-year retrospective data, severe cardiac toxicity with AWBI for left-sided breast cancers may not be increased compared with conventional WBI. Additionally, recent data implies that even more accelerated WBI scheduling may be non-inferior to standard 3- or 5-week schedules.

ACCELERATED PARTIAL-BREAST IRRADIATION

Clinical Context and Therapy Purpose

The purpose of APBI in individuals who have early-stage breast cancer is to provide a treatment option that is an alternative to or an improvement on existing therapies.

Systematic Reviews

A number of RCTs and nonrandomized comparative studies have evaluated interstitial, external-beam, or intraoperative APBI compared with conventional WBI. Several meta-analyses have evaluated evidence on APBI, with various methods grouped in the same review.²⁷⁻³⁴ Conclusions cannot be drawn from these meta-analyses because methods varied and were not evaluated individually. The review authors were consistent in concluding that additional data from RCTs are needed. In 2020, Viani et al published a systematic review and update meta-analysis of partial- versus whole-breast radiotherapy for early breast cancer that included a subgroup analysis assessing the potential effectiveness of APBI technique - intraoperative radiotherapy (IORT), brachytherapy, or EBRT.³⁵ Results revealed no significant difference in local recurrence with APBI and WBI when using brachytherapy (p=0.051), EBRT (p=0.25), or mixed techniques (p=0.89) at 5 years; however, a significant increase in local recurrence was noted with IORT use (p=0.014). At 7- and 10-years follow-up, the difference in local recurrence within the IORT subgroup disappeared. Additionally, an analysis of overall mortality revealed no difference between 5, 7, and 10 years of follow-up for any subgroup. Korzets et al (2019) revealed similar results from a subgroup analysis of APBI modality within a systematic review and meta-analysis that evaluated toxicity and clinical outcomes of partial-

versus WBI for early-stage breast cancer.³⁶ These authors concluded that the highest risk of local recurrence was seen with IORT, whereas when EBRT was used the odds for local recurrence were equivalent to WBI. The IORT studies included a larger number of patients with high-grade disease and nodal involvement, which may partially explain the increased local recurrence rate with this modality.

Interstitial Brachytherapy

Randomized Controlled Trials

GEC-ESTRO was a European multicenter noninferiority RCT with 5-year results (Table 3). Primary results were published in 2016, late-side effects in 2017, and quality of life in 2018.³⁷⁻³⁹ The primary study end point was the first incidence of local ipsilateral breast cancer recurrence within the 5-year observation period and the noninferiority margin was 3% difference. At 5 years, the associated cumulative incidence of local recurrence was 0.92% (95% CI, 0.12% to 1.73%) in the conventional WBI group and 1.44% (95% CI, 0.51% to 2.38%) in the APBI group (Table 3). The difference between groups was within the non-inferiority margin. There was no grade 4 skin toxicity. Grade 2 and 3 skin toxicity was 10.7% with WBI and 6.9% with APBI (p=0.02).

Ten-year outcomes of the GEC-ESTRO trial were published by Strnad et al in 2023.⁴⁰ At a median follow-up of 10.36 years, rates of local recurrence were 1.58% (95% CI, 1.99 to 5.03) in the WBI group and 3.51% (95% CI, 1.99 to 5.03) in the APBI group (Table 3). Significantly fewer treatment-related grade 3 late adverse effects were observed in the APBI group (1%) compared to the WBI group (4%). No grade 4 adverse events or treatment-related deaths were reported. Additional publications with detailed analyses of late adverse effects (e.g., cosmesis) and quality of life outcomes are planned by the investigators.

Table 3. Summary of Key RCT Characteristics

Study	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
GEC-ESTRO	EU	16	2004-2009	1328 patients ≥40 y with BCS for stage 0-IIa breast cancer, lesions ≤3 cm in diameter, clear margins ≥2 mm in any direction, and no lymph or blood vessel invasion	655 patients given APBI using interstitial brachytherapy	673 patients given WBI at 50 Gy in daily fractions of 1.8-2.0 Gy over 5 wk

APBI: accelerated partial breast irradiation; BCS: breast-conserving therapy; Gy: gray; RCT: randomized controlled trial; WBI: whole-breast irradiation.

Table 4. Summary of Key RCT Results

Study	Local Recurrence, n (%)	Overall Survival	Grade 2 to 3 Late Skin Toxicity	Excellent-to-Good Cosmetic Results, n (%)	Global Health Status (SD)
GEC-ESTRO					
5-year Outcomes					
n	1184	1184	1184	1007	537

WBI	5 (0.92)	95.5%	5.7%	408 (90)	66.0 (21.8)
APBI	9 (1.44)	97.27%	3.2%	503 (93)	66.2 (22.2)
Diff (95% CI)	0.52% (-0.72% to 1.75%)	1.72% (-0.44% to 3.88%)			-0.2 (-4.0 to 3.6)
p	NS	0.11	0.080	0.12	0.94
10-year Outcomes					
n	772	772	688	688	NR
WBI	9 (1.58)	89.5%	3%	121/ 313 (34)	NR
APBI	21 (3.51)	90.5%	1%	188/446 (45)	NR
Difference (95% CI)	1.93%		Difference (95% CI)	1.93%	
p	.074	.50	.70	NR	NR

APBI: accelerated partial breast irradiation; CI: confidence interval; Diff: difference; RCT: randomized controlled trial; RRWBI: whole-breast irradiation; SD: standard deviation.

Major limitations in relevance and design and conduct are shown in Tables 5 and 6, respectively. This information is synthesized as a summary of the body of evidence following each table and provides conclusions on the sufficiency of the evidence supporting the position statement.

Table 5. Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Follow-Up ^e
GEC-ESTRO				1. Overall survival was not a primary outcome	

The study limitations stated in this table are those notable in the current review; this is not a comprehensive limitations assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

Table 6. Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
GEC-ESTRO		1-3. Not blinded				1. No prespecified noninferiority analysis on survival outcomes

The study limitations stated in this table are those notable in the current review; this is not a comprehensive limitations assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

^b Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

Nonrandomized Studies

Kim et al (2023) reported 5-year survival outcomes in 223 patients with early-stage breast cancer prospectively enrolled in a single-center registry.⁴¹ Patients were treated with breast-conserving surgery and individualized applicator-based brachytherapy administered via a Strut Adjusted Volume Implant (SAVI) device. After a median follow-up of 63 months, recurrence was reported in 19 (8.5%) patients with ipsilateral breast tumor recurrence occurring in 17 (7.6%). Rates of 5-year ipsilateral breast tumor-free survival, disease-free survival, and OS were 92.2%, 91.1%, and 97.8%, respectively. Significantly higher rates of 5-year ipsilateral breast tumor-free survival were noted in postmenopausal women (93.6% vs. 66.4%; $p=.04$), patients with body mass index $<30 \text{ kg/m}^2$ (97.5% vs. 88.1%; $p=.02$), and endocrine therapy adherence in postmenopausal women (97.5% vs. 88.6%; $p=.02$). Eight (3.6%) developed a grade 2 or higher treatment-related complication within 9 months of brachytherapy completion.

Ajkay et al (2015) reported retrospectively on 5-year adverse events in patients with early-stage breast cancer treated at a single center.⁴² Of 417 patients who received BCS and radiotherapy, 271 received brachytherapy (34 Gy in 10 fractions; 90% MammoSite, 9% Contura, 1% strut-adjusted volume implant) and 146 received WBI using 3-dimensional conformal radiotherapy (45-50.4 Gy in 25-28 fractions with 10-16 Gy boost). Median follow-up was 4.8 years in the brachytherapy group and 4.1 years in the WBI group. The estimated 5-year overall incidence of any adverse event was greater in the brachytherapy group (72%) than in the WBI group (52%; $p<0.001$). For prespecified adverse events of interest, estimated 5-year incidences of infectious skin complications, abscess, telangiectasia, and breast pain were similar between groups. Estimated 5-year incidences of seroma (47% vs. 19%, $p<0.001$) and fat necrosis (40% vs. 24%, $p<0.001$) were greater in the brachytherapy group, respectively.

Section Summary: Interstitial Brachytherapy

The 2015 GEC-ESTRO RCT reported 5-year follow-up data and found that interstitial brachytherapy was noninferior to WBI on rates of local breast cancer recurrence, when applying a noninferiority margin of 3%. Ten-year follow-up data published in 2023 reported a nonsignificant 1.93% difference in the cumulative incidence of local recurrence between groups.

Intraoperative Radiotherapy

Systematic Reviews

Ravani et al (2024) conducted a meta-analysis of patient-level data from RCTs that compared partial breast irradiation, IORT, and WHI in patients with early-stage breast cancer who underwent breast-conserving surgery.⁴³ A total of 11 trials was included ($N=15,460$), and the trials had a median follow-up of 9 years. About half of the included patients (49.6%) received WBI, while 35% received partial breast irradiation, and 15.3% received IORT. Patients who received IORT had a higher risk of recurrence (HR, 1.46; 95% CI, 1.23 to 1.72; $p<.01$) compared to WBI with similar risk of OS ($p=.81$) and disease-free survival ($p=.31$). The authors concluded that IORT was inferior to WBI.

Wang et al (2021) conducted a systematic review and meta-analysis to compare the efficacy and safety of IORT to WBI in patients with early-stage breast cancer receiving breast-conserving surgery.⁴⁴ A total of 10 RCTs representing 5,698 patients were included for

analysis. The IORT group was associated with a significantly higher risk of local recurrence (relative risk [RR], 2.11; 95% CI, 1.13 to 3.94; $p=.019$). Pooled analyses did not find any statistically significant differences in overall survival, recurrence-free survival, distant metastasis-free survival, and cancer-specific survival between groups. While the risk of skin toxicity was significantly lower in the IORT group compared to the WBI group (RR, 0.28; 95% CI, 0.16 to 0.49; $p<.0001$), the incidence of fat toxicity, scar calcification, and edema were 3.1, 1.3, and 2.4 times higher than that in the WBI group, respectively. The authors concluded that IORT is not a better alternative to WBI.

Randomized Controlled Trials

One RCT, reported by Vaidya et al (2010, 2014) compared intraoperative radiotherapy (IORT) with WBI in 2232 women.^{45,46} Radiotherapy was delivered to the tumor bed using the Intrabeam® device, which provides a point source of 50 kV energy x-rays at the center of a spherical applicator, for 20 to 45 minutes. It was specifically developed for IORT. The TARGIT-A (Risk-adapted Targeted Intraoperative Radiotherapy) trial was a noninferiority trial with 28 centers in 9 countries and a sample size of 3451. (In 2010, the trial was extended for 2 more years to allow accrual in sub-protocols.) An ITT approach was used. Patients were not blinded to treatment choice. As anticipated in advance, 14% of those in the IORT arm received EBRT as well, because of unfavorable pathologic features determined after surgery (e.g., lobular carcinoma). The predefined noninferiority margin was an absolute difference of 2.5% between groups for pathologically confirmed, ipsilateral local recurrence. The most recent article reported 5-year results, defined as results for patients with 5-years of follow-up or "if they were seen the year before database lock."⁴⁶ Median follow-up for all patients was 2 years and 5 months (IQR, 12-52 months), and 1222 patients (35%) had a median follow-up of 5 years. Estimated 5-year risks for ipsilateral local recurrence were 3.3% (95% CI, 2.1 to 5.1) in the TARGIT group and 1.3% (95% CI, 0.7 to 2.5; $p=0.042$) in the whole-breast radiotherapy group. Mortality was similar between the 2 groups (2.6% TARGIT vs. 1.9%; $p=0.56$). However, there were significantly fewer non-breast cancer deaths in the TARGIT group than in the whole-breast radiotherapy group (1.4% [95% CI, 0.8 to 2.5] vs. 3.5% [95% CI, 2.3 to 5.2]; $p<0.001$), with fewer deaths from cardiovascular causes and other cancers in the TARGIT group. In the group that received IORT plus whole-breast radiotherapy, the mortality rate was higher at 8% (95% CI, 3.7 to 17.5), but the percentage of women with local recurrences (0.9%; 95% CI, 0.1 to 6.1) was similar to those who received only IORT. Non-inferiority was established for the whole intraoperative cohort and for those who received IORT alone, but not for those patients who underwent both types of radiotherapy. There was no significant difference between the IORT and WBI groups in predefined 6-month wound-related complications. However, grade 3 or 4 radiotherapy-related skin complications were more common in the WBI group (13/1730 vs. 4/1731; $p=0.029$). In 2016, the full final report of the TARGIT-A trial was published concluding that "for patients with breast cancer (women who are aged ≥ 45 years with hormone-sensitive invasive ductal carcinoma that is up to 3.5 cm in size), targeted IORT concurrent with lumpectomy within a risk-adapted approach is as effective as, safer than, and less expensive than postoperative EBRT."⁴⁷

In a parallel study to TARGIT-A, Vaidya and colleagues (2020) randomly assigned 1153 patients who had undergone breast cancer excision to either conventional fractionated whole breast EBRT over 3 to 6 weeks or undergo a further operation to deliver delayed radiotherapy (as a single dose via Intrabeam) to the wound by reopening the original incision.⁴⁸ Results at 5 years revealed local recurrence rates of 3.96% for delayed IORT versus 1.05% for EBRT; a difference of 2.9% with an upper 90% CI of 4.4, which crossed the noninferiority

margin of 2.5%. Of note, at a median follow-up of 9 years, there are no significant differences between the 2 treatment approaches with regard to local recurrence-free survival, invasive local recurrence-free survival, mastectomy-free survival, distant disease-free survival, breast cancer mortality, and overall survival. The authors concluded that the results of this trial clearly show that the preferred timing of IORT use is during the initial surgical excision of breast cancer setting, not in the delayed setting; however, if immediate IORT is not possible the data from this trial may assist clinicians and patients who want to avoid a prolonged postoperative EBRT course.

Another form of IORT, called electron intraoperative radiotherapy (ELIOT), uses electrons (Veronesi et al, 2010, 2013).⁴⁹ The 2013 ELIOT trial compared IORT plus ELIOT to WBI.⁵⁰ With a sample size of 1305 patients and median follow-up of 5.8 years (IQR, 4.1-7.7 years), 35 (4.4%) patients in the intraoperative group and 4 (0.4%) patients in the WBI group developed ipsilateral breast tumor recurrences (HR=9.3; 95% CI, 3.3 to 26.3; $p<0.001$). There was no statistically significant difference in 5-year OS. For women with data on adverse skin events (IORT=464, WBI=412), there were significantly fewer events among women who received IORT ($p<0.001$). This was an equivalence trial with a prespecified limit of 7.5% for local recurrence in the IORT group only. Therefore, although the criterion for equivalence was satisfied, ipsilateral breast recurrence rate was significantly higher in the IORT group. A subsequent review of the ELIOT trial noted that, of 69 women who had 4 or more positive lymph nodes, those randomized to WBI ($n=38$) received concurrent axillary radiation; for those randomized to ELIOT ($n=31$), axillary irradiation was delayed 6 to 12 weeks.⁹ These reviewers also characterized ELIOT data as “still early” and noted that long-term results are needed to assess net health benefit. Orrechia et al (2021) reported 15-year results of the ELIOT trial that confirmed the 5-year findings.⁵¹ After a median follow-up of 12.4 years (interquartile range, 9.7 to 14.7 years), ipsilateral breast tumor recurrence had occurred in 70 patients (11%) in the IORT group 16 patients (2%) in the WBI group (hazard ratio, 4.62; 95% CI, 2.68 to 7.95; $p<.0001$). Fifteen-year OS was 83.4% in the IORT group and 82.4% in the WBI group. The authors concluded that low risk patients may be appropriate for IORT since the higher rate of ipsilateral breast cancer recurrence in the ELIOT trial did not lead to an increase in OS.

Registry Trials

Ding et al (2025) conducted a retrospective propensity score-matched cohort study using the Surveillance, Epidemiology, and End Results (SEER) database to compare survival outcomes associated with IORT versus WBI in patients with early-stage breast cancer.⁶⁹ The study included adult women diagnosed between 2010 and 2019 with unilateral invasive ductal carcinoma staged T1–2N0M0, estrogen receptor-positive disease, and treatment with breast-conserving surgery followed by either IORT or WBI. After exclusions, 83,474 patients were included, of whom 1,760 received IORT and 81,714 received WBI. Propensity score matching was performed in a 1:4 ratio using covariates including age, race, marital status, tumor grade, T stage, progesterone receptor, HER2 status, and chemotherapy exposure, resulting in a matched cohort of 8,800 patients. The primary endpoints were cancer-specific survival and OS. Median follow-up was 57 months (interquartile range, 12 to 119 months). Kaplan-Meier analyses demonstrated no significant differences in cancer-specific survival or OS between IORT and WBI before or after matching. In multivariable Cox regression analysis before matching, WBI was not associated with significantly different cancer-specific survival (HR, 1.23; 95% CI, 0.74 to 2.05; $p=.43$) or OS (HR, 1.30; 95% CI, 0.96 to 1.75; $p=.09$) compared with IORT. Positive estrogen receptor status (HR, 0.73; 95% CI, 0.57 to 0.92), positive progesterone receptor status (HR, 0.60; 95% CI, 0.52 to 0.69), and HER2-positive status (HR,

0.61; 95% CI, 0.51 to 0.74) were associated with improved cancer-specific survival, whereas age >65 years (HR, 1.82; 95% CI, 1.53 to 2.17) and T2 disease (HR, 3.72; 95% CI, 1.91 to 7.23) were associated with worse cancer-specific survival.

Cohort Studies

Silverstein et al (2025) reported on the results of their 15-year IORT registry trial and the rationale for stopping the trial.⁶⁸ Intraoperative radiation therapy (IORT) delivers the full course of radiation therapy during the initial surgical excision. In the early 2000s, IORT was a promising method of breast cancer treatment de-escalation, offering many advantages. Patients with early-stage breast cancer were entered into an institutional review board (IRB) approved registry. They were analyzed by intention to treat and by various subgroups, including those who received local treatment to the area of the primary tumor only versus those who received additional whole breast treatment. A total of 1785 patients with early-stage breast cancers entered the registry. With a median follow-up of 85 months, the local recurrence rate (LRR) at 5-years for the entire cohort was 4.44%. Among 1527 who received local treatment only, the LRR at 5-years was 5.09% compared with 1.13% for 301 patients who received additional whole breast treatment ($p = 0.001$). For patients aged ≥ 65 with luminal A tumors $\leq 20\text{mm}$, the 5-year LRR dropped to 2.32%. The authors concluded that IORT is extremely convenient for the patient and offers many advantages when compared with other methods of partial or whole breast treatment. Overall, the LRR for IORT is 4-5 times higher than many competing forms of whole or accelerated partial breast irradiation but still relatively low. A more select choice of patients for IORT can lower the LRR. Following the decision by ASTRO in 2024 not to recommend IORT unless within a clinical trial, and the above findings of a higher local recurrence rate in those treated with IORT, the group made the decision to discontinue their IORT program.

Section Summary: Intraoperative Radiotherapy

Several RCTs have been published, but they have not demonstrated that outcomes after intraoperative radiotherapy (IORT) are noninferior to WBI. Five-year results from the TARGIT-A RCT showed increased ipsilateral local recurrence with APBI compared with whole-breast radiotherapy. In a parallel study to TARGIT-A, delayed IORT was also associated with an increase in local recurrence rates at 5 years as compared to EBRT. In another RCT that used a different technology (ELIOT), recurrence rate with IORT was statistically greater than that with WBI. In a large registry study including 83,474 patients, no significant differences in cancer-specific survival or OS were observed between IORT and WBI.

External-Beam APBI

Systematic Reviews

An Agency for Healthcare Research and Quality (AHRQ) comparative effectiveness review assessed partial-breast irradiation (PBI) versus WBI in patients undergoing breast-conserving therapy for early-stage breast cancer.⁵² The analysis found no significant differences between PBI and WBI in terms of ipsilateral breast tumor recurrence, overall survival, or cancer-free survival at both 5 and 10-year follow-up with a high strength of evidence. In pre-specified subgroup analyses, AHRQ found no difference between APBI and non-accelerated PBI and observed that the results were consistent across various PBI modalities, including 3D conformal external beam radiation therapy (3DCRT), intensity-modulated radiation therapy (IMRT), and multi-catheter interstitial brachytherapy, when each was evaluated against WBI. PBI was associated with significantly fewer acute adverse events, while late toxicity rates were similar between groups (moderate strength of evidence).

Randomized Controlled Trials

Rodriguez et al (2013), reported on 102 patients were randomized to WBI, with or without a boost to the tumor bed, or APBI.⁵³ The primary end point was local recurrence within 5 years. In this noninferiority trial, the sample size was calculated to detect a 10% difference between treatment arms, with a power of 80% and a significance level of 0.05. The APBI group was significantly younger than the WBI group (mean age [SD], 67.1 years vs. 70.1 years; $p=0.009$). After a median follow-up of 5 years, there were no recurrences in either group, nor was there a statistically significant difference in survival. Investigators noted that the sample size may have been insufficient to detect a true difference in local control. Ninety percent (46/51) of APBI patients had acute skin effects, mostly grade 1; all patients in the WBI group had acute skin effects, and most were grade 2. Grade 1 and 2 late effects were reported with some changes in the relative positions of the treatment groups over time. Li et al (2021) reported long-term results of this trial (median, 10.3 years).⁵⁴ Rates of recurrence (4%) and disease-free survival (84%) were the same in both groups. Estimated 12-year OS was also similar between groups ($81.8\pm7.4\%$ APBI vs. $89.9\pm4.3\%$ WBI; $p>.05$). Grade 1 and 2 fibrosis was numerically more common than the ABPI group ($n=10$) than the WBI group ($n=4$; $p=.18$).

Olivotto et al (2013) reported interim results of the multicenter randomized RAPID trial (Randomized Trial of Accelerated Partial Breast Irradiation).⁵⁵ The sample size was 2135, and median follow-up was 3 years. Most patients were older than 50 years and had estrogen receptor–positive tumors less than 1.5 cm in diameter. An interim article reported on cosmetic and toxicity results. An accelerated regimen was used for WBI, and 21% of these patients received a boost to the tumor bed. APBI patients were more likely than WBI patients to have adverse cosmesis at 3 years, whether reported by physicians ($p<0.001$), nurses ($p<0.001$), or patients ($p<0.05$). As for late toxicities, 1.4% of APBI patients had a grade 3 adverse event versus none of the WBI patients. Telangiectasia and breast induration were more common among APBI patients ($p<0.001$). Although the primary outcome was ipsilateral local breast tumor recurrence, there were too few events to trigger an efficacy analysis. In 2019, Whelan et al published longer term results from RAPID.⁵⁶ Results from this analysis revealed similar ipsilateral breast tumor recurrence rates at 8 years between the groups (hazard ratio 1.27; 90% CI, 0.84 to 1.91) and no difference in OS (hazard ratio 1.18; 95% CI, 0.84 to 1.64).

In Livi et al (2015), 520 patients with early breast cancer were randomized to APBI using intensity-modulated radiotherapy or WBI.⁵⁷ The local recurrence rate at 5 years was 1.5% (3 cases) in the APBI group, there were 7 deaths in the WBI group and 1 in the APBI group ($p=0.057$). The 5-year OS was 96.6% for the WBI group and 99.4% for the APBI group. Long-term results (mean, 10.5 years; range, 1.4 to 14.8 years) were published by Meattini et al (2020).⁵⁸ The 10-year cumulative ipsilateral breast tumor recurrence rate was 2.5% with APBI and 3.7% with WBI (hazard ratio, 1.56; 95% CI, 0.55 to 4.37; $p=.40$). A similar number of deaths occurred in both groups (24 ABPI vs. 25 WBI) and the 10-year point estimate for OS was the same in both groups (91.9%; hazard ratio, 0.95; 95% CI, 0.50 to 1.79; $p=.86$).

Vicini et al (2019) completed a phase 3, equivalence, multicenter, RCT that compared APBI to WBI after breast-conserving surgery for early-stage breast cancer that enrolled 4216 patients.⁵⁹ Results revealed that, at a median follow-up of 10.2 years, ABPI did not meet the criteria for equivalence to WBI with regard to controlling ipsilateral breast tumor recurrence (hazard ratio, 1.22; 90% CI, 0.94 to 1.58); however, the absolute difference in the 10-year cumulative incidence of ipsilateral recurrence was $<1\%$ (4.6% APBI vs. 3.9% WBI).

Significantly more evaluable patients in the APBI group had recurrence-free interval events than patients in the WBI group (hazard ratio, 1.33; 95% CI, 1.04 to 1.69; $p=.02$); distant disease-free survival, OS, and disease-free survival were not different between the groups. The trial had broad eligibility criteria but was not designed to test equivalence in patient subgroups or outcomes from varying APBI techniques.

Polgar et al (2021) reported 20-year results of an RCT that compared APBI with either EBRT or high-dose interstitial brachytherapy ($n=128$) and WBI ($n=130$) in patients with early-stage breast cancer who had undergone breast-conserving surgery.⁶⁰ Patient accrual was stopped early and the study did not have sufficient power for the difference that was seen in the primary outcome (ipsilateral breast tumor recurrence). Median follow-up was 17 years (range, 1.5 to 21.2 years). Tumor recurrence rates were similar with APBI and WBI (9.6% vs. 7.9%, respectively; $p=.59$). Overall survival for 20 years was also similar between groups (59.5% vs. 59.7%, respectively; $p=.90$). Similar rates of grade 2 to 3 skin toxicity ($p=.32$) and fibrosis ($p=.16$) were reported in both groups.

Meduri et al (2023) reported 5-year outcomes of the IRMA trial.⁶¹ The IRMA trial randomly assigned women with early-stage breast cancer to breast-conserving surgery and WBI ($n=1657$) or twice-daily external-beam APBI ($n=1602$). At a median follow-up of 5.6 years, significantly higher rates of adverse cosmesis were reported in the APBI group (14% vs. 9.8%; $p=.012$) compared to WBI. Grade 3 late soft tissue (2.8% APBI vs. 1% WBI; $p<.0001$) and bone toxicities (1.1% APBI vs. 0% WBI) were significantly higher in the APBI arm. No significant differences in late skin or lung toxicities were observed. Five-year OS was 97.4% with APBI compared to 97.2% with WBI. The investigators plan to publish primary endpoint results (i.e., ipsilateral breast tumor recurrence) in a future study.

Tables 7 and 8 detail key characteristics and results of the RCTs summarized in this section.

Table 7. Summary of Key RCT Characteristics-External Beam APBI vs. WBI

Trial	Countries	Sites	Dates	Participants	Interventions	
Meduri et al (2023)	Italy	35	2007-2019	Age 49 years or older, with stage I-IIA breast cancer; tumor size <3 cm in diameter; with negative margins after breast-conserving therapy and a clinical target volume <30% of whole breast volume	APBI: 38.5 Gy total in 10 fractions (3.85 Gy/fraction), twice-daily for five consecutive days N=1602	WBI: 50.0 Gy in 25 fractions over 5 weeks N=1657
Polgar et al (2021)	Hungary	1	1998-2004	Low risk invasive breast carcinoma with negative margins after BCS	APBI: 5.2 Gy fractions given twice daily for 7 fractions with brachytherapy or 50 Gy total dose fractions over 5 weeks with EBRT N=128	WBI: 50 Gy given in 25 fractions over 5 weeks N=130
Vicini et al (2019)	U.S., Canada,	154	2005-2013	Over age 18, lumpectomy for	APBI: 34 Gy with	WBI: 50 Gy per day in

	Ireland, Israel			stage 0 cancer or stage I or II invasive adenocarcinoma of the breast with no distant metastases, life expectancy of at least 10 y; surgical resection margins needed to be cancer free	brachytherapy or 38.5 Gy with EBRT in 10 fractions given twice daily, at least 6 hours apart, on 5 treatment days within an 8-day period N=2107	25 total fractions spread over 5 weeks N=2109
Livi et al (2015) Meattini et al (2020)	Italy	1	2005-2013	Over age 40, maximum tumor size 25mm	APBI: 30 GY to the tumor bed in five daily fractions N=260	WBI: 50 Gy in 25 fractions, followed by a boost on the tumor bed of 10 Gy in five fractions N=260
Olivotto et al (2013) Whelan et al (2019)	Canada, Australia, New Zealand	33	2006-2011	Invasive ductal carcinoma or DCIS treated with BCS with microscopically clear margins and negative axillary nodes by sentinel node biopsy, or axillary dissection for those with invasive disease, or by clinical examination for those with DCIS alone	APBI: 38.5 Gy in 10 fractions treated twice daily over 5 to 8 days with a minimum interfraction interval of 6 hours N=1070	WBI: 42.5 Gy in 16 fractions or 50 Gy in 25 fractions. Boost irradiation of 10 Gy in 4-5 daily fractions after WBI was based on criteria such as young age or close margins N=1065
Rodriguez et al (2013) Li et al (2021)	Spain	1	NR	Invasive ductal carcinoma; age 60 or older; unifocal tumor; primary tumor size ≤30 mm	37.5 Gy in 3.75 Gy per fraction delivered twice daily. N=51	WBI: 48 Gy in daily fractions of 2 Gy, with or without additional 10 Gy to the tumor bed N=51

RCT: randomized controlled trial; APBI: accelerated partial breast irradiation; WBI: whole breast irradiation; Gy: Grays; N: sample size; BCS: breast-conserving surgery; DCIS: ductal carcinoma in situ.

Table 8. Summary of Key RCT Results-External Beam APBI vs. WBI

Study	Local Recurrence	Overall Survival	Toxicity
Meduri et al (2023)	Ipsilateral tumor recurrence at 5 years	5-year OS	Grade ≥3 late toxicity
N		3225	3225
APBI	NR	97.17%	Skin: 0.5% Soft Tissue: 2.8% (p<.0001) Lung: 0.1% Bone: 1.1% (p<.0001)
WBI	NR	97.44%	Skin: 0.4% Soft Tissue: 1.0% Lung: 0.2% Bone: 0.0%
Polgar et al (2021)	Ipsilateral tumor recurrence at 20 years	20-year OS	Grade 2-3 late toxicity

N	258		
APBI	9.6%	59.5%	Skin: 13.6% Fibrosis: 9.4%
WBI	7.9%	59.7%	Skin: 11.8% Fibrosis: 9.4%
Vicini et al (2019)	Ipsilateral tumor recurrence (first recurrence)	10-year point-estimate	CTCAE toxicity grade
N	4025		4109
ABPI	4%	90.6%	Grade 1: 40% Grade 2: 44% Grade 3: 10%
WBI	3%	91.3%	Grade 1: 31% Grade 2: 59% Grade 3: 7%
Livi et al (2015) Meattini et al (2020)	Ipsilateral tumor recurrence at 5 years	Number of deaths at 5 years	Acute skin toxicity (≥grade 2) at 5 years
N	520		520
APBI	1.5%	1	37.7%
WBI	1.4%	7	2%
Olivotto et al (2013) Whelan et al (2019)	Ipsilateral tumor recurrence at 8 years	Deaths	Grade 2 or 3 toxicity at 3 years
N	--	140	1070
APBI	3%	76	1.4%
WBI	2.8%	64	0%
Rodriguez et al (2013); Li et al (2021)			Acute and late toxicity
N	102	102	Acute: 102 4 years: 70
APBI	4%	81.8%	Acute: 46/51 (90.2%) 4 years: 16%, all grade 1
WBI	4%	89.9%	Acute: 51/51 (100%) 4 years: 11%, all grade 1

RCT: randomized controlled trial; APBI: accelerated partial breast irradiation; WBI: whole breast irradiation; Gy: Grays; N: sample size; BCS: breast-conserving surgery; DCIS: ductal carcinoma in situ. CTCAE: Common Terminology Criteria for Adverse Events

Relevance and study design limitations are summarized in Tables 9 and 10. The studies were limited by a lack of long-term follow-up data, small sample size, and incomplete follow-up data.

Table 9. Relevance Limitations-External Beam APBI vs. WBI

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Follow-Up ^e
Meduri et al (2023)				1. Primary endpoint (ipsilateral breast tumor recurrence) NR	1. 5-year outcomes
Polgar et al (2021)		3. 69% of patients received high-dose brachytherapy, 33% of patients received EBRT			
Vicini et al (2019)	4. Absence of HER2 data	3. 73% of patients had 3DCRT as			

	for enrolled patients with invasive breast cancer	their APBI technique; 27% underwent brachytherapy as the ABPI technique (either single-entry or multi-catheter)			
Livi et al (2015); Meattini et al (2020)				1.Overall survival NR	
Olivotto et al (2013) Whelan et al (2019)				1.Too few events for efficacy analysis of the primary outcome (local recurrence)	
Rodriguez et al (2013); Li et al (2021)					

APBI: accelerated partial breast irradiation; NR: not reported; NCT: national clinical trial; WBI: whole breast irradiation.

The evidence limitations stated in this table are those notable in the current review; this is not a comprehensive limitations assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

Table 10. Study Design and Conduct Limitations-External Beam APBI vs. WBI

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Meduri et al (2023)		1, 2. Not blinded to treatment assignment and outcome assessment				
Polgar et al (2021)		1,2	1. Protocol not registered			
Vicini et al (2019)		1,2				
Livi et al (2015); Meattini et al (2020)						
Olivotto et al (2013) Whelan et al (2019)		1		1.335/2135 (15.7%) completed 5-year assessment		

Rodriguez et al (2013); Li et al (2021)			1. Protocol not registered	1. Toxicity outcomes reported on 70/102 patients (68.6%)	3. May have been underpowered to detect difference in local recurrence rates	Trial terminated early due to cosmesis benefit
---	--	--	----------------------------	--	--	--

APBI: accelerated partial breast irradiation; NCT: national clinical trial; WBI: whole breast irradiation

The study limitations stated in this table are those notable in the current review; this is not a comprehensive limitations assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

^b Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

Section Summary: External Beam APBI

A comparative-effectiveness review from the Agency for Healthcare Research and Quality reported with high-strength evidence that partial-breast irradiation (PBI) offers equivalent ipsilateral recurrence, overall survival, and cancer-free survival rates to whole-breast irradiation (WBI) through 10 years of follow-up, and its pre-specified subgroup analysis revealed no outcome differences between APBI and conventional PBI. Across 6 randomized trials with 5 to 20-year follow-up, cumulative ipsilateral breast-tumor recurrence with APBI ranged 2.7% to 9.6% versus 2.8% to 7.9% with WBI; hazard ratios for recurrence or disease-free survival met pre-specified non-inferiority margins in most studies, although the largest trial was found non-equivalent despite an absolute 10-year difference of <1 %. Overall survival and distant disease-free survival were not significantly different in observations from 5 to 20 years. Several studies have reported that APBI produces higher rates of acute skin reactions, adverse cosmesis, and fibrosis/telangiectasia; however, serious grade ≥3 toxicities remain uncommon (<3%) and are comparable across modalities.

LOCAL BOOST BRACHYTHERAPY

Clinical Context and Therapy Purpose

The purpose of local boost brachytherapy with WBI in individuals who have early-stage breast cancer is to provide a treatment option that is an alternative to or an improvement on existing therapies.

Systematic Reviews

No RCTs were identified on local boost brachytherapy. However, there were 7 nonrandomized studies comparing 2 types of local boost radiotherapy: brachytherapy (n=2033) and EBRT (n=1557); all patients also received BCS and WBI. The combination of brachytherapy with local boost, BCS, and WBI prevented local tumor recurrence and salvage mastectomy in 95% to 97% of patients at 5 years and 88% to 92% of patients at 10 years. Five-year survival in the 5 studies reporting this outcome ranged from 83% to 96%. Data from uncontrolled studies reported similar rates of local control and 5-year survival.

Section Summary: Local Boost Brachytherapy

For women undergoing BCS plus WBI as initial treatment for stage I or II breast cancer, nonrandomized comparative studies have shown similar outcomes with brachytherapy with local boost and with EBRT with local boost.

NONINVASIVE BREAST BRACHYTHERAPY

Clinical Context and Therapy Purpose

The purpose of noninvasive breast brachytherapy in individuals who have early-stage breast cancer is to provide a treatment option that is an alternative to or an improvement on existing therapies.

AccuBoost for image-guided breast irradiation, also called noninvasive breast brachytherapy, has been used for local boost around the tumor bed. The AccuBoost system provides image-guided radiotherapy before each treatment to ensure that radiation is directed at the treatment target. The breast is placed between mammography paddles, where images are taken and radiation is delivered using a distinct applicator. The paddles prevent motion during treatment. Radiation is delivered from 1 side of the breast to the other or from the top of the breast to the bottom of the breast. This is proposed to reduce radiation exposure to adjacent tissues, including the heart and lung (Hamid et al, 2012).⁶² No long-term studies are available to confirm this.

Review of Evidence

No systematic reviews or RCTs of noninvasive breast brachytherapy for individuals with early-stage breast cancer were identified.

Nonrandomized Studies

One comparative study on noninvasive breast brachytherapy. This 2013 matched retrospective study assessed patients receiving the boost dose using AccuBoost or electron beams (a type of EBRT) [Leonard et al, 2013].⁶³ Each of 47 AccuBoost patients was compared with 2 controls matched on age, stage, chemotherapy use, fractionation, and when possible, breast size, comorbidities, and smoking status. Main differences between the 2 treatment groups were in radiation doses received and timing of radiotherapy administration. The percentage of patients with a WBI dose (accompanying the boost dose) of 50 to 50.4 Gy was 68% in the AccuBoost group and 37% in the electron-treated group ($p < 0.001$). Also, a greater proportion of patients in the electron-treated group received the boost dose after WBI, rather than during WBI or starting before and ending during WBI (99% for the electron-treated group vs. 6% for the AccuBoost group). Approximately 60% of patients had stage I breast cancer, and approximately 25%, ductal carcinoma in situ. With median follow-up of 13.6 months, skin and subcutaneous tissue toxicity occurred less often among patients treated with AccuBoost than among those treated with electron beam ($p = 0.046$). Locoregional control rates were 99% or greater in both groups. Study limitations included the between-group differences in dose and timing of boost, as well as selection bias and the study's retrospective design.

Section Summary: Noninvasive Breast Brachytherapy

One nonrandomized comparative study was identified. The comparative study was retrospective matched comparison of noninvasive breast brachytherapy or electron-beam radiotherapy to provide boost radiation to the tumor bed. The study was subject to selection bias, relatively short follow-up, and use of a retrospective design.

SUPPLEMENTAL INFORMATION

CLINICAL INPUT FROM PHYSICIAN SPECIALTY SOCIETIES, ACADEMIC MEDICAL CENTERS AND EXTERNAL STAKEHOLDERS (PRACTICING CLINICIANS AND RESEARCH BASED EXPERTS IN THE FIELD)

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

2017 Input

Clinical input was sought to help determine whether the use of accelerated whole breast irradiation (AWBI) for individuals with node-negative, early-stage breast cancer with clear surgical margins would provide a clinically meaningful improvement in net health outcome and whether the use is consistent with generally accepted medical practice. In response to requests, clinical input was received from 5 respondents, including no specialty society-level response(s) including physicians with academic medical center affiliation; 1 physician-level responses identified through a specialty society; 4 physician-level responses identified through an academic medical center.

For individuals who have node-negative, early-stage breast cancer with clear surgical margins who receive AWBI, clinical input supports this use provides a clinically meaningful improvement in net health outcome and indicates this use is consistent with generally accepted medical practice. Input was limited to the policy statement on AWBI. Three of 4 academic medical centers and the physician specialty society agreed with the statement as a whole. Reviewers suggested other eligibility criteria but there was no consensus on specific criteria.

PRACTICE GUIDELINES AND POSITION STATEMENTS

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they are issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

National Comprehensive Cancer Network

Current National Comprehensive Cancer Network (NCCN) guidelines (version 5.2026) on breast cancer state:

APBI/PBI is endorsed for any patient without germline BRCA 1/2 mutations who meets the criteria outlined in the 2024 ASTRO guidelines. Patients aged ≥ 40 years are recommended for APBI/PBI if they have:

- Invasive ER-positive ductal carcinoma measuring ≤ 2 cm (pT1 disease), grade 1–2, with negative margin widths, no lymph-vascular space invasion, and negative nodes; OR
- Ductal carcinoma in situ (DCIS) measuring size ≤ 2 cm with low-intermediate grade with negative margins. APBI offers comparable local control and comparable or improved cosmesis to whole breast radiotherapy when delivered with the following dose regimens. The APBI regimens have not been compared directly but 30 Gy/5 fractions is preferred based on the highest rated cosmesis outcomes.

Radiotherapy Dosing:

- External Beam Radiation Therapy APBI
 - 30 Gy/5 fractions QOD (preferred); Intensity-modulated radiation therapy/Volumetric modulated arc therapy protocol mandated;
 - 40 Gy/15 fractions
- Brachytherapy APBI (including balloon/interstitial)
 - 34 Gy/10 fractions BID;
 - 32 Gy/8 fractions BID;
 - 30.1 Gy/7 fractions BID"

For whole-breast radiotherapy, the NCCN recommends a “hypofractionated dose of 40–42.5 Gy in 15–16 fractions; in selected cases 45–50.4 Gy in 25–28 fractions may be considered. A boost to the tumor bed is recommended in patients at higher risk for recurrence. The boost can be given sequentially after whole breast RT or as a simultaneous integrated boost. Typical boost doses when given sequentially are 10–16 Gy in 4–8 fractions. When given concurrently, the whole breast should receive 40 Gy in 15 fractions, and the lumpectomy site should receive 48 Gy in 15 fractions. Available data support the use of ultra-hypofractionated regimens (28.5 Gy in 5 once-a-week fractions or 26 Gy and 5 daily fractions). The optimal way to incorporate a tumor bed boost and the long-term toxicity associated with its delivery in the ultra-hypofractionated setting is yet to be established. Lumpectomy cavity boost can be delivered using enface electrons, photons, or brachytherapy.

American Society for Radiation Oncology et al

The American Society for Radiation Oncology (ASTRO, 2024) published guidance on the use of PBI in individuals with early-stage invasive breast cancer or DCIS and issued the following recommendations:

Indications for PBI as an alternative to WBI in early-stage invasive breast cancer:

1. PBI is recommended for patients with early-stage invasive breast cancer with all of the following factors: Grade 1-2 disease, ER-positive histology, Age ≥ 40 years, Tumor size ≤ 2 cm. Strength of recommendation: Strong; Quality of evidence: High (for grade, histology, & age ≥ 40 years) and Moderate (for age 40-49 years and size)
2. PBI is conditionally recommended for patients with early-stage invasive breast cancer with the following factors: Grade 3 disease or ER-negative histology or Size $>2 - \leq 3$ cm. Strength of recommendation: Conditional; Quality of evidence: Low
3. PBI is conditionally **not** recommended for patients with early-stage invasive breast cancer with any of the following factors: HER2-positive tumors not receiving anti-HER2 therapy, Lymphovascular invasion, or Lobular histology. Strength of recommendation: Conditional; Quality of evidence: Expert opinion
4. PBI is **not** recommended for patients with early-stage invasive breast cancer with any of the following factors: Positive lymph nodes, Positive surgical margins, Known germline BRCA1/2 mutation, and Age <40 years. Strength of recommendation: Strong; Quality of evidence: Expert opinion

Indications for PBI as an alternative to WBI in DCIS:

1. PBI is recommended for patients with DCIS with all of the following factors: Low to intermediate grade, Age ≥ 40 years, Size ≤ 2 cm. Strength of recommendation: Strong; Quality of evidence: Expert opinion.
2. PBI is conditionally recommended for patients with DCIS with the following factors: High grade, Size $>2 - \leq 3$ cm. Strength of recommendation: Conditional; Quality of evidence: Expert opinion.

3. PBI is **not** recommended for patients with DCIS with any of the following factors:
Positive surgical margins, Known germline BRCA1/2 mutation, Age <40 years. Strength of recommendation: Strong; Quality of evidence: Expert opinion.

Appropriate PBI techniques with respect to rates of Ipsilateral Breast Recurrence:

1. For patients with early-stage invasive breast cancer or DCIS receiving PBI, 3-D CRT is recommended. Strength of recommendation: Strong; Quality of evidence: High
2. For patients with early-stage invasive breast cancer or DCIS receiving PBI, IMRT is recommended. Strength of recommendation: Strong; Quality of evidence: Moderate
3. For patients with early-stage invasive breast cancer or DCIS receiving PBI, multicatheter brachytherapy is recommended. Strength of recommendation: Strong; Quality of evidence: Moderate
4. For patients with early-stage invasive breast cancer or DCIS receiving PBI, single-entry catheter brachytherapy is conditionally recommended. Strength of recommendation: Conditional; Quality of evidence: Moderate
5. For patients with early-stage invasive breast cancer receiving PBI, electron IORT is **not** recommended, unless part of a clinical trial or multi-institutional registry. Strength of recommendation: Strong; Quality of evidence: Moderate
6. For patients with early-stage invasive breast cancer receiving PBI, kV IORT alone (without WBI) is **not** recommended, unless part of a clinical trial or multi-institutional registry. Strength of recommendation: Strong; Quality of evidence: Low

ASTRO (2023), American Society of Breast Surgeons (2018), and the American Brachytherapy Society (2018) have issued various consensus statements for the selection of patients for APBI (summarized in Table 11).

Table 11. Professional Medical Society Criteria for Performing APBI

Factor	ASTRO" Suitable" (2016)	ASTRO" Cautionary" (2016)	ASTRO" Unsuitable" (2016)	ASTRO Recommended (2023)	ASTRO Conditionally Recommended (2023)	ASTRO Conditionally Not Recommended (2023)	ASTRO Not Recommended (2023)	ASBS	ABS
Patient factors									
Age	≥50 y	40 to 49 y; ≥50 y if patient has at least 1 of the pathologic factors and does not have any "unsuitable" factors	<40 y; 40 to 49y and do not meet the criteria for cautionary	≥40 y	NR	NR	<40 y	≥45 y for all tumor types	≥45 y
BRCA1 and BRCA2 variants	Not present	NR	Present	NR	NR	NR	Present	Patients should not be treated if they have a BRCA genetic mutation	NR
Pathologic factors									
Tumor size	≤2 cm	2.1 to 3.0 cm	>3 cm	≤2 cm	>2 to ≤3 cm	NR	NR	≤3 cm	≤3 cm
Tumor stage	Tis or T1	T0 or T2	T3-4	NR	NR	NR	NR	NR	Tis, T1, T2 (≤3cm)
Margins	Negative ≥2 mm	Close (<2mm)	Positive	NR	NR	NR	Positive	No tumor on ink for invasive tumors or tumors	Negative (no tumor on ink for invasive)

								involved with DCIS; ≥ 2 mm for DCIS	≥ 2 mm for DCIS)
Grade	Any	NR	NR	1 or 2	3	NR	NR	NR	NR
LVSI	No	Limited/focal	Extensive	NR	NR	Yes	NR	Allowed as long as it is focal	Not present
ER status	Positive	Negative	NR	Positive	Negative	HER2 positive and not receiving anti-HER2 therapy	NR	Positive or negative	Positive or negative
Multicentricity	Unicentric	NR	Present	NR	NR	NR	NR	NR	NR
Multifocality	Clinically unifocal; total size, ≤ 2.0 cm	Clinically unifocal; size, 2.1-3.0 cm	Clinically multifocal or microscopically multifocal; size, ≥ 3 cm	NR	NR	NR	NR	Multifocal disease is allowed as long as the combined area of tumor is ≤ 3 cm	NR
Histology	Invasive ductal or other favorable subtypes	Invasive lobular	NR	Invasive	Invasive	Invasive lobular	NR	All invasive subtypes; DCIS	All invasive subtypes and DCIS
Pure DCIS	Not allowed ^a	≤ 3 cm if "suitable" criteria not fully met	> 3 cm	NR	NR	NR	NR	≤ 3 cm	≤ 3 cm

ABS: American Brachytherapy Society; ALND: axillary lymph node dissection; APBI: accelerated partial-breast irradiation; ASBS: American Society of Breast Surgeons; ASTRO: American Society for Radiation Oncology; DCIS: ductal carcinoma in situ; EIC: extensive intraductal component; ER: estrogen receptor; LCIS: lobular carcinoma in situ; LVSI: lymphovascular space invasion; NR: not reported; SN: sentinel node; SNB: sentinel node biopsy.

^a Strongly encouraged to enroll in NSABP B-39/RTOG 04-13 trial.

^b Allowed if screen-detected, low to intermediate nuclear grade, ≤ 2.5 cm size, and resected with margins negative at ≥ 3 mm.

^c Lymphovascular space invasion is considered a contraindication for accelerated partial-breast irradiation.

The ASTRO (2018) updated its guidelines on fractionation for whole-breast irradiation. The consensus-based guidelines conclude that accelerated whole-breast irradiation may be used for any age, and any stage provided the intent is to treat the whole breast without any additional field, and with any chemotherapy.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 12.

Table 12. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing			
WBI vs. APBI with or without tumor bed boost in DCIS			
NCT04669873	Clinical Trial, Randomized, Open Label, With an Active Comparator to Assess the Efficacy and Safety of Using Accelerated Partial Irradiation Versus Standard or Hypofractionated Irradiation of the Entire Breast in Patients with Initial Breast Cancer After Conservative Surgery	36	Dec 2026 (recruiting)
Intraoperative Radiotherapy			
NCT06375798	Clinical Study of Breast Conserving Surgery Combined with Intraoperative Radiotherapy for Early Breast Cancer	620	Nov 2026 (recruiting)

NCT01644669	Safety and efficacy study of the Xofigo® Axxent® eBx™ IORT system	1200	Dec 2029 (active not recruiting)
External-beam APBI			
NCT01247233	Standard or hypofractionated radiotherapy vs. accelerated partial breast irradiation (APBI) for breast cancer (SHARE)	1006	Oct 2025 (active, not recruiting)
NCT06185205	Accelerated Super-Hypofractionated Breast Brachytherapy-ASHBY Trial	60	Jan 2033 (recruiting)
NCT01185132	Intensity modulated radiotherapy (IMRT) vs. 3D-conformal accelerated partial breast irradiation (APBI) for early-stage breast cancer after lumpectomy (2009-APBI)	660	Jul 2028 (recruiting)
APBI (multimodality)			
NCT05914831	Ultra-hypofractionated for Whole Breast Irradiation (WBI) Compared to Partial Breast Irradiation (PBI): A Single-Institution Prospective Phase 2 Trial	100	May 2033 (recruiting)
NCT05472792	Comparison of Adjuvant Monotherapy with Endocrine Therapy or Accelerated Partial Breast Irradiation following Lumpectomy for Low-Risk Breast Cancer Patients Over 65 (CAMERAN) (CAMERAN)	90	May 2032 (recruiting)
NCT00185744	Accelerated Partial Breast Irradiation Following Lumpectomy for Breast Cancer	400	Mar 2029 (active not recruiting)
NCT04852887	A Phase III Clinical Trial Evaluating De-Escalation of Breast Radiation for Conservative Treatment of Stage I, Hormone Sensitive, HER-2 Negative, Oncotype Recurrence Score Less Than or Equal to 18 Breast Cancer	1670	Jul 2041 (recruiting)

APBI: accelerated partial-breast irradiation; DCIS: ductal carcinoma in situ; NCT: national clinical trial; WBI: whole-breast irradiation

Regulatory Status:

In 2002, the MammoSite® Radiation Therapy System (Proxima Therapeutics; Alpharetta, GA), the first device specifically designed for breast brachytherapy,¹³ was cleared for marketing by the U. S. Food and Drug Administration (FDA) through the 510(k) process. Its intended use is “to provide brachytherapy when the physician chooses to deliver intracavitary radiation to the surgical margins following lumpectomy for breast cancer.”¹⁴

Since 2002, several other devices for breast brachytherapy have cleared for marketing by the FDA through the 510(k) process. The FDA determined that several devices (e.g., Axxent® Electronic Brachytherapy System [Xoft; San Jose, CA], Strut-Adjusted Volume Implant [SAVI™] Applicator Kit [Biolument (now Cianna Medical); Aliso Viejo, CA], Contura® Multi-Lumen Balloon Source Applicator for Brachytherapy [SenoRx; Aliso Viejo, CA], ClearPath™ Adjustable Multi-Catheter Source Applicator [North American Scientific; Chatsworth, CA], Intrabeam® System [Carl Zeiss Surgical, GmbH; Oberkochen, Germany]) were substantially equivalent to predicate devices. Each includes an FDA-required warning that the safety and effectiveness of the device “as a replacement for whole-breast irradiation in the treatment of breast cancer has not been established.”

Although the Intrabeam® System (discussed in the Intraoperative Radiotherapy subsection) is subject to FDA regulation, it does not fall under the regulatory purview of the U.S. Nuclear

Regulatory Commission. In some states, participation of radiation oncologists in delivering radiation is not required.

Government Regulations

National:

There is no national coverage determination on this topic. In the absence of an NCD, coverage decisions are left to the discretion of local Medicare carriers.

Local:

Wisconsin Physicians Service Insurance Corporation has a local coverage determination on brachytherapy (L30320); however, it is not specific for treatment of the breast. Policy was retired 6/1/15 and not replaced.

(The above Medicare information is current as of the review date for this policy. However, the coverage issues and policies maintained by the Centers for Medicare & Medicare Services [CMS, formerly HCFA] are updated and/or revised periodically. Therefore, the most current CMS information may not be contained in this document. For the most current information, the reader should contact an official Medicare source.)

Related Policies

- Brachytherapy for Prostate Cancer
 - Brachytherapy of the Femoropopliteal Arterial System after Percutaneous Transluminal Angioplasty (PTA)
 - Electronic Brachytherapy
 - Intraoperative Radiation Therapy
-

References

1. Moran MS, Schnitt SJ, Giuliano AE, et al. Society of Surgical Oncology-American Society for Radiation Oncology consensus guideline on margins for breast-conserving surgery with whole-breast irradiation in stages I and II invasive breast cancer. *Int J Radiat Oncol Biol Phys*. Mar 01, 2014; 88(3): 553-64. PMID 24521674
2. National Cancer Institute, Surveillance Epidemiology and End Results Program. Cancer Stat Facts: Female Breast Cancer. n.d.; <https://seer.cancer.gov/statfacts/html/breast.html>. Accessed May 23, 2025.
3. Darby S, McGale P, Correa C, et al. Effect of radiotherapy after breast-conserving surgery on 10-year recurrence and 15-year breast cancer death: meta-analysis of individual patient data for 10,801 women in 17 randomised trials. *Lancet*. Nov 12, 2011; 378(9804): 1707-16. PMID 22019144
4. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Breast Cancer. Version 5.2026. https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf. Accessed June 2026.
5. Albert JM, Pan IW, Shih YC, et al. Effectiveness of radiation for prevention of mastectomy in older breast cancer patients treated with conservative surgery. *Cancer*. Oct 01, 2012; 118(19): 4642-51. PMID 22890779

6. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Accelerated radiotherapy after breast-conserving surgery for early-stage breast cancer. TEC Assessments. 2013;Volume 27:Tab 6.
7. Kaidar-Person O, Yarnold J, Offersen BV, et al. Is current evidence about intraoperative partial breast irradiation sufficient for broad implementation in clinical practice? *Eur J Surg Oncol*. Jul 2014; 40(7): 791-3. PMID 24721228
8. Silverstein MJ, Fastner G, Maluta S, et al. Intraoperative radiation therapy: a critical analysis of the ELIOT and TARGIT trials. Part 2--TARGIT. *Ann Surg Oncol*. Nov 2014; 21(12): 3793-9. PMID 25138079
9. Silverstein MJ, Fastner G, Maluta S, et al. Intraoperative radiation therapy: a critical analysis of the ELIOT and TARGIT trials. Part 1--ELIOT. *Ann Surg Oncol*. Nov 2014; 21(12): 3787-92. PMID 25160734
10. Athas WF, Adams-Cameron M, Hunt WC, et al. Travel distance to radiation therapy and receipt of radiotherapy following breast-conserving surgery. *J Natl Cancer Inst*. Feb 02, 2000; 92(3): 269-71. PMID 10655446
11. Farrow DC, Hunt WC, Samet JM. Geographic variation in the treatment of localized breast cancer. *N Engl J Med*. Apr 23, 1992; 326(17): 1097-101. PMID 1552910
12. Nattinger AB, Kneusel RT, Hoffmann RG, et al. Relationship of distance from a radiotherapy facility and initial breast cancer treatment. *J Natl Cancer Inst*. Sep 05, 2001; 93(17): 1344-6. PMID 11535710
13. Dragun AE, Huang B, Tucker TC, et al. Disparities in the application of adjuvant radiotherapy after breast-conserving surgery for early-stage breast cancer: impact on overall survival. *Cancer*. Jun 15, 2011; 117(12): 2590-8. PMID 21656737
14. Berliner E. Adopting medical technology. *Med Decis Making*. Nov 2014; 34(8): 948-50. PMID 25224365
15. Food and Drug Administration. 510(K) summary of safety and effectiveness: MammoSite Radiation Therapy System. 2002; <https://www.accessdata.fda.gov/scripts/cdrh/devicesatfda/index.cfm?db=pmn&id=K011690>. Accessed May 23, 2025. In supplemental documentation folder.
16. Valle LF, Agarwal S, Bickel KE, et al. Hypofractionated whole breast radiotherapy in breast conservation for early-stage breast cancer: a systematic review and meta-analysis of randomized trials. *Breast Cancer Res Treat*. Apr 2017; 162(3): 409-417. PMID 28160158
17. Bentzen SM, Agrawal RK, Aird EG, et al. The UK Standardisation of Breast Radiotherapy (START) Trial B of radiotherapy hypofractionation for treatment of early breast cancer: a randomised trial. *Lancet*. Mar 29, 2008; 371(9618): 1098-107. PMID 18355913
18. Whelan T, MacKenzie R, Julian J, et al. Randomized trial of breast irradiation schedules after lumpectomy for women with lymph node-negative breast cancer. *J Natl Cancer Inst*. Aug 07, 2002; 94(15): 1143-50. PMID 12165639
19. Whelan TJ, Pignol JP, Levine MN, et al. Long-term results of hypofractionated radiation therapy for breast cancer. *N Engl J Med*. Feb 11, 2010; 362(6): 513-20. PMID 20147717
20. Hopwood P, Haviland JS, Sumo G, et al. Comparison of patient-reported breast, arm, and shoulder symptoms and body image after radiotherapy for early breast cancer: 5-year follow-up in the randomised Standardization of Breast Radiotherapy (START) trials. *Lancet Oncol*. Mar 2010; 11(3): 231-40. PMID 20138809
21. Haviland JS, Owen JR, Dewar JA, et al. The UK Standardisation of Breast Radiotherapy (START) trials of radiotherapy hypofractionation for treatment of early

- breast cancer: 10-year follow-up results of two randomised controlled trials. *Lancet Oncol.* Oct 2013; 14(11): 1086-1094. PMID 24055415
22. Shaitelman SF, Schlembach PJ, Arzu I, et al. Acute and Short-term Toxic Effects of Conventionally Fractionated vs Hypofractionated Whole-Breast Irradiation: A Randomized Clinical Trial. *JAMA Oncol.* Oct 2015; 1(7): 931-41. PMID 26247543
 23. Brunt AM, Haviland JS, Sydenham M, et al. Ten-Year Results of FAST: A Randomized Controlled Trial of 5-Fraction Whole-Breast Radiotherapy for Early Breast Cancer. *J Clin Oncol.* Oct 01, 2020; 38(28): 3261-3272. PMID 32663119
 24. Murray Brunt A, Haviland JS, Wheatley DA, et al. Hypofractionated breast radiotherapy for 1 week versus 3 weeks (FAST-Forward): 5-year efficacy and late normal tissue effects results from a multicentre, non-inferiority, randomised, phase 3 trial. *Lancet.* May 23, 2020; 395(10237): 1613-1626. PMID 32580883
 25. Chan EK, Woods R, McBride ML, et al. Adjuvant hypofractionated versus conventional whole breast radiation therapy for early-stage breast cancer: long-term hospital-related morbidity from cardiac causes. *Int J Radiat Oncol Biol Phys.* Mar 15, 2014; 88(4): 786-92. PMID 24606848
 26. Chan EK, Woods R, Virani S, et al. Long-term mortality from cardiac causes after adjuvant hypofractionated vs. conventional radiotherapy for localized left-sided breast cancer. *Radiother Oncol.* Jan 2015; 114(1): 73-8. PMID 25227961
 27. Kong L, Cheng J, Ding X, et al. Efficacy and safety of accelerated partial breast irradiation after breast-conserving surgery: a meta-analysis of published comparative studies. *Breast J.* 2014; 20(2): 116-24. PMID 24372818
 28. Marta GN, Macedo CR, Carvalho Hde A, et al. Accelerated partial irradiation for breast cancer: systematic review and meta-analysis of 8653 women in eight randomized trials. *Radiother Oncol.* Jan 2015; 114(1): 42-9. PMID 25480094
 29. Lv Y, He L, Wang C, et al. A systematic review of clinical outcomes and radiotherapy-associated toxicity in multicatheter accelerated partial breast irradiation. *Medicine (Baltimore).* Feb 2019; 98(6): e14407. PMID 30732191
 30. Liu L, Yang Y, Guo Q, et al. Comparing hypofractionated to conventional fractionated radiotherapy in postmastectomy breast cancer: a meta-analysis and systematic review. *Radiat Oncol.* Jan 17, 2020; 15(1): 17. PMID 31952507
 31. Hausmann J, Budach W, Corradini S, et al. No Difference in Overall Survival and Non-Breast Cancer Deaths after Partial Breast Radiotherapy Compared to Whole Breast Radiotherapy-A Meta-Analysis of Randomized Trials. *Cancers (Basel).* Aug 17, 2020; 12(8). PMID 32824414
 32. Xiang X, Ding Z, Feng L, et al. A meta-analysis of the efficacy and safety of accelerated partial breast irradiation versus whole-breast irradiation for early-stage breast cancer. *Radiat Oncol.* Feb 02, 2021; 16(1): 24. PMID 33531014
 33. Shah C, Jia X, Hobbs BP, et al. Outcomes with Partial Breast Irradiation vs. Whole Breast Irradiation: A Meta-Analysis. *Ann Surg Oncol.* Sep 2021; 28(9): 4985-4994. PMID 33393051
 34. Hickey BE, Lehman M. Partial breast irradiation versus whole breast radiotherapy for early breast cancer. *Cochrane Database Syst Rev.* Aug 30, 2021; 8(8): CD007077. PMID 34459500
 35. Viani GA, Arruda CV, Faustino AC, et al. Partial-breast irradiation versus whole-breast radiotherapy for early breast cancer: A systematic review and update meta-analysis. *Brachytherapy.* 2020; 19(4): 491-498. PMID 32340902
 36. Korzets Y, Fyles A, Shepshelovich D, et al. Toxicity and clinical outcomes of partial breast irradiation compared to whole breast irradiation for early-stage breast cancer: a

- systematic review and meta-analysis. *Breast Cancer Res Treat.* Jun 2019; 175(3): 531-545. PMID 30929116
37. Strnad V, Ott OJ, Hildebrandt G, et al. 5-year results of accelerated partial breast irradiation using sole interstitial multicatheter brachytherapy versus whole-breast irradiation with boost after breast-conserving surgery for low-risk invasive and in-situ carcinoma of the female breast: a randomised, phase 3, non-inferiority trial. *Lancet.* Jan 16, 2016; 387(10015): 229-38. PMID 26494415
 38. Polgár C, Ott OJ, Hildebrandt G, et al. Late side-effects and cosmetic results of accelerated partial breast irradiation with interstitial brachytherapy versus whole-breast irradiation after breast-conserving surgery for low-risk invasive and in-situ carcinoma of the female breast: 5-year results of a randomised, controlled, phase 3 trial. *Lancet Oncol.* Feb 2017; 18(2): 259-268. PMID 28094198
 39. Schäfer R, Strnad V, Polgár C, et al. Quality-of-life results for accelerated partial breast irradiation with interstitial brachytherapy versus whole-breast irradiation in early breast cancer after breast-conserving surgery (GEC-ESTRO): 5-year results of a randomised, phase 3 trial. *Lancet Oncol.* Jun 2018; 19(6): 834-844. PMID 29695348
 40. Strnad V, Polgár C, Ott OJ, et al. Accelerated partial breast irradiation using sole interstitial multicatheter brachytherapy compared with whole-breast irradiation with boost for early breast cancer: 10-year results of a GEC-ESTRO randomised, phase 3, non-inferiority trial. *Lancet Oncol.* Mar 2023; 24(3): 262-272. PMID 36738756
 41. Kim H, Hieken TJ, Abraha F, et al. Long-term outcomes of intraoperatively-placed applicator brachytherapy for rapid completion of breast conserving treatment: An analysis of prospective registry data. *Clin Transl Radiat Oncol.* Jul 2023; 41: 100639. PMID 37251618
 42. Ajkay N, Collett AE, Bloomquist EV, et al. A comparison of complication rates in early-stage breast cancer patients treated with brachytherapy versus whole-breast irradiation. *Ann Surg Oncol.* Apr 2015; 22(4): 1140-5. PMID 25319575
 43. Ravani LV, Calomeni P, Wang M, et al. Comparison of partial-breast irradiation and intraoperative radiation to whole-breast irradiation in early-stage breast cancer patients: a Kaplan-Meier-derived patient data meta-analysis. *Breast Cancer Res Treat.* Jan 2024; 203(1): 1-12. PMID 37736843
 44. Wang L, Sun M, Yang S, et al. Intraoperative Radiotherapy Is Not a Better Alternative to Whole Breast Radiotherapy as a Therapeutic Option for Early-Stage Breast Cancer. *Front Oncol.* 2021; 11: 737982. PMID 34976796
 45. Vaidya JS, Joseph DJ, Tobias JS, et al. Targeted intraoperative radiotherapy versus whole breast radiotherapy for breast cancer (TARGIT-A trial): an international, prospective, randomised, non-inferiority phase 3 trial. *Lancet.* Jul 10, 2010; 376(9735): 91-102. PMID 20570343
 46. Vaidya JS, Wenz F, Bulsara M, et al. Risk-adapted targeted intraoperative radiotherapy versus whole-breast radiotherapy for breast cancer: 5-year results for local control and overall survival from the TARGIT-A randomised trial. *Lancet.* Feb 15, 2014; 383(9917): 603-13. PMID 24224997
 47. Vaidya JS, Wenz F, Bulsara M, et al. An international randomised controlled trial to compare TARGeted Intraoperative radioTherapy (TARGIT) with conventional postoperative radiotherapy after breast-conserving surgery for women with early-stage breast cancer (the TARGIT-A trial). *Health Technol Assess.* Sep 2016; 20(73): 1-188. PMID 27689969
 48. Vaidya JS, Bulsara M, Saunders C, et al. Effect of Delayed Targeted Intraoperative Radiotherapy vs Whole-Breast Radiotherapy on Local Recurrence and Survival: Long-

- term Results From the TARGIT-A Randomized Clinical Trial in Early Breast Cancer. *JAMA Oncol.* Jul 01, 2020; 6(7): e200249. PMID 32239210
49. Veronesi U, Orecchia R, Luini A, et al. Intraoperative radiotherapy during breast conserving surgery: a study on 1,822 cases treated with electrons. *Breast Cancer Res Treat.* Nov 2010; 124(1): 141-51. PMID 20711810
 50. Veronesi U, Orecchia R, Maisonneuve P, et al. Intraoperative radiotherapy versus external radiotherapy for early breast cancer (ELIOT): a randomised controlled equivalence trial. *Lancet Oncol.* Dec 2013; 14(13): 1269-77. PMID 24225155
 51. Orecchia R, Veronesi U, Maisonneuve P, et al. Intraoperative irradiation for early breast cancer (ELIOT): long-term recurrence and survival outcomes from a single centre, randomised, phase 3 equivalence trial. *Lancet Oncol.* May 2021; 22(5): 597-608. PMID 33845035
 52. Shumway DA, Corbin KS, Farah MH, Viola KE, Nayfeh T, Saadi S, Shah V, Hasan B, Shah S, Mohammed K, Riaz IB, Prokop LJ, Wang Z, Murad MH. Partial Breast Irradiation for Breast Cancer [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (US); 2023 Jan. Report No.: 23-EHC001. PMID: 36753597
 53. Rodríguez N, Sanz X, Dengra J, et al. Five-year outcomes, cosmesis, and toxicity with 3-dimensional conformal external beam radiation therapy to deliver accelerated partial breast irradiation. *Int J Radiat Oncol Biol Phys.* Dec 01, 2013; 87(5): 1051-7. PMID 24161420
 54. Li X, Sanz J, Foro P, et al. Long-term results of a randomized partial irradiation trial compared to whole breast irradiation in the early stage and low-risk breast cancer patients after conservative surgery. *Clin Transl Oncol.* Oct 2021; 23(10): 2127-2132. PMID 33880724
 55. Olivotto IA, Whelan TJ, Parpia S, et al. Interim cosmetic and toxicity results from RAPID: a randomized trial of accelerated partial breast irradiation using three-dimensional conformal external beam radiation therapy. *J Clin Oncol.* Nov 10, 2013; 31(32): 4038-45. PMID 23835717
 56. Whelan TJ, Julian JA, Berrang TS, et al. External beam accelerated partial breast irradiation versus whole breast irradiation after breast conserving surgery in women with ductal carcinoma in situ and node-negative breast cancer (RAPID): a randomised controlled trial. *Lancet.* Dec 14, 2019; 394(10215): 2165-2172. PMID 31813635
 57. Livi L, Meattini I, Marrazzo L, et al. Accelerated partial breast irradiation using intensity-modulated radiotherapy versus whole breast irradiation: 5-year survival analysis of a phase 3 randomised controlled trial. *Eur J Cancer.* Mar 2015; 51(4): 451-463. PMID 25605582
 58. Meattini I, Marrazzo L, Saieva C, et al. Accelerated Partial-Breast Irradiation Compared with Whole-Breast Irradiation for Early Breast Cancer: Long-Term Results of the Randomized Phase III APBI-IMRT-Florence Trial. *J Clin Oncol.* Dec 10, 2020; 38(35): 4175-4183. PMID 32840419
 59. Vicini FA, Cecchini RS, White JR, et al. Long-term primary results of accelerated partial breast irradiation after breast-conserving surgery for early-stage breast cancer: a randomised, phase 3, equivalence trial. *Lancet.* Dec 14, 2019; 394(10215): 2155-2164. PMID 31813636
 60. Polgár C, Major T, Takácsi-Nagy Z, et al. Breast-Conserving Surgery Followed by Partial or Whole Breast Irradiation: Twenty-Year Results of a Phase 3 Clinical Study. *Int J Radiat Oncol Biol Phys.* Mar 15, 2021; 109(4): 998-1006. PMID 33186620
 61. Meduri B, Baldissera A, Iotti C, et al. Cosmetic Results and Side Effects of Accelerated Partial-Breast Irradiation Versus Whole-Breast Irradiation for Low-Risk Invasive

- Carcinoma of the Breast: The Randomized Phase III IRMA Trial. J Clin Oncol. Apr 20, 2023; 41(12): 2201-2210. PMID 36623246
62. Hamid S, Rocchio K, Arthur D, et al. A multi-institutional study of feasibility, implementation, and early clinical results with noninvasive breast brachytherapy for tumor bed boost. Int J Radiat Oncol Biol Phys. Aug 01, 2012; 83(5): 1374-80. PMID 22209153
 63. Leonard KL, Hepel JT, Styczynski JR, et al. Breast boost using noninvasive image-guided breast brachytherapy vs. external beam: a 2:1 matched-pair analysis. Clin Breast Cancer. Dec 2013; 13(6): 455-9. PMID 24099648
 64. Shaitelman SF, Anderson BM, Arthur DW, et al. Partial Breast Irradiation for Patients with Early-Stage Invasive Breast Cancer or Ductal Carcinoma In Situ: An ASTRO Clinical Practice Guideline. Pract Radiat Oncol. 2024; 14(2): 112-132. PMID 37977261
 65. American Society of Breast Surgeons. Consensus statements: accelerated partial breast irradiation. Revised 2018. <https://www.breastsurgeons.org/about/statements/>. Accessed May 23, 2025. In supplemental information folder.
 66. Shah C, Vicini F, Shaitelman SF, et al. The American Brachytherapy Society consensus statement for accelerated partial-breast irradiation. Brachytherapy. 2018; 17(1): 154-170. PMID 29074088
 67. Smith BD, Bellon JR, Blitzblau R, et al. Radiation therapy for the whole breast: Executive summary of an American Society for Radiation Oncology (ASTRO) evidence-based guideline. Pract Radiat Oncol. 2018; 8(3): 145-152. PMID 29545124
 68. Silverstein MJ, Kim B, Lin K, Lloyd S, Snyder L, Khan S, Hossino D, Chen P. Intraoperative Radiation Therapy (IORT) for Breast Cancer: The Final Analysis of a Prospective Cohort of 1828 Cases. Ann Surg Oncol. 2025 Aug;32(8):5563-5571. doi: 10.1245/s10434-025-17546-9. Epub 2025 May 31. PMID: 40450167; PMCID: PMC12222354.
 69. Ding L, Mo H, Yang Y, et al. Impact of intraoperative radiotherapy vs. whole-breast external beam radiotherapy for early breast cancer: a propensity score matching study based on SEER database. Gland Surg. Jun 30, 2025; 14(6): 1012-1023. PMID 40671772

The articles reviewed in this research include those obtained in an Internet based literature search for relevant medical references through June 2026, the date the research was completed.

Joint BCBSM/BCN Medical Policy History

Policy Effective Date	BCBSM Signature Date	BCN Signature Date	Comments
6/25/02	6/25/02	6/25/02	Joint policy established
7/20/05	7/20/05	7/28/05	Updated policy; added references to "MammoSite" RT system and accelerated partial breast irradiation.
5/1/07	2/9/07	2/9/07	Routine maintenance; policy retired.
7/1/12	4/10/12	5/18/12	Policy taken out of retirement to address breast brachytherapy done without prior breast conserving surgery (BSC). Brachytherapy of the breast alone (without prior BCS) is experimental and investigational.
9/1/13	6/19/13	6/26/13	Routine maintenance/code update. Policy updated to mirror BCBSA, including title change. Policy statement and criteria for accelerated whole breast radiation changed from "negative surgical margins" to "technically clear surgical margins." Added APBI to the exclusions. Added procedure codes 77261-77263; 77280-77295; 77326-77328 and 0182T.
1/1/14	10/17/13	10/25/13	Removed 0182T from policy as it does not apply to this topic.
9/1/15	6/16/15	7/16/15	Routine maintenance. Routine updated of rationale and references. Deleted codes 77326-8; added codes 77316-8 effective 1/1/15
9/1/16	6/21/16	6/21/16	Routine maintenance. Updated references.
9/1/17	6/20/17	6/20/17	Routine maintenance. References updated. Deleted codes 77776, 77777 and 77787, no longer valid.

5/1/18	2/20/18	2/20/18	Changes made to policy statement to align with ASTRO guidelines. Changes to inclusion/exclusion criteria for alignment. Added code 19294 as established. Added table depicting stages of breast cancer.
5/1/19	2/19/19		Routine policy maintenance. No changes in policy status.
5/1/20	2/18/20		Rationale updated, references # 26, 27, 29, 30, and 38-40 added. Outdated references removed. No change in policy status.
5/1/21	2/16/21		Updated rationale, references # 22, 23, 39 and 48 added. NCCN/ASTRO guidelines section updated.
5/1/22	2/15/22		Updated rationale, references 46, 48, 52 and 54 added. No changes in policy status.
5/1/23	2/21/23		Routine policy maintenance, no changes in policy status. (ds)
5/1/24	4/23/24		Changes made to inclusion/exclusion sections. MPS simplified. Title change. section. Routine policy maintenance. Vendor managed: Evicore (ds)
5/1/25	2/18/25		Routine maintenance, no change in status. Vendor managed: Evicore (ds)
5/1/26	2/17/26		Updated rationale, references added. IORT now E/I, MPS statement changed to E/I. Code 19294 moved to E/I. Title change to "Accelerated Breast Irradiation and Brachytherapy Boost after Breast-Conserving Surgery for EARLY-STAGE Breast Cancer". Formatting changes made. Vendor managed: Evicore (ds, mk)
11/1/26	8/18/26		Added electronic brachytherapy to exclusions. Rationale updated, one reference added. Vendor managed: Evicore (ds, mk)

Next Review Date:

1st Qtr. 2027

BLUE CARE NETWORK BENEFIT COVERAGE

POLICY:

ACCELERATED BREAST IRRADIATION AND BRACHYTHERAPY BOOST AFTER BREAST-CONSERVING SURGERY FOR EARLY-STAGE BREAST CANCER

I. Coverage Determination:

Commercial HMO (includes Self-Funded groups unless otherwise specified)	Covered; criteria apply.
BCNA (Medicare Advantage)	See government section.
BCN65 (Medicare Complementary)	Coinsurance covered if primary Medicare covers the service.

II. Administrative Guidelines:

- N/A