

---

## Medical Policy



Nonprofit corporations and independent licensees  
of the Blue Cross and Blue Shield Association

**Joint Medical Policies are a source for BCBSM and BCN medical policy information only. These documents are not to be used to determine benefits or reimbursement. Please reference the appropriate certificate or contract for benefit information. This policy may be updated and is therefore subject to change.**

---

**\*Current Policy Effective Date: 1/1/25**  
(See policy history boxes for previous effective dates)

### **Title: Transcatheter Arterial Chemoembolization of Hepatic Tumors (TACE)**

---

#### **Description/Background**

##### **TRANSCATHETER ARTERIAL CHEMOEMBOLIZATION**

Transcatheter arterial chemoembolization (TACE) is a minimally invasive procedure performed by interventional radiologists who inject highly concentrated doses of chemotherapeutic agents into the tumor tissues and embolic agent(s) to restrict tumor blood supply. The embolic agent(s) causes ischemia and necrosis of the tumor and slows anticancer drug washout. The most common anticancer drugs used in published TACE studies for hepatocellular carcinoma include doxorubicin (36%), followed by cisplatin (31%), epirubicin (12%), mitoxantrone (8%), and mitomycin C (8%).<sup>1</sup>

The TACE procedure requires hospitalization for placement of a hepatic artery catheter and workup to establish eligibility for chemoembolization. Before the procedure, the patency of the portal vein must be demonstrated to ensure an adequate posttreatment hepatic blood supply. With the patient under local anesthesia and mild sedation, a superselective catheter is inserted via the femoral artery and threaded into the hepatic artery. Angiography is then performed to delineate the hepatic vasculature, followed by injection of the embolic chemotherapy mixture. Embolic material varies but may include a viscous collagen agent, polyvinyl alcohol particles, or ethiodized oil. Typically, only 1 lobe of the liver is treated during a single session, with subsequent embolization procedures scheduled 5 days to 6 weeks later. In addition, because the embolized vessel recanalizes, chemoembolization can be repeated as many times as necessary.

##### **Adverse Events**

Transcatheter arterial chemoembolization of the liver has been associated with potentially life-threatening toxicities and complications, including severe postembolization syndrome, hepatic insufficiency, abscess, or infarction. Transcatheter arterial chemoembolization has been investigated to treat resectable, unresectable, and recurrent hepatocellular carcinoma, cholangiocarcinoma, liver metastases, and in the liver transplant setting. Treatment alternatives include resection when possible, other locally ablative techniques (eg, radiofrequency ablation,

cryoablation), and chemotherapy administered systemically or by hepatic artery infusion. Hepatic artery infusion involves the continuous infusion of chemotherapy with an implanted pump, while TACE is administered episodically. Hepatic artery infusion does not involve the use of embolic material.

---

## **Regulatory Status**

Chemoembolization for hepatic tumors is a medical procedure and, as such, is not subject to regulation by the U.S. Food and Drug Administration. However, the embolizing agents and drugs are subject to Food and Drug Administration approval.

---

## **Medical Policy Statement**

The safety and effectiveness of transcatheter arterial chemoembolization of hepatic tumors has been established and may be considered an established therapeutic option in select conditions.

---

## **Inclusionary and Exclusionary Guidelines**

### **Inclusions:**

- Unresectable hepatocellular cancer confined to liver not associated with portal vein thrombosis and liver function not characterized as Child-Pugh class C.
- Liver metastasis in symptomatic patients with metastatic neuroendocrine tumors whose symptoms persist despite therapy who are not candidates for surgical liver resection
- Liver metastasis in patients with liver-dominant metastatic uveal melanoma
- Bridge to transplant in patients with hepatocellular cancer where the intent is to prevent further tumor growth and to maintain a patient's candidacy for liver transplant
- If criteria for TACE is met, TACE in combination with radiofrequency ablation may be considered a treatment option, when recommended by the oncology specialist

### **Exclusions:**

- Treatment of liver metastases from any other tumors or to treat hepatocellular cancer that does not meet the criteria noted above including recurrent hepatocellular carcinoma
- As a neoadjuvant or adjuvant therapy in hepatocellular cancer that is considered resectable
- Treatment of hepatocellular tumors prior to liver transplantation except as noted above
- Treatment of unresectable cholangiocarcinoma

---

**CPT/HCPCS Level II Codes** *(Note: The inclusion of a code in this list is not a guarantee of coverage. Please refer to the medical policy statement to determine the status of a given procedure.)*

**Established codes:**

37243                      75894

**Other codes (investigational, not medically necessary, etc.):**

N/A

---

**Rationale**

This evidence review was informed by a TEC Assessment (2000) that assessed use of transcatheter arterial chemoembolization (TACE) for hepatic tumors.<sup>2</sup>

Evidence reviews assess the clinical evidence to determine whether the use of a technology improves the net health outcome. Broadly defined, health outcomes are length of life, quality of life, and ability to function-including benefits and harms. Every clinical condition has specific outcomes that are important to patients and to managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of a technology, 2 domains are examined: the relevance and the quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

**Transcatheter Arterial Chemoembolization for Unresectable and Resectable Hepatocellular Carcinoma**

In 2020, an estimated 105,765 people in the U.S. lived with hepatocellular carcinoma (HCC) or intrahepatic cholangiocarcinoma (ICC).<sup>3</sup> Of the primary intrahepatic cancers, HCC and ICC account for 90% and 10% of cases, respectively. The number of new cases of HCC and ICC are estimated at 9.5 per 100,000 men and women per year. The number of deaths are estimated at 6.6 per 100,000 men and women per year.

**TRANSCATHETER ARTERIAL CHEMOEMBOLIZATION FOR UNRESECTABLE HEPATOCELLULAR CARCINOMA CONFINED TO THE LIVER AND NOT ASSOCIATED WITH PORTAL VEIN THROMBOSIS**

## Clinical Context and Therapy Purpose

The purpose of TACE is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as other locally ablative techniques (eg, radiofrequency ablation [RFA], cryoablation), systemic therapy, and supportive care, in individuals with unresectable HCC confined to the liver and not associated with portal vein thrombosis.

The following PICO was used to select literature to inform this review.

## Populations

The relevant population of interest is individuals with unresectable HCC confined to the liver and not associated with portal vein thrombosis.

## Interventions

The therapy being considered is TACE.

TACE of the liver is a proposed alternative to conventional systemic or intra-arterial chemotherapy and to various nonsurgical ablative techniques, to treat resectable and nonresectable tumors. TACE combines the infusion of chemotherapeutic drugs with particle embolization. Tumor ischemia secondary to the embolization raises the drug concentration compared with infusion alone, extending the retention of the chemotherapeutic agent and decreasing systemic toxicity.

## Comparators

Comparators of interest include other locally ablative techniques (eg, RFA, cryoablation), systemic therapy, and supportive care.

## Outcomes

The general outcomes of interest are overall survival (OS), disease-specific survival, quality of life, treatment-related mortality, and treatment-related morbidity. (Table 1)

**Table 1. Outcomes of Interest for Individuals With Unresectable Hepatocellular Carcinoma Confined to Liver and Not Associated with Portal Vein Thrombosis**

Outcomes	Details
Overall survival	[Timing: ≥ 5 years]
Disease-specific survival	Progression-free survival/complete response Local tumor control Time to secondary therapy [Timing for disease-specific survival: 14 weeks to 2 years]

OS: overall survival.

## Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies
- To assess longer term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought

- Studies with duplicative or overlapping populations were excluded.

## REVIEW OF EVIDENCE

### Systematic Reviews

Systematic reviews have compared TACE with hepatic resection and concluded that hepatic resection is superior to TACE for eligible patients.<sup>4,5</sup> For patients with unresectable HCC, the evidence is less but does include some systematic reviews. Table 2 provides a comparative breakdown of 25 studies included in systematic reviews of TACE versus another intervention for unresectable HCC. These studies were published from 1990 to 2011.

A Cochrane review by Oliveri et al (2011) included 9 trials involving 645 patients treated with TACE or transarterial embolization for unresectable HCC.<sup>6</sup> Six of these trials compared TACE with control treatments. Reviewers concluded that all trials were biased, larger trials should be conducted, and that, despite the fact that TACE has been advocated as standard locoregional treatment, there was no firm evidence to support or refute its use in patients with unresectable HCC.

Xie et al (2012) conducted a meta-analysis of 13 studies on treatment for unresectable HCC using chemoembolization (1233 patients) or microsphere embolization (597 patients, using a glass or resin hepatic artery infusion [HAI]).<sup>7</sup> Microsphere embolization treatment resulted in statistically significant longer OS (hazard ratio [HR], 0.73; 95% confidence interval [CI], 0.60 to 0.88;  $p < .001$ ) and time to progression (HR=0.61; 95% CI, 0.41 to 0.89;  $p = .01$ ) than chemoembolization. However, this meta-analysis included uncontrolled observational studies, which limits interpretation.

**Table 2. Comparison of Trials and Studies Included in the Systematic Reviews**

Study	Xie et al (2012) <sup>7</sup>	Oliveri et al (2011) <sup>6</sup>
Ahmad et al (2005) <sup>8</sup>		

Hao et al (2000)<sup>18.</sup>

patients treated with TACE, who had a median PFS of 32 weeks (range, 16 to 70 weeks) versus 26 weeks (range, 14 to 54 weeks) for patients treated with systemic chemotherapy (p=.03). Median OS did not differ significantly between TACE (38 weeks) and chemotherapy (32 weeks; p=.08), except for patients with a serum albumin greater than 3.3 g/dL (60 weeks versus 36 weeks; p=.003). Treatment-related mortality was 4% in the TACE arm and 0% in the chemotherapy arm.

**Table 3. Summary of Key Randomized Controlled Trial Characteristics**

Study	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Bush et al (2016) <sup>33</sup>	U.S.	1	NR	69 patients with clinical or pathologic diagnosis of HCC using either Milan or San Francisco transplant criteria; race or ethnicity of participants were not described	TACE	Proton beam radiotherapy
Mabed et al (2009) <sup>34</sup>	Egypt	1	2003-2005	100 patients with unresectable HCC; race or ethnicity of participants were not described	TACE	Systemic chemotherapy

HCC: hepatocellular carcinoma; NR: not reports; RCT: randomized controlled trial; TACE: transcatheter arterial chemoembolization.

**Table 4. Summary of Key Randomized Controlled Trial Results**

Study	PFS	Overall Survival (%)	Response Rate, n %	TRM, %
Bush et al (2016) <sup>33</sup>	<i>PFS at 2 years</i>		<i>Pathologic complete response after liver transplant</i>	
TACE	31	30 (59) mo (entire group)	1/10 (10)	
Proton beam therapy	48	30 (59) mo (entire group)	3/12 (25)	
95% CI	NR	20.7 to 39.3 mo		
p	.06	NR	.38	
Mabed et al (2009) <sup>34</sup>	<i>Median PFS</i>		<i>Partial response<sup>a</sup></i>	
TACE	32 wks	38 wks	16 (32)	4
Range	16 to 70 wks	22 to 70 weeks		
Systemic chemotherapy	26 wks	32 wks	5 (10)	0
Range	14 to 54 wks	26 to 68 weeks		
p	.03	.08	.007	NR

CI: confidence interval; NR: not reported; PFS: progression-free survival; TACE: transcatheter arterial chemoembolization; TRM: treatment-related mortality.

<sup>a</sup>defined as a decrease of 50% or more in the product of 2 perpendicular diameters of the largest tumour nodule for at least 4 weeks without the appearance of new lesions or progression of lesions

**Table 5. Study Relevance Limitations**

Study	Population <sup>a</sup>	Intervention <sup>b</sup>	Comparator <sup>c</sup>	Outcomes <sup>d</sup>	Follow-Up <sup>e</sup>
-------	-------------------------	---------------------------	-------------------------	-----------------------	------------------------

Bush et al (2016) <sup>33</sup>	3. Patients required to meet Milan or San Francisco criteria for liver transplant to enroll in the trial, and some patients in each group underwent liver transplant after treatment			3. Treatment-related toxicities were only reported in detail for patients who were hospitalized due to complications, and investigators used days of hospitalization as a surrogate to quantify significant toxicity (reported difficulty adjudicating significant events as treatment-related or not treatment-related)	
Mabed et al (2009) <sup>34</sup>	2. Study population is unclear		2. Doxorubicin is not a recommended systemic therapy option in current treatment guidelines; appropriateness of dosing regimen used in the trial is unclear		

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment. a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest.

c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively.

d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not established and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported.

e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

**Table 6. Study Design and Conduct Limitations**

Study	Allocation <sup>a</sup>	Blinding <sup>b</sup>	Selective Reporting <sup>c</sup>	Data Completeness <sup>d</sup>	Power <sup>e</sup>	Statistical <sup>f</sup>
Bush et al (2016) <sup>33</sup>		1,2. No blinding was reported			3. Power estimates led investigators to plan enrollment of 110 patients per treatment arm to identify differences of 15% or greater in 2-year progression-free survival; only 69 patients total were included in this interim analysis	
Mabed et al (2009) <sup>34</sup>		1,2. No blinding was reported				

The evidence limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

PFS: progression-free survival.

a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

b Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

### **Nonrandomized Observational Studies**

Shen et al (2019) published a retrospective, single-center study comparing stereotactic body radiation therapy (SBRT) and TACE as treatments for unresectable HCC of 3 to 8 cm.<sup>35</sup> One hundred eighty-eight patients received either TACE (n=142) or SBRT (n=46) between 2008 and 2017. Before propensity score matching, the 3-year in-field control rates were 63.0% and 73.3% for TACE and SBRT, respectively, while 3-year OS rates were 47.4% and 22.9%. After propensity score matching, 3-year in-field control rates were 55.6% and 77.5% (p=.007), and 3-year OS rates were 13.0% and 55.0% (p<.001), both favoring SBRT. This study was limited by its retrospective nature, long look-back period, and possibility for treatment selection bias.

Biederman et al (2018) published a retrospective, single-center study comparing radiation segmentectomy and TACE as treatments for unresectable, solitary HCC of 3 cm or less.<sup>36</sup> One hundred twelve patients, of whom 57 received TACE, were treated between 2012 and 2016. Results were reported both before and after conducting propensity score matching using the nearest neighbor algorithm (1:1). Before propensity score matching, the complete response rate was 49.1% for TACE and 81.2% for radiation segmentectomy (odds ratio [OR], 2.2; 95% CI, 1.4 to 3.3; p<.001). Median time to secondary therapy was 246 days for TACE and 700 days for radiation segmentectomy (HR, 0.71; 95% CI, 0.55 to 0.92; p=0.009); there was no significant difference in OS (p=.29). After matching, radiation segmentectomy still had significantly better results for complete response (p=0.005) and time to secondary therapy (p=.001), and there was again no significant difference in OS (p=.71). The study was limited by its retrospective nature and the possibility of treatment selection bias.

Multiple noncomparative prospective single-center cohort studies, which included patients with unresectable HCC not suitable for curative treatment with Child-Pugh class A cirrhosis, have reported a favorable impact of TACE on objective response rate or 1-, 3-, and 5-year OS rates.<sup>37,38,39</sup> The largest of these studies published from Japan reported results from an 8-year prospective cohort.<sup>38</sup> In this study, 8510 patients with unresectable HCC underwent TACE using an emulsion of lipiodol and anticancer agents followed by gelatin sponge particles as an initial treatment. The mean follow-up was 1.77 years. Median and 1-, 3-, and 5-year OS rates with TACE were 34 months, 82%, 47%, and 26%, respectively.

### **Section Summary: Transcatheter Arterial Chemoembolization for Unresectable Hepatocellular Carcinoma Confined to the Liver and Not Associated with Portal Vein Thrombosis**

There is evidence from 1 RCT that survival with TACE is at least as good as with systemic chemotherapy.

### **TRANSCATHETER ARTERIAL CHEMOEMBOLIZATION FOR RESECTABLE HEPATOCELLULAR CARCINOMA AS NEOADJUVANT OR ADJUVANT THERAPY**

Although hepatic resection is potentially curative, local recurrence rates after surgery are still high and those rates have led to the use of neoadjuvant and adjuvant systemic therapy approaches to improve outcomes.

### **Clinical Context and Therapy Purpose**

The purpose of neoadjuvant or adjuvant TACE is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as other locally ablative techniques (eg, RFA, cryoablation) and systemic therapy, in patients with resectable HCC.

The following PICO was used to select literature to inform this review.

### **Populations**

The relevant population of interest is individuals with resectable HCC.

### **Interventions**

The therapy being considered is neoadjuvant or adjuvant TACE.

### **Comparators**

Comparators of interest include surgery alone, other locally ablative techniques (eg, RFA, cryoablation) and systemic therapy.

### **Outcomes**

The general outcomes of interest are OS, disease-specific survival, quality of life, treatment-related mortality, and treatment-related morbidity (Table 7).

**Table 7. Outcomes of Interest for Individuals with Resectable Hepatocellular Carcinoma Treated with Neoadjuvant or Adjuvant Transcatheter Arterial Chemoembolization**

Outcomes	Details
Overall survival (OS)	[Timing: Up to 5 years]
Disease-specific survival	Intra- and extrahepatic recurrence [Timing: Up to 5 years] Recurrence-free survival (RFS) [Timing: Up to 5 years]

OS: overall survival; RFS: recurrence-free survival;

### **Study Selection Criteria**

See information under first indication.

# NEOADJUVANT THERAPY

## REVIEW OF EVIDENCE

### Systematic Reviews

Chan et al (2023) performed a meta-analysis of 12 studies (N=3960) that compared preoperative neoadjuvant TACE with resection in patients with large HCC ( $\geq 5$  cm).<sup>40</sup> All but 2 of the included studies were retrospective. There was no significant difference in OS between the 2 treatments. Disease-free survival was more common with TACE than resection alone (HR, 0.79; 95% CI, 0.63 to 0.99;  $p=.04$ ), but operating time and blood loss were also significantly higher with TACE.

Si et al (2016) reported results of a meta-analysis of RCTs that evaluated the impact of neoadjuvant TACE compared to surgery alone.<sup>41</sup> Individually, 2 of the 5 RCTs concluded no effect (no reduction in postoperative recurrence or effect on survival) while 3 suggested unfavorable effect (higher dropouts from definitive surgery, higher prevalence of intraoperative lesions, delayed definitive surgery). None of the studies were graded as low risk of bias in any of the 5 domains of the Cochrane risk of bias tool. Meta-analysis reported no difference between the 2 groups on OS (HR=1.25; 95% CI, 0.92 to 1.68), disease-free survival (DFS) rate (HR=0.95; 95% CI, 0.76 to 1.19), and perioperative mortality rate (OR=0.70; 95% CI, 0.22 to 2.30).

Zhou et al (2013) conducted a meta-analysis of 21 studies evaluating preoperative TACE.<sup>42</sup> Included were 4 RCTs and 17 nonrandomized studies (N=3210). Preoperative TACE was given to 1431 patients, with the remaining 1779 serving as controls. In 18 studies, 5-year DFS for preoperative TACE ranged from 7.0% to 57.0% and from 8.0% to 48.8% in the controls. In 16 studies, 5-year OS for preoperative TACE ranged from 15.4% to 62.7% and from 19.0% to 62.5% in the controls. In pooled analyses, there were no significant improvements with preoperative TACE versus controls in 5-year DFS (32.1% versus 30.0%,  $p=.17$ ) or OS rates (40.2% versus 45.2%,  $p=.37$ ). Intra- and extrahepatic recurrence also did not differ significantly across pooled analyses (51.2% versus 53.6% and 12.9% versus 10.3%, respectively,;  $p=.19$ ).

Chua et al (2010) conducted a systematic review of neoadjuvant TACE for resectable HCC.<sup>43</sup> The authors evaluated 18 studies, including 3 randomized trials and 15 observational studies, some of which are detailed in the following section. The review comprised 3927 patients, 1293 of whom underwent neoadjuvant TACE. Reviewers' conclusions were that TACE could be used safely and resulted in high rates of pathologic responses but did not appear to improve DFS in the TACE group. No conclusions could be drawn about OS differences between the TACE and non-TACE groups due to the heterogeneity of the results across studies.

Table 8 provides a comparative breakdown of RCTs included in select systematic reviews.

**Table 8. Comparison of Randomized Controlled Trials Included in Systematic Reviews**

Study	Si et al (2016) <sup>41,</sup>	Zhou et al (2013) <sup>42,</sup>	Chua et al (2010) <sup>43,</sup>
-------	--------------------------------	----------------------------------	----------------------------------

Kaibori et al (2012)<sup>44,</sup>

Cui et al (2003)<sup>[46](#)</sup>

Neoadjuvant TACE did not provide benefit compared with curative therapy alone in subgroup analysis.

Choi et al (2007) studied 273 patients who underwent curative resection for HCC, 120 of whom had preoperative TACE.<sup>50</sup> The 1-, 3-, and 5-year DFS rates were 76.0%, 57.7%, and 51.3%, in the TACE group and 70.9%, 53.8%, and 46.8% in the non-TACE group, respectively. The differences between the TACE and non-TACE groups were not statistically significant.

### **Subsection Summary: Transcatheter Arterial Chemoembolization for Resectable Hepatocellular Carcinoma as Neoadjuvant Therapy**

Randomized and nonrandomized trials have evaluated TACE as neoadjuvant therapy to hepatic resection in HCC. The highest quality RCTs did not report differences in the survival rates when TACE was added to hepatic resection. Meta-analyses of these studies also did not report differences in outcomes on pooled analyses.

## **ADJUVANT THERAPY**

### **REVIEW OF EVIDENCE**

#### **Systematic Reviews**

Liang et al (2020) published a systematic review and meta-analysis that included 9 RCTs and 15 nonrandomized controlled trials (N=6977) that evaluated adjuvant TACE in patients undergoing liver resection with HCC.<sup>51</sup> Overall survival was based on 6 RCTs and 15 nonrandomized controlled trials, while DFS was reported in 7 RCTs and 6 nonrandomized trials. Compared with surgery alone, use of adjuvant TACE resulted in prolonged OS (HR, 0.67; 95% CI, 0.60 to 0.76;  $p < .001$ ) and DFS (HR, 0.71; 95% CI, 0.61 to 0.84;  $p < .001$ ). The authors noted that 9 nonrandomized controlled trials were at relatively moderate risk of bias and 6 were at relatively serious risk of bias. Among the RCTs, 4 had unknown risk of bias while 5 had high risk of bias. Key RCTs are discussed in the next section.

Liao et al (2017) reported results of a meta-analysis that included 8 RCTs and 12 retrospective studies with a total of 3191 patients (779 in RCT, 2412 in observational studies).<sup>52</sup> Five of the 8 RCTs reported OS and 7 reported recurrence-free survival (RFS). A discussion of key RCTs is presented in the next section. Results showed that adjuvant TACE was associated with improved OS (HR, 0.70; 95% CI, 0.63 to 0.78;  $p < .001$ ) and RFS (HR, 0.69; 95% CI, 0.63 to 0.76;  $p < .001$ ). Results were also similar between the RCTs and retrospective studies for OS (HR, 0.66 and 0.71, respectively) and RFS (HR, 0.66 and 0.70, respectively). Meta-regression revealed that OS was similar among patients treated with various combinations of chemotherapeutic drugs. Majority of the RCTs were rated as moderate risk of bias due to lack of blinding and allocation concealment.

#### **Randomized Controlled Trials**

Li et al (2006) reported the results of an RCT in which 112 patients with HCC and portal vein tumor thrombosis (PVTT) and no extrahepatic metastasis were randomized to surgery ( $n=37$ ), surgery plus TACE ( $n=35$ ), or surgery plus TACE plus portal vein chemotherapy ( $n=40$ ).<sup>53</sup> Race or ethnicity of participants were not described. Staging of HCC was not reported. Portal vein thrombus extirpation was performed at the time of surgery. Although the trial was randomized, no details for randomization including allocation concealment were provided for this single-center trial. Power calculations were also not reported. The DFS curve differed significantly across the 3 groups, as estimated using the Kaplan-Meier method (both  $p < .05$ ).

Overall survival was not reported. Patients who received surgery plus TACE plus portal vein chemotherapy showed a higher DFS rate than those who received surgery only ( $p < .05$ ). There were no statistical differences between patients who received surgery plus TACE and those who received surgery only or between those who received surgery plus TACE plus portal vein chemotherapy and those who received surgery plus TACE (both  $p > .05$ ). The 1-, 3-, and 5-year DFS rates for surgery only were 50.7%, 17.8%, and 0%, respectively; for surgery plus TACE, rates were 62.3%, 23.7%, and 4.0%, respectively; and in surgery plus TACE and portal vein chemotherapy, rates were 74.4%, 46.1%, and 11.5%, respectively. Tumor size, tumor number, PVTT location, and treatment modalities were independent prognostic factors ( $p < .05$ ). Adverse events were mostly related to the surgery, catheters, and local chemotherapy, and included liver decompensation (15.0%), catheter obstruction (11.6%), and nausea and loss of appetite (22.1%).

In the same year, a nearly identical RCT with a larger sample size ( $N=131$ ) was published by the same group.<sup>25</sup> Similarities between the 2 RCTs were same Chinese hospital, same enrollment time period (1998 to 2001), same trial arms (surgery alone or surgery plus TACE or surgery plus TACE plus portal vein chemotherapy), same outcomes (DFS) and same author group. Correspondence with the authors about the study overlap did not yield a response.

Zhong et al (2009) reported on the results of an RCT in which 118 patients with stage IIIA HCC (multiple tumors  $>5$  cm or tumor involving a major branch of the portal or hepatic vein) were randomized to hepatectomy followed by TACE ( $n=59$ ) or to hepatectomy alone ( $n=59$ ).<sup>54</sup> Race or ethnicity of participants were not described. Three patients were excluded from the final analysis (2 from the adjuvant arm, 1 from the hepatectomy arm). Although the trial was randomized, no details on randomization including allocation concealment were provided in this single-center trial. With a sample size of 56 in each arm, the trial was adequately powered (80%) to detect a 20% difference in 5-year survival. The demographic data were well matched between arms. The incremental median OS advantage for adjuvant TACE treatment was 9 months compared with surgery alone (23.0 months vs 14.0 months, respectively,  $p=.048$ ). Confidence intervals around median estimates and HR for death were not reported.

Peng et al (2009) reported the results of a RCT assessing 126 patients with HCC and PVTT who were randomized to liver resection plus PVTT removal ( $n=63$ ) or liver resection plus adjuvant TACE ( $n=63$ ).<sup>55</sup> Race or ethnicity of participants were not described. Staging of HCC was not reported. Twelve patients in the TACE group and 10 patients in the control group were lost during follow-up, and the final analysis included 104 patients. Although the trial was randomized, no details for randomization including allocation concealment were provided in this single-center trial. Power calculations were also not reported. The median OS for TACE adjuvant arm was 13 months (95% CI, 6.3 to 19.8 months) compared with 9 months (95% CI, 6.9 to 11.1 months) in the control arm ( $p < .05$ ). The HR for death was not reported. In addition, 80% of patients had tumor recurrence in the liver, with no significant differences between groups.

### **Subsection Summary: Transcatheter Arterial Chemoembolization for Resectable Hepatocellular Carcinoma as Adjuvant Therapy**

Multiple RCTs and retrospective observational studies, as well as meta-analyses have evaluated TACE as adjuvant therapy to hepatic resection in HCC. Results of the meta-analysis, which included RCTs and retrospective studies, showed that adjuvant TACE was associated with a 30% to 33% relative reduction in the hazard of death and a 29% and 31% relative reduction in the hazard of DFS and recurrence, respectively. However, the meta-

analysis counted the nearly identical RCT's published by Li et al (2006) as separate RCTs. Absent any conclusive evidence that these 2 RCTs are distinct trials, the survival estimates of the meta-analysis are likely over-estimated due to double counting. Further, the entire body of RCTs is comprised of single-center trials from China published in open access journals with inadequate reporting of study procedures (eg, randomization, allocation concealment), patient characteristics (stage of HCC), results (lack of HRs or CIs, inadequate description of the impact of interventions subsequent to recurrence on study endpoints). Well-conducted multi-centric trials from U.S. or Europe, with adequate randomization procedures, blinded assessments, centralized oversight and published in peer-reviewed journals, are required.

**COMBINATION TREATMENT OF LOCOREGIONAL RESECTABLE AND UNRESECTABLE HEPATOCELLULAR CARCINOMA**

**Transcatheter Arterial Chemoembolization Plus Radiofrequency Ablation for Resectable Hepatocellular**

**Clinical Context and Therapy Purpose**

The purpose of TACE plus RFA is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as surgery alone, in patients with resectable hepatocellular cancer.

The following PICO was used to select literature to inform this review.

***Populations***

The relevant population of interest is individuals with resectable HCC.

***Interventions***

The therapy being considered is TACE plus RFA.

***Comparators***

Comparators of interest include surgery alone.

***Outcomes***

The general outcomes of interest are OS, disease-specific survival, quality of life, treatment-related mortality, and treatment-related morbidity (Table 9).

**Table 9. Outcomes of Interest for Individuals with Resectable Hepatocellular Carcinoma Treated with TACE Plus Radiofrequency Ablation**

Outcomes	Details
Overall survival	[Timing: Up to 5 years]
Disease-specific survival	Recurrence-free survival [Timing: Up to 5 years]

OS: overall survival; RFS: recurrence-free survival;

**Study Selection Criteria**

See information under first indication.

**REVIEW OF EVIDENCE**

## Systematic Reviews

Gui et al (2020) published a meta-analysis of data from 1 RCT and 8 retrospective studies to compare TACE plus RFA to surgery alone.<sup>56</sup> Key studies from this meta-analysis, including the single RCT, are summarized below. A total of 867 patients were treated with TACE plus RFA and 1025 patients were treated with surgery. Rates of 1-, 3-, and 5-year OS were not significantly different between treatments. At 1 year, DFS was not significantly different between treatments, and surgery alone demonstrated better DFS at 3 years (OR, 0.78; 95% CI, 0.62 to 0.98;  $p=.03$ ) and 5 years (OR, 0.74; 95% CI, 0.58 to 0.95;  $p=.02$ ). However, in a subgroup analysis of propensity score-matched studies, 3- and 5-year DFS were not significantly different between treatments. This difference in findings may be due to selection bias in the non-matched studies.

## Randomized Controlled Trials

Liu et al (2016) published a RCT in which 200 patients with solitary HCC nodule of 5 cm or less, or up to 3 nodules of 3 cm or less in size (Milan criteria) deemed treatable by partial hepatectomy or TACE plus RFA and liver function characterized as Child-Pugh grade A or B were randomized to surgical resection or to TACE plus RFA.<sup>57</sup> Race or ethnicity of participants were not described. Tumor sizes ranged from 0.6 to 5.0 cm, with a median of 3 cm in the surgical resection group and 2.8 cm in the TACE plus RFA group. Overall survival ( $p=.007$ ) and RFS ( $p=.026$ ) were significantly higher in the surgical resection group (Table 10). Local tumor progression occurred in 1 patient in the surgical resection group and in 18 patients in the TACE plus RFA group ( $p<.001$ ). There were no significant differences in recurrence or OS between the 2 groups for HCC lesions 3 cm or smaller, but there were significant benefits for surgery in recurrence ( $p=.032$ ) and OS ( $p=.012$ ) in patients with lesions larger than 3 cm. Tumor size was an independent prognostic factor for RFS (HR, 1.76;  $p=.006$ ) along with hepatitis B virus DNA and platelet count. Hepatitis B virus DNA was a significant risk factor for length of OS. Complications were higher in the surgical resection group (23.0%) than in the TACE plus RFA group (11.0%;  $p=.24$ ). It was unclear in this trial whether TACE plus RFA was as effective as surgical resection for these small tumors.

**Table 10. Survival Rates After Surgical Resection or Transcatheter Arterial Chemoembolization Plus Radiofrequency Ablation for Resectable Hepatocellular Carcinoma**

Outcomes	1 Year, %	3 Years, %	5 Years, %
Overall survival			
Surgical resection group	97.0	83.7	61.9
TACE plus RFA group	96.0	67.2	45.7
Recurrence-free survival			
Surgical resection group	94.0	68.2	48.4
TACE plus RFA group	83.0	44.9	35.5

Adapted from Liu et al (2016).<sup>57</sup>

OS: overall survival; RFA: radiofrequency ablation; RFS: recurrence-free survival; TACE: transcatheter arterial chemoembolization.

## Retrospective Studies

Ako et al (2018) published a retrospective analysis of 100 patients with HCC who received TACE followed by RFA 20 or more days later.<sup>58</sup> All patients were treated at a single center in Japan between 2001 and 2014. Tumor size reduction was observed in 69% of patients (median reduction rate, 16.2%). Tumor size was unchanged in 3% of patients or increased by 28%. In a univariate analysis, the tumor size at first treatment and the time between therapies were both significantly related to tumor reduction ( $p<.01$  and  $p=.02$ , respectively). The study was limited by its retrospective nature, relatively small population size, potential patient selection bias, and 2 different modalities used to measure tumors, possibly influencing size perception.

Haochen et al (2018) published a retrospective single-center study of 3.1- to 5 cm HCC nodules treated at a university hospital in China, with TACE followed by imaging-guided RFA 2 to 4 weeks later.<sup>59</sup> Two hundred sixteen nodules (162 patients) treated between 2008 and 2016 were identified. Follow-up was performed at 1, 3, 6, and 12 months after TACE plus RFA. Two hundred seven (95.8%) nodules were completely eliminated after 1 to 3 sessions of TACE plus RFA, and 180 (83.3%) nodules were completely eliminated after 1 session. Besides its retrospective nature, no study limitations were reported.

Bholee et al (2017) published a retrospective matched case-control study comparing TACE plus radiofrequency ablation (TACE plus RFA) and hepatectomy as treatments for HCC within Milan criteria.<sup>60</sup> A total of 222 patients were included; 74 individuals treated with TACE plus RFA between 2006 and 2010 at a university cancer center in China, were matched with 148 controls (ratio 1:2) treated with hepatectomy. The 1-, 3-, and 5-year OS for TACE plus RFA was 94.6%, 75.1%, and 55.3%, respectively, and 91.2%, 64.4% and 47.7%, respectively, for hepatectomy ( $p=.488$ ). The 1-, 3-, and 5-year DFS for TACE plus RFA was 87.8%, 48.3%, and 33.5%, respectively, and 68.9%, 49.2%, 40.9%, respectively, for hepatectomy ( $p=.619$ ). The study was limited by possible selection bias due to its nonrandomized design, relatively small population size, and the fact that some patients who received TACE plus RFA did not have histological diagnoses.

### **Section Summary: Transcatheter Arterial Chemoembolization Plus Radiofrequency Ablation for Resectable Hepatocellular Carcinoma**

One RCT has evaluated combination of TACE plus RFA as primary treatment for resectable HCC. The trial failed to show superiority in survival benefit with combination treatment over surgery for HCC lesions 3 cm or smaller. Further, the ad hoc subgroup analysis showed a significant benefit for surgery in recurrence and OS in patients with lesions larger than 3 cm. It cannot be determined from this trial whether TACE plus RFA is as effective as surgical resection for these small tumors. Several retrospective studies have compared TACE with surgical resection; results were inconsistent for which treatment produces better outcomes. A meta-analysis of data from retrospective studies and the sole available RCT did not find significant survival benefits with TACE plus RFA compared to surgery alone. Although TACE plus RFA does not have significant benefit over surgery, there may be clinical situations in which it may be the treatment of choice.

## **TRANSCATHETER ARTERIAL CHEMOEMBOLIZATION PLUS RADIOFREQUENCY ABLATION FOR UNRESECTABLE HEPATOCELLULAR CARCINOMA**

### **Clinical Context and Therapy Purpose**

The purpose of TACE plus RFA is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as RFA alone, in individuals with unresectable HCC.

The following PICO was used to select literature to inform this review.

**Populations**

The relevant population of interest is individuals with unresectable HCC.

**Interventions**

The therapy being considered is TACE plus RFA.

**Comparators**

Comparators of interest include RFA alone.

**Outcomes**

The general outcomes of interest are OS, disease-specific survival, quality of life, treatment-related mortality, and treatment-related morbidity (Table 11).

**Table 11. Outcomes of Interest for Individuals With Unresectable Hepatocellular Carcinoma Treated with Transcatheter Arterial Chemoembolization Plus Radiofrequency Ablation**

Outcomes	Details
Overall survival	[Timing: Up to 5 years]
Disease-specific survival	Local tumor progression [Timing: Up to 3 years]

OS: overall survival

**Study Selection Criteria**

See information under first indication.

**REVIEW OF EVIDENCE**

**Systematic Reviews**

Multiple meta-analyses have recently compared the impact of TACE plus RFA with either treatment alone on progression, RFS, and OS with up to 5 years of follow-up.<sup>61,62,63,64</sup> While many of these meta-analyses have used standard methodologies to pool estimates, including indirect network analysis as well as an assessment of study quality and publication bias, the fundamental flaws in the pooled RCTs render the results of meta-analysis uncertain. For example, Lan et al (2016) reported a network meta-analysis of a combined treatment approach using RFA and TACE but pooled survival estimates from studies that, while individually homogeneous, were collectively heterogeneous in terms of patient populations.<sup>61</sup> In addition, Peng et al (2012)<sup>65</sup> reported on the results of a RCT that enrolled patients with previously treated recurrent HCC tumor 5 cm or smaller while Morimoto et al (2010)<sup>66</sup> enrolled treatment-naïve patients with a solitary tumor measuring 3.1 to 5 cm and Shibata et al (2009)<sup>67</sup> enrolled patients tumor smaller than 3 cm without specifying if patients were treatment-naïve or -experienced. Two of the 5 meta-analyses also included results from the first RCT that demonstrated combination treatment was better than RFA alone.<sup>68</sup> However, that article was retracted in 2009 because of questions about data integrity and reporting.<sup>69</sup>

**Randomized Controlled Trials**

To assess the nature of the evidence that makes the case for combined use of TACE and RFA in HCC, BCBSA reviewed the current RCTs<sup>65,69,70,71</sup> published after 2009 (an arbitrary threshold). All trials were conducted in China and all but 1 were reported in open access journals.<sup>71</sup> In many of these trials where survival was assessed, trialists reported the results of log-rank testing only which would indicate whether there were differences between the survival times of the 2 groups but would not allow other explanatory variables to be taken into account.<sup>65,66,67</sup> No explanations were provided for not reporting results of a semiparametric (Cox) or parametric (exponential, Weibull) model testing for survival analysis.

### **Locoregional Treatment-Naive Therapy for Tumors Less Than 7 cm**

Yi et al (2014) reported the results of a RCT assessing 94 HCC patients with no previous treatment for HCC except liver resection and a solitary tumor measuring 7 cm or smaller or multiple lesions each measuring less than 3 cm.<sup>70</sup> Patients were randomized to sequential TACE plus RFA or microwave ablation (MWA; n=47) or RFA or MWA alone (n=47). The hazard of death was statistically significantly lower in the combined arm versus RFA or MWA alone arm (HR=0.53; 95% CI, 0.33 to 0.82; p=.002). The 5-year OS rate was 62% in the combined arm and 45% in the RFA or MWA alone arm. No subgroup analysis stratified by lesion size was reported.

Peng et al (2013) reported on the results of an adequately powered trial evaluating 189 HCC patients with no previous treatment and a solitary tumor measuring 7 cm or less or fewer than 3 lesions each measuring less than 3 cm.<sup>71</sup> Patients were randomized to receive sequential TACE plus RFA (n=94) or to RFA alone (n=95). Overall survival and RFS were longer in the TACE plus RFA group (HR, 0.56; 95% CI, 0.34 to 0.82; p=.002) than in the RFA group alone (HR, 0.58; 95% CI, 0.37 to 0.90; p=.009). Corresponding OS rates in the 2 groups were 92.6% and 85.3% at 1 year, 66.6% and 61.8% at 2 years, and 59.0% and 45.0% at 4 years, respectively. The major limitation of this well-conducted trial was the generalizability of findings. Over 50% of patients enrolled in the trial had a single lesion with tumor size less than 3 cm (median size, 3.43 cm) even though patients with multiple lesions and tumor measuring up to 7 cm were allowed to enroll. Further, this single center trial conducted in China might not generalize to patients in Western countries.

Morimoto et al (2010) reported the results of a smaller RCT in which 37 HCC treatment-naive patients with a solitary tumor measuring 3.1 to 5 cm were randomized to sequential TACE plus RFA (n=19) or to RFA alone (n=18).<sup>66</sup> While the rates of local tumor progression at end of the third year were significantly lower in the combined arm (6%) than in the RFA alone arm (39%, p=.012), there was no difference in the 3-year survival rates (93% versus 80%, respectively, p=.369). In addition to having same statistical limitations as Peng et al (2012),<sup>64</sup> the Morimoto trial had a small sample size with inadequate power to detect a difference in survival.<sup>65</sup>

### **Locoregional Treatment-Experienced Therapy for Tumors Less Than 5 cm**

Peng et al (2012) also reported on 139 patients with recurrent HCC (after curative treatment with RFA or hepatectomy but not liver transplantation) and tumors measuring up to 5 cm in diameter who were randomized to sequential TACE plus RFA (n=69) or to RFA alone (n=70).<sup>65</sup> A p value less than .008 was considered statistically significant due to multiple comparisons. There were no statistically significant differences in the OS in the combined arm (94%, 69%, and 46%) versus RFA alone arm (82%, 47%, and 36%; p=.037) at 1, 2, and 5 years, respectively. The RFS rates were statistically significantly greater in the combined arm compared with RFA alone arm (80%, 45%, and 40% versus 64%, 18%, and 18% respectively; p=.005). Hazard ratios and confidence intervals were not reported. Further, subgroup analyses

showed that OS was longer for combined arm versus the RFA alone arm among patients with tumors measuring 3.1 to 5.0 cm ( $p=.002$ ) but not for tumors 3 cm or smaller ( $p=.478$ ).

### **Section Summary: Transcatheter Arterial Chemoembolization Plus Radiofrequency Ablation for Unresectable Hepatocellular Carcinoma**

Multiple meta-analyses and RCTs have shown a consistent benefit in survival and RFS favoring combination treatment with TACE plus RFA versus RFA alone. Results of these meta-analyses are difficult to interpret because the pooled data included heterogeneous patient populations and, in a few cases, included data from a study that was retracted due to reporting veracity. Since 2009, several smaller studies, most of which are from China, have reported outcomes favoring the combination treatment of TACE plus RFA. In 2013, a larger well-conducted RCT showed the relative reduction in the hazard of death by 44% and a 14% difference in favor of combination therapy in a proportion of patients surviving at 4 years.

### **Transcatheter Arterial Chemoembolization as a Bridge to Liver Transplant**

Transcatheter arterial chemoembolization has been explored in various settings as a technique to prevent tumor progression in patients on the liver transplant waiting list, to downstage tumors so a patient may be considered a better candidate for liver transplantation, and to decrease the incidence of posttransplant recurrence in patients with larger (T3) tumors. All uses are in part related to the United Network for Organ Sharing (UNOS) liver allocation policy, which prioritizes patients for receiving donor livers. The UNOS policy and the 3 treatment settings are discussed further here.

#### **United Network for Organ Sharing Liver Allocation System**

In 2002, UNOS introduced the Model for End-Stage Liver Disease (MELD) system for allocating new livers to adults awaiting a transplant.<sup>72,73</sup> The MELD score is a continuous disease severity scale incorporating bilirubin, prothrombin time (ie, international normalized ratio), and creatinine into an equation, producing a number that ranges from 6 (less ill) to 40 (gravely ill). Aside from those in fulminant liver failure, donor livers are prioritized to those with the highest MELD score. This system accurately predicts the risk of dying from liver disease except for those with HCC, who often have low MELD scores because bilirubin, international normalized ratio, and creatinine levels are near normal. Therefore, patients with HCC are assigned additional allocation points according to the size and number (T stage) of tumor nodules as follows:

T1: 1 nodule greater than 1 cm and 1.9 cm or smaller

T2: 1 nodule between 2 and 5 cm, or 2 or 3 nodules each 1 cm or greater and up to 3 cm

T3: 1 nodule larger than 5 cm, or 2 or 3 nodules with at least 1 larger than 3 cm.

Patients with T1 lesions are considered at low risk of death on the waiting list, while those with T3 lesions are at high risk of posttransplant recurrence and are generally not considered transplant candidates. Patients with T2 tumors have an increased risk of dying while on the waiting list compared with those who had T1 lesions and are an acceptable risk of posttransplant tumor recurrence. Therefore, UNOS criteria, which were updated in 2022, prioritize only T2 HCC patients who meet specified staging, laboratory and imaging criteria by awarding exception scores in place of the calculated MELD score.<sup>73</sup> This definition of T2 lesions is often referred to as the Milan criteria, in reference to a key study by Mazzaferro et al (1996) that examined the recurrence rate of HCC according to the size of the initial tumor.<sup>74</sup> Liver transplantation for those with T3 HCC is not prohibited, but these patients do not receive

priority on the waiting list. All patients with HCC awaiting transplantation are reassessed at 3-month intervals. Those whose tumors have progressed and are no longer T2 tumors lose the additional allocation points.

Additionally, nodules identified through imaging of cirrhotic livers are given an Organ Procurement and Transplantation Network class 5 designation. Class 5B and 5T nodules are eligible for automatic priority. Class 5B criteria consist of a single nodule 2 cm or larger and up to 5 cm (T2 stage) that meets specified imaging criteria. Class 5T nodules have undergone subsequent locoregional treatment after being automatically approved on initial application or extension. A single class 5A nodule (>1 cm and <2 cm) corresponds to T1 HCC and does not qualify for automatic priority. However, combinations of class 5A nodules are eligible for automatic priority if they meet stage T2 criteria. Nodules less than 1 cm are considered indeterminate and are not considered for additional priority.

The UNOS allocation system provides strong incentives to use locoregional therapies to downsize tumors to T2 status and to prevent progression while on the waiting list. In a report from a national conference in the U.S., Pomfret et al (2010) addressed the need to characterize better the long-term outcomes of liver transplantation for patients with HCC and to assess the justification for continuing the policy of assigning increased priority for candidates with early-stage HCC on the U.S. transplant waiting list.<sup>75</sup> There was a general consensus for developing a calculated continuous HCC priority score for ranking HCC candidates on the list that would incorporate the calculated MELD score,  $\alpha$ -fetoprotein, tumor size, and rate of tumor growth and that only candidates with at least stage T2 tumors would receive additional HCC priority points. The report addressed the role of locoregional therapy to downstage patients from T3 to T2 and stated that the results of downstaging before liver transplantation are heterogeneous, with no upper limits for tumor size and number before downstaging across studies, and the use of different end points for downstaging before transplantation. The UNOS criteria specify that certain patients may undergo downstaging with locoregional therapy in order to qualify for a MELD exception score. Downstaging is possible in patients with 1 lesion between 5 and 8 cm; patients with 2 or 3 lesions with at least 1 lesion greater than 3 cm, no lesion greater than 5 cm, and a total diameter of all lesions of 8 cm or less; and patients with 4 or 5 lesions that are less than 3 cm each and less than or equal to 8 cm total. Patients must meet T2 criteria after downstaging in order to qualify for an exception score. Patients with T2 lesions and elevated  $\alpha$  fetoprotein (>1000 ng/mL) may also undergo locoregional therapy in order to qualify for a MELD exception score ( $\alpha$  fetoprotein must be below 500 ng/mL after treatment in order to qualify for an exception score).

### **Clinical Context and Therapy Purpose**

The purpose of pretransplant TACE is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as other locally ablative techniques (eg, RFA, cryoablation) and systemic therapy, in patients with 1 to 3 small HCC tumors seeking to prevent tumor growth and maintain candidacy for liver transplant.

The following PICO was used to select literature to inform this review.

### ***Populations***

The relevant population of interest is individuals with 1 to 3 small HCC tumors seeking to prevent tumor growth and maintain candidacy for a liver transplant.

### ***Interventions***

The therapy being considered is pretransplant TACE.

**Comparators**

Comparators of interest include other locally ablative techniques (eg, radiofrequency ablation, cryoablation), and systemic therapy.

**Outcomes**

The general outcomes of interest are OS, disease-specific survival, quality of life, treatment-related mortality, and treatment-related morbidity.

**Table 12. Outcomes of Interest for Individuals Awaiting Liver Transplant Who Are Treated with Transcatheter Arterial Chemoembolization**

Outcomes	Details
Overall survival	[Timing: Up to > 7 years]
Disease-specific survival	Tumor recurrence [Timing: Up to 5 years]

OS: overall survival

**Study Selection Criteria**

See information under first indication.

**REVIEW OF EVIDENCE**

**Systematic Reviews**

Butcher et al (2022) reported on a meta-analysis evaluating long-term survival and postoperative complications of pre-liver transplantation TACE in HCC.<sup>76</sup> Twenty-one high-quality non-randomized controlled trials (N=8242) were included. In all included studies, patients underwent or did not undergo TACE based on clinical recommendations while on the transplant waiting list. Overall, individuals treated with TACE had similar survival and postoperative outcomes as non-TACE patients, however, they had worse prognostic features at baseline. In terms of baseline characteristics, tumor diameter was significantly larger in TACE patients (3.49 cm vs. 3.15 cm; p=.02) compared to control groups and time on the transplant waiting list was significantly longer in TACE patients (4.87 months vs. 3.46 months; p=.05), while MELD scores were significantly higher in non-TACE patients (10.81 vs. 12.35; p=.005). There were no significant differences in 3-year OS, 5-year OS, or 3-year DFS between those who received TACE and those who did not. Based on the worse prognostic features at baseline, administration of TACE to patients with poorer prognosis while awaiting liver transplantation may lead to comparable survival outcomes between those who do not receive TACE but have better prognosis characteristics. Interpretation of results is limited, as all studies pooled were nonrandomized with considerable heterogeneity among outcomes. Additionally, waitlist dropout rates could not be analyzed due to inadequate data.

Si et al (2017) reported on a meta-analysis evaluating the correlation between preoperative TACE and liver transplant.<sup>77</sup> This meta-analysis included 2902 patients (721 had TACE plus liver transplant, 2181 had liver transplant alone) from 7 retrospective cohort studies and 5 case-control studies. It is unclear as to how patients were selected in the control arm (ie, those who did not receive TACE) in the individual studies. Further, it is not clear if the authors of the meta-analysis extracted unadjusted or adjusted estimates from individual studies. Because all studies were observational, it is important to know how the TACE groups differed at baseline

from the control groups, particularly with respect to prognostic factors, and whether statistical controls were used (if any beyond case-control matching) to adjust the hazard ratio estimates in the primary studies. Results of the meta-analysis showed no difference in OS (HR, 1.05; 95% CI, 0.65 to 1.72;  $p=.83$ ), but a higher rate of vascular complications (RR, 2.01; 95% CI, 1.23 to 3.27;  $p=.005$ ) and a reduction in DFS (HR, 1.66; 95% CI, 1.02 to 2.70;  $p=.04$ ) with those receiving TACE compared with those who did not. Reviewers hypothesized that vascular complications resulting from repeated intubations and toxic damage of chemotherapeutic drugs could seriously affect the function of transplanted liver and that early hepatic artery thrombosis after liver transplant might result in graft loss. The meta-analysis also reported regional differences in TACE outcomes between Asia and Western countries potentially related to differences in the mechanisms of hepatocarcinogenesis (alcoholic liver cirrhosis in the western countries versus hepatitis B in the Asian subcontinent). Subgroup analysis of OS showed that hazard of death was higher in 2 Asian studies (HR, 2.65; 95% CI, 1.49 to 4.71) than in 4 European studies (HR, 1.01; 95% CI, 0.74 to 1.37). Similarly, the hazard of death varied by whether the studies were retrospective cohort (HR, 1.66) or case-control studies (HR, 0.84) and whether they were higher (HR, 1.46) or lower quality (HR, 0.70) studies. Given that all studies pooled were nonrandomized with considerable heterogeneity and directional differences in the outcomes based on geography and study designs, interpretation of the results is uncertain.

### **Prospective Studies**

Graziadei et al (2003) reported on 48 patients with HCC awaiting transplantation; all underwent TACE every 6 to 8 weeks until complete response or a donor organ became available.<sup>78</sup> None was removed from the list due to tumor progression after a mean waiting time of 178 days. Of the 48 patients, 41 underwent a liver transplant. The 1-, 2-, and 5-year intention-to-treat survival rates were 98%, 98%, and 94%, respectively. Tumor recurrence was only reported in 1 (2.4%) patient. Maddala et al (2004) reported on dropout rates for 54 patients who received TACE while awaiting transplantation.<sup>79</sup> During a median waiting time of 211 days (range, 28 to 1099 days), the dropout rate was 15%. Obed et al (2007) reported on 20 patients with nonprogressing lesions after TACE who had liver transplantation; median survival in this group was 92.3 months.<sup>80</sup>

### **Transcatheter Arterial Chemoembolization to Downstage Hepatocellular Carcinoma Prior to Transplant or to Reduce Recurrence in Those With T3 Lesions (Bridge to Transplant)**

Published literature reflects an ongoing discussion whether the United Network for Organ Sharing allocation criteria (see Background) should be expanded to include patients with larger tumors. Some patients with T3 lesions are cured with a liver transplant, although most experience tumor recurrence. For example, in the seminal study by Mazzaferro et al (1996),<sup>74</sup> the 4-year RFS rate was 92% in those who met the Milan criteria (T2 lesion) compared with 59% in those who did not; additional studies confirm this difference in RFS rate.

However, other institutions have reported similar outcomes with expanded criteria. Yao at University of California at San Francisco (UCSF) reported similar RFS after transplant in patients with T2 tumors and a subset of those with T3 tumors.<sup>81</sup> This T3 subset was defined as a single lesion 6.5 cm or smaller or no more than 3 lesions with none greater than 3 cm, with a sum of tumor diameters 8 cm or smaller. These expanded criteria are known as “the UCSF criteria.”

Lewandowski et al (2009) compared the efficacy of radioembolization with chemoembolization in downstaging 86 patients with HCC from stage T3 to T2.<sup>23</sup> Patients were treated with yttrium-90 (Y90) microspheres (n=43) or TACE (n=43). Median tumor size was similar between treatment groups (5.7 cm for TACE versus 5.6 cm for radioembolization). Partial response rates were 61% and 37% for radioembolization and TACE, respectively, with downstaging from T3 to T2 in 58% of patients treated with radioembolization and 31% with TACE ( $p<.05$ ).

Gabr et al (2017) published a prospective, single-center comparative study analyzing posttransplant outcomes for patients with HCC bridged or downstaged to orthotopic liver transplantation by TACE or Y90 radioembolization.<sup>82</sup> One hundred seventy-two patients (TACE=79, Y90=93) treated between 2003 and 2013 were identified; a classification into the TACE or Y90 group was based on the first liver-directed therapy received. Median posttransplant follow-up was 26.1 months. For TACE, 6 (8%) of 79 patients experienced tumor recurrence and 8 (9%) of 92 patients who received Y90 experienced tumor recurrence. There were no significant differences in RFS (TACE, 77 months versus Y90, 79 months;  $p=.71$ ) and OS (TACE, 87.2 months versus Y90, median not reached at 100 months;  $p=.42$ ) between groups. The study was limited by its relatively small sample size, inherent selection bias since transplanted patients usually exhibit more favorable biology and response, and lack of etiology of death for some patients.

### **Section Summary: Transcatheter Arterial Chemoembolization as a Bridge to Liver Transplant**

There is a lack of comparative trials assessing TACE as a bridge to liver transplantation. Several small prospective studies have demonstrated that TACE can prevent dropouts from the transplant list. The evidence on vascular complications and long-term survival is conflicting and limited to retrospective case-control and cohort studies. Two meta-analyses of these studies have shown no difference in OS among patients who received TACE as a bridging therapy and those who did not prior to transplant. The older meta-analysis did show a higher rate of vascular complications and a reduction in DFS with TACE, but the more recent meta-analysis did not demonstrate a difference in DFS. The more recent meta-analysis (Butcher et al [2022]) demonstrated no difference between groups despite the TACE group having worse prognostic characteristics at baseline. The significant limitations of the meta-analyses, including lack of clarity on the use of unadjusted or adjusted estimates from individual studies, lack of randomized data, considerable heterogeneity and directional differences based on geography and study designs, limit the interpretation of results. The consequences of dropping from a transplant list is likely death and, therefore, any strategy that delays progression with an acceptable safety profile is beneficial, and available data have demonstrated that for TACE. However, the relative efficacy and safety of various locoregional treatments as a bridge therapy or to downstage HCC have not been evaluated in an RCT setting.

### **Transcatheter Arterial Chemoembolization for Unresectable Intrahepatic Cholangiocarcinoma**

As mentioned earlier, an estimated 100,476 people in the U.S. lived with HCC or ICC in 2019.<sup>3</sup> Surgical resection represents the only form of curative therapy for ICC. However, most ICC patients are not surgical candidates due to their advanced disease at diagnosis, which is caused by the lack of symptoms until late in disease progression. The overall prognosis of ICC is far worse than for extrahepatic cholangiocarcinoma because of its late presentation. Most patients with ICC qualify for palliative therapy, including systemic chemotherapy and

radiotherapy. However, such palliative options afford little to no survival benefit over supportive therapy alone, because ICC responds poorly to such existing therapies.<sup>83</sup> Survival prognosis for patients with unresectable ICC is poor, with a median survival of 3 to 6 months if left untreated.<sup>84</sup>

**Clinical Context and Therapy Purpose**

The purpose of TACE is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as other locally ablative techniques (eg, RFA, cryoablation) and systemic therapy, in patients with unresectable ICC.

The following PICO was used to select literature to inform this review.

**Populations**

The relevant population of interest is individuals with unresectable ICC.

**Interventions**

The therapy being considered is TACE.

**Comparators**

Comparators of interest include other locally ablative techniques (eg, RFA, cryoablation) and systemic therapy.

**Outcomes**

The general outcomes of interest are OS, disease-specific survival, quality of life, treatment-related mortality, and treatment-related morbidity.

**Table 13. Outcomes of Interest for Individuals with Unresectable Intrahepatic Cholangiocarcinoma Treated with Transcatheter Arterial Chemoembolization**

Outcomes	Details
Overall survival	[Timing: > 22 months]
Disease-specific survival	Timing: Up to 5 years]

OS: overall survival

**Study Selection Criteria**

See information under first indication.

**REVIEW OF EVIDENCE**

**Systematic Reviews**

Lv et al (2022) conducted a systematic review of 11 observational studies that compared TACE versus non-TACE interventions in patients with ICC.<sup>85</sup> Some of the included studies had patients who underwent surgical resection. Among patients who received palliative TACE for nonresectable ICC (N=3 studies), OS was significantly higher than in patients who received supportive treatment (HR, 0.35; 95% CI, 0.25 to 0.47; p<.00001). Survival at 1 year was also significantly higher with palliative TACE versus non-TACE interventions (63.9% vs. 9.2%; p<.00001). Interpreting these results is limited by a lack of information about the non-TACE interventions.

Edeline et al (2021) conducted a systematic review of nonrandomized studies of locoregional treatments (ie, TACE, radioembolization, HAI, EBRT, and ablation) for unresectable ICC.<sup>86</sup> There were 22 cohorts (n=1145) for TACE, 27 cohorts (n=1232) for radioembolization, 16 cohorts (n=331) for HAI, 18 cohorts (n=541) for EBRT, and 15 cohorts (n=645) for ablation. The mean weighted OS was 15.9 months (95% CI, 12.9 to 19.0) for TACE, 14.1 months (95% CI, 12.1 to 16.0) for radioembolization, 21.3 months (95% CI, 15.4 to 27.1) for HAI, 18.9 months (95% CI, 14.2 to 23.5) for EBRT, and 30.2 months for ablation (95% CI, 21.8 to 38.6).

Mosconi et al (2021) conducted a systematic review and meta-analysis of 31 studies (N=1695) with either TACE or transarterial radioembolization for unresectable ICC.<sup>87</sup> Median survival after TACE was 14.2 months (95% CI, 11.6 to 17.6) versus 13.5 months (95% CI, 11.4 to 16.1) after transarterial radioembolization. Radiologic objective response was similar between groups (20.6% vs. 19.3%, respectively). Adverse events were more common with TACE (58.5%) than with transarterial radioembolization (43.0%). Substantial heterogeneity between groups limits interpretation of these results.

Boehm et al (2015) conducted a meta-analysis of 20 studies (N=657) on the hepatic artery therapies of TACE, hepatic arterial infusion (HAI) and Y90 for ICC.<sup>88</sup> Median OS was lowest for TACE (12.4 months) and drug-eluting bead TACE (12.3 months) compared with HAI (22.8 months) and Y90 (13.9 months). Complete and partial responses to therapy were also lowest with TACE (17.3%) compared with Y90 (27.4%) and HAI (56.9%). Transcatheter arterial chemoembolization had lower grade 3 and 4 toxicity (0.26 events per patient) than HAI (0.35 events per patient).

### **Nonrandomized Observational Studies**

Knüppel et al (2012) reviewed 195 patients with intrahepatic (57%) or extrahepatic (43%) cholangiocarcinoma. Patients received chemotherapy or a combination of photodynamic therapy or TACE with chemotherapy.<sup>89</sup> Some patients underwent surgical resection. Patients who only received palliative care (no surgery) survived 9.8 months longer with combination chemotherapy and TACE (n=14) than with chemotherapy alone (n=81) (median survival for chemotherapy plus TACE, 22.0 months vs. chemotherapy alone, 12.2 months; p=.039). Survival was not reported for extrahepatic versus ICC.

Park et al (2011) reviewed the medical and imaging records of 155 patients with unresectable ICC treated with TACE between 1996 and 2009.<sup>83</sup> Patients who had undergone local or systemic therapy were excluded. Seventy-two patients underwent TACE and 83 received supportive care, based on physician and patient preference. Survival was the primary end point. Baseline patient and tumor characteristics were well-balanced between groups. Most patients had stage III or IV disease. Tumor multiplicity was single and multiple or diffuse in 43% and 57% of the TACE patients, respectively, and in 53% and 47% of the supportive group, respectively. Maximum tumor size in the TACE group was 8.1 cm and 7.8 cm in the supportive group. The median number of sessions per patient in the TACE group was 2.5 (range, 1 to 17 sessions). After TACE, the incidences of significant ( $\geq$  grade 3) hematologic and nonhematologic toxicities were 13% and 24%, respectively, and no patients died within 30 days of TACE. Across a range of outcomes, TACE outperformed supportive care. For example, Kaplan-Meier survival analysis showed a median survival in the TACE group of 12.2 months versus 3.3 months in the supportive therapy group (p<.001). Survival rates differed significantly between groups according to the presence or absence of extrahepatic metastases. In patients with the liver-only disease, median survival was 13.3 months (95% CI,

9.2 to 17.4 months) for the TACE group and 4 months (95% CI, 3 to 5 months;  $p < .001$ ) for the supportive treatment group. In patients with extrahepatic metastases, median survival was 11.3 months (95% CI, 8.9 to 13.7 months) for the TACE group and 3.2 months for the supportive treatment group (95% CI, 2.6 to 3.8 months;  $p < .001$ ).

### **Section Summary: Transcatheter Arterial Chemoembolization for Unresectable Intrahepatic Cholangiocarcinoma**

RCTs evaluating the benefit of adding TACE to standard of care for patients with unresectable ICC are lacking. Results from retrospective studies have reported survival benefit with TACE over the standard of care; however, systematic reviews comparing TACE to other locoregional therapies are conflicting..

## **TRANSCATHETER ARTERIAL CHEMOEMBOLIZATION FOR SYMPTOMATIC UNRESECTABLE NEUROENDOCRINE TUMORS**

Neuroendocrine tumors are a heterogeneous group of typically slow-growing tumors with an indolent course, with the capacity to synthesize and secrete hormones. Liver metastases may result in significant hormonal symptoms and are associated with a poor prognosis.

### **Clinical Context and Therapy Purpose**

The purpose of TACE is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as other locally ablative techniques (eg, RFA, cryoablation) and systemic therapy, in patients with symptomatic metastatic neuroendocrine tumors despite systemic therapy and who are not candidates for surgical resection.

The following PICO was used to select literature to inform this review.

### ***Populations***

The relevant population of interest is individuals with symptomatic metastatic neuroendocrine tumors despite systemic therapy and who are not candidates for surgical resection.

Systemic chemotherapy for these tumors has shown modest response rates of limited duration, and although somatostatin analogues are usually effective at controlling symptoms, the disease eventually becomes refractory. Therefore, liver-directed therapies aim to reduce tumor burden, to lower hormone levels, and to palliate symptoms in patients with unresectable neuroendocrine metastases.

### ***Interventions***

The therapy being considered is TACE.

### ***Comparators***

Comparators of interest include other locally ablative techniques (eg, RFA, cryoablation) and systemic therapy.

### ***Outcomes***

The general outcomes of interest are OS, disease-specific survival, symptoms, quality of life, treatment-related mortality, and treatment-related morbidity.

### **Table 14. Outcomes of Interest for Individuals with Unresectable Metastatic Neuroendocrine Tumors Treated with Transcatheter Arterial Chemoembolization**

Outcomes	Details
Overall survival	[Timing: Up to 5 years]
Disease-specific survival	Freedom from disease progression [Timing: Up to 3 years]
Quality of life	Symptomatic relief [Timing: Up to 3 years]

OS: overall survival

## Study Selection Criteria

See information under first indication.

## REVIEW OF EVIDENCE

### Systematic Reviews

Tai et al (2020) published a systematic review and meta-analysis comparing TACE to transarterial bland embolization in 8 studies (N=504) in patients with neuroendocrine tumors.<sup>90</sup> Seven of the included studies were retrospective cohort studies, and 1 small RCT was included. No differences between groups were found in OS at 1 year (OR, 0.72; 95% CI, 0.27 to 1.94), 2 years (OR, 0.69; 95% CI, 0.43 to 1.11), or 5 years (OR, 0.91; 95% CI, 0.37 to 2.24). In addition, PFS was not different between groups at 1 year (OR, 0.71; 95% CI, 0.38 to 1.55), 2 years (OR, 0.83; 95% CI, 0.33 to 2.06), or 5 years (OR, 0.91; 95% CI, 0.37 to 2.24). The authors noted that the quality of evidence is limited due to the rarity of neuroendocrine tumors. In addition, other factors (age, sex, performance status, tumor grade, volume of hepatic metastasis) may have influenced OS.

A literature review by Nazario and Gupta (2010) summarized the experience with TACE (and transarterial embolization).<sup>91</sup> They evaluated multiple nonrandomized, retrospective reports that demonstrated reduced tumor burden, lowered hormone levels, and palliation of symptoms with these interventions. Radiologic responses ranging from 25% to 95% and symptomatic responses ranging from 53% to 100% were reported. Five-year OS rates varied from 14% to 75%, likely a reflection of the heterogeneity of the patient populations and treatment regimens used.

### Nonrandomized Observational Studies

Ruutinen et al (2007) reported on a retrospective study of 67 patients who underwent 219 embolization procedures: 23 patients received primarily bland embolization, and 44 primarily received TACE.<sup>92</sup> Patients with disease relapse were retreated when feasible. Ten (15%) of 67 patients were lost to follow-up. Toxicities of grade 3 or 4 occurred after 25% of chemoembolization procedures and 22% of bland embolization procedures. Rates of freedom from disease progression at 1, 2, and 3 years were numerically but not statistically superior for TACE (49%, 49%, and 35%) compared with bland embolization (0%, 0%, and 0%,  $p=.16$ ). Patients treated with chemoembolization also experienced longer symptomatic relief (15 months) than those who received bland embolization (7.5 months,  $p=.14$ ). Posttherapy survival rates at 1, 3, and 5 years were 86%, 67%, and 50% for TACE and 68%, 46%, and 33% for bland embolization 7.5 months;  $p=.18$ ). These results are consistent with those reported by Gupta et al (2003) on a retrospective series of 81 patients given hepatic artery embolization or chemoembolization, which resulted in symptomatic and radiographic responses in most patients with carcinoid metastases to the liver.<sup>93</sup> Osborne et al (2006) reported on a nonrandomized study of 59 patients with neuroendocrine tumors who received cytorreduction

or embolization for symptomatic hepatic metastases.<sup>94</sup> Both duration of symptom relief (35 months versus 22 months) and survival (43 months vs. 24 months) favored the cytoreduction approach.

**Section Summary: Transcatheter Arterial Chemoembolization for Symptomatic Unresectable Neuroendocrine Tumors**

For patients with unresectable neuroendocrine tumors, there is a lack of RCT evidence assessing TACE. Uncontrolled trials have reported that TACE reduces symptoms and tumor burden, and improves hormone profile. Generally, the response rates exceed 50% and includes patients with massive hepatic tumor burden. Despite the uncertain benefit on survival, the use of TACE to palliate the symptoms associated with hepatic neuroendocrine metastases can provide a clinically meaningful improvement in the net health outcome.

**Transcatheter Arterial Chemoembolization for Liver-Dominant Metastatic Uveal Melanoma**

Uveal melanoma (also called ocular melanoma) is the most common primary ocular malignancy in adults and shows a strong predilection for liver metastases.

**Clinical Context and Therapy Purpose**

The purpose of TACE is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as other locally ablative techniques (eg, RFA, cryoablation), in individuals with liver-dominant metastatic uveal melanoma.

The following PICO was used to select literature to inform this review.

***Populations***

The relevant population of interest is individuals with liver-dominant metastatic uveal melanoma.

Even with successful treatment of the primary tumor, up to 50% of patients will subsequently develop systemic metastases, with liver involvement in up to 90% of these patients. Metastatic uveal melanoma is resistant to systemic chemotherapy, leading to the evaluation of locoregional treatment modalities to control tumor progression in the liver, including TACE.

***Interventions***

The therapy being considered is TACE.

***Comparators***

Comparators of interest include other locally ablative techniques (eg, RFA, cryoablation).

***Outcomes***

The general outcomes of interest are OS, disease-specific survival, quality of life, treatment-related mortality, and treatment-related morbidity.

**Table 15. Outcomes of Interest for Individuals with Liver-Dominant Metastatic Uveal Melanoma Treated with Transcatheter Arterial Chemoembolization**

Outcomes	Details
----------	---------

Overall survival	[Timing: Up to > 2 years]
------------------	---------------------------

OS: overall survival

## Study Selection Criteria

See information under first indication.

## REVIEW OF EVIDENCE

### Systematic Reviews

A literature review by Rowcroft et al (2020) summarized published studies on liver-directed therapies in patients with hepatic metastases from uveal melanoma.<sup>95</sup> Median OS with TACE ranged from 5 to 29 months in 17 prospective and retrospective observational studies that included a total of 647 patients.

### Nonrandomized Observational Studies

Huppert et al (2010) reported on a single-arm prospective study of 14 patients with hepatic metastases from uveal melanoma who underwent TACE.<sup>96</sup> Patients received a mean of 2.4 treatments (34 total treatments). Responses were partial for 8 (57%) patients, stable for 4 (29%) patients, and tumor progression for 2 (14%) patients. Median time to progression was 8.5 months (range, 5 to 35 months), and median survival after the first TACE treatment was 14.5 months in responders and 10 months in nonresponders ( $p=.18$ ). Survival rates were 86% at 6 months, 50% at 12 months, 28% at 18 months, and 14% at 24 months after the first TACE treatment. A survival advantage was most pronounced for patients with tumors occupying less than 25% of the liver volume ( $n=7$ ); that subgroup had a median survival of 17 months versus 11 months in the 7 patients with more than 25% involvement of the liver ( $p=.02$ ). The authors stated that, compared with no treatment, survival after detection of liver metastases was 2 to 7 months, with a median 1-year survival rate less than 30%. Response rates for systemic chemotherapy were less than 10%, and 20% to 50% with immunochemotherapy, but with only a median survival of 5 to 9 months and serious toxicity.

Sharma et al (2008) reported the results of a retrospective single cohort study that assessed the use of TACE for melanoma metastatic to the liver in a series of 20 patients (17 with ocular melanoma) treated between 2004 and 2007.<sup>97</sup> The 20 patients underwent 46 TACE sessions (mean, 2.4 sessions; range, 1 to 5 sessions). Mean and median OS times were 334 days and 271 days, respectively. There were no deaths within 30 days of treatment. The authors noted TACE resulted in longer survival than had been noted among historical controls. This work built on results reported by Bedikian et al (1995), which showed that TACE had a 36% response rate (cisplatin chemoembolization) compared with a 1% response rate to systemic chemotherapy.<sup>98</sup>

Patel et al (2005) reported the results of a prospective single cohort study of TACE for treatment of hepatic metastasis from uveal melanoma.<sup>99</sup> In this study, 18 of the 24 patients experienced regression or stabilization of hepatic metastases for at least 6 weeks. Overall response rates (complete responses and partial responses) for the intention-to-treat population and for patients evaluable for response were 16.7% and 20.4%, respectively. The median OS of the entire intention-to-treat group of patients was 5.2 months; for patients with complete responses or partial response in hepatic metastases, it was 21.9 months; for patients with stable disease, 8.7 months; and for patients with disease progression, 3.3 months.

**Section Summary: Transcatheter Arterial Chemoembolization for Liver-Dominant Metastatic Uveal Melanoma**

For patients with liver-dominant metastatic uveal melanoma, there is a lack of RCT evidence for TACE likely due to rarity of this condition. Noncomparative prospective and retrospective case series have reported improvements in tumor response and survival compared to historical controls treated with systemic therapy. Given the very limited treatment response from systemic therapy and the rarity of this condition, the existing evidence may support conclusions that TACE meaningfully improves outcomes for patients with hepatic metastases from uveal melanoma.

**Transcatheter Arterial Chemoembolization for Other Unresectable Hepatic Metastases**

**Clinical Context and Therapy Purpose**

The purpose of TACE is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as other locally ablative techniques (eg, RFA, cryoablation) and systemic therapy, in patients with unresectable hepatic metastases from other types of primary tumors (eg, colorectal, breast).

The following PICO was used to select literature to inform this review.

**Populations**

The relevant population of interest is individuals with unresectable hepatic metastases from other types of primary tumors (eg, colorectal, breast).

**Interventions**

The therapy being considered is TACE.

**Comparators**

Comparators of interest include other locally ablative techniques (eg, RFA, cryoablation) and systemic therapy.

**Outcomes**

The general outcomes of interest are OS, disease-specific survival, quality of life, treatment-related mortality, and treatment-related morbidity (Table 16).

**Table 16. Outcomes of Interest for Individuals with Other Unresectable Hepatic Metastases**

Outcomes	Details
Overall survival	[Timing: Up to 3 years]
Disease-specific survival	Progression-free survival [Up to >15 months] Local tumor control [Up to >15 months]

OS: overall survival; PFS: progression-free survival.

**Study Selection Criteria**

See information under first indication.

**REVIEW OF EVIDENCE**

# Metastatic Colorectal Cancer

## Systematic Reviews

Sugumar et al (2024) conducted a systematic review and meta-analysis of several liver-directed therapies, including TACE, for the treatment of colorectal cancer liver metastases.<sup>100</sup> The authors included 7 articles with TACE plus systemic chemotherapy, and 55 articles with other therapies. At 12 months, OS was 83% with TACE, which decreased to 14% at 36 months. Progression-free survival with TACE was 66% at 6 months, followed by 20% at 12 months, and 3% at 36 months. No statistical comparisons between treatments were reported.

Zacharias et al (2015) published a meta-analysis on hepatic-artery based therapies for colorectal metastases.<sup>101</sup> Techniques included TACE, HAI chemotherapy, and radioembolization. Ninety studies reported on outcomes of HAI-based therapy. Eight studies were RCTs, including 1 RCT of TACE. On combined analysis, OS for patients treated with TACE was 15.2 months, compared to 21.4 months with HAI and 29.4 months with radioembolization. Differences between groups were not statistically significant. The grade 3 or 4 toxicity rates were 40% in the HAI group, 19% in the radioembolization group, and 18% in the TACE group. This review included retrospective studies along with prospective studies and RCTs, so interpretation of these combined analyses may be limited.

Richardson et al (2013) reported on a systematic review (1 RCT, 5 observational studies) of TACE for unresectable colorectal liver metastasis.<sup>102</sup> Median survival times ranged from 15.2 to 25 months. The most common adverse events were postembolization syndrome (abdominal pain, nausea, vomiting) followed by hypertension.

Swierz et al (2020) reported the results of a Cochrane that assessed benefits and harms of TACE compared with no intervention or placebo in patients with liver metastases irrespective of the location of primary tumor.<sup>103</sup> Only 1 RCT published in 1990 fulfilled inclusion criteria. It randomized 61 patients with colorectal liver metastases were randomized into hepatic artery embolization, HAI chemotherapy, and no active therapeutic intervention. Reviewers judged this trial to have a risk of bias on the basis of lack of sequence generation and lack of allocation concealment or blinding. Results of the trial with respect to mortality were inconclusive. Reviewers concluded that in patients with liver metastases, the evidence regarding benefits and harms of TACE versus no active treatment is lacking, and more high-quality RCTs are necessary to draw conclusions about TACE in this setting.

Table 17 provides a comparative breakdown of studies included in the highest quality systematic reviews (eg, reviews that only considered RCTs and/or prospective trials).

**Table 17. Comparison of Trials and Studies Included in Select Systematic Reviews**

Study	Swierz et al (2020) <sup>103</sup> ,	Richardson et al (2013) <sup>102</sup> .
-------	--------------------------------------	--

Hunt et al (1990)<sup>104</sup>



were present in 43% and 33%, respectively. Of 36 patients, 21 underwent TACE, with a median survival of 7.7 months measured from the first TACE treatment. Median survival was 6.9 months in the radioembolization group ( $p=.27$ ). Survival results were comparable with other studies addressing colorectal cancer and TACE (range, 7 to 10 months). The 1-, 2-, and 5-year survival rates were 43%, 10%, and 0%, respectively, for the chemoembolization group and 34%, 18%, and 0%, respectively, for the radioembolization group.

## **Metastatic Breast Cancer**

### **Systematic Review**

Rivera et al (2021) published a systematic review of various liver directed therapies, including TACE, for treatment of breast cancer liver metastases.<sup>115</sup> The systematic review included 8 retrospective and prospective studies (N=362) that evaluated TACE; however, no RCTs were identified. Pooled median OS was 19.6 months (based on 6 studies) and 1-year survival ranging from 32% to 88.8% (based on 4 studies) with use of TACE.

### **Nonrandomized Trial**

Vogl et al (2010) published a study that was not included in the systematic review. The authors reported on the efficacy of repeated TACE treatments in 208 patients with unresectable hepatic metastases from breast cancer.<sup>116</sup> A total of 1068 chemoembolizations were performed (mean, 5.1 sessions per patient; range, 3 to 25). Patients received 1 of the chemotherapeutic agents alone (mitomycin-C or gemcitabine) or in combination. Tumor response was evaluated by magnetic resonance imaging using Response Evaluation Criteria in Solid Tumors (RECIST) criteria. For all chemotherapy protocols, local tumor control was 13% (27/208); stable disease, 50.5% (105/208); and progressive disease, 36.5% (76/208). The 1-, 2-, and 3-year survival rates after TACE were 69%, 40%, and 33%, respectively. Median and mean survival times from the beginning of the TACE sessions were 18.5 months and 30.7 months, respectively. Treatment with mitomycin-C only showed median and mean survival times of 13.3 months and 24 months; and with gemcitabine, 11 months and 22.3 months, respectively. With combination mitomycin-C and gemcitabine, median and mean survival times were 24.8 months and 35.5 months, respectively.

### **Section Summary: Transcatheter Arterial Chemoembolization for Other Unresectable Hepatic Metastases**

For other types of hepatic metastases, the largest amount of evidence assesses colorectal cancer. Multiple RCTs and numerous nonrandomized studies that have compared TACE with alternatives. The nonrandomized studies have indicated that TACE can stabilize 40% to 60% of treated patients but whether this translates into a prolonged survival benefit relative to systemic chemotherapy alone is uncertain. Two small RCTs have reported that TACE results in statistically significant improvements in response rates and PFS. Whether this translates into a prolongation of survival relative to systemic chemotherapy alone is uncertain. For cancers other than colorectal, the evidence is extremely limited and no conclusions can be made.

## **SUMMARY OF EVIDENCE**

### **Unresectable and Resectable Hepatocellular Carcinoma**

For individuals who have unresectable hepatocellular carcinoma (HCC) confined to the liver and not associated with portal vein thrombosis who receive transcatheter arterial chemoembolization (TACE), the evidence includes several randomized controlled trials

(RCTs), large observational studies, and systematic reviews. Relevant outcomes are overall survival (OS), disease-specific survival, quality of life, and treatment-related mortality and morbidity. Evidence from 1 RCT has suggested that survival with TACE is at least as good as with systemic chemotherapy. One systematic review has highlighted possible biases associated with RCTs that compared TACE with no therapy. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have resectable HCC who receive neoadjuvant or adjuvant TACE, the evidence includes several RCTs and systematic reviews. Relevant outcomes are OS, disease-specific survival, quality of life, and treatment-related mortality and morbidity. Studies have shown little to no difference in OS rates with neoadjuvant TACE compared with surgery alone. A meta-analysis found no significant improvements in survival or recurrence with preoperative TACE for resectable HCC. While both RCTs and the meta-analyses that evaluated TACE as adjuvant therapy to hepatic resection in HCC reported positive results, the quality of individual studies and the methodologic issues related to the meta-analyses preclude certainty when interpreting the results. Well-conducted multicentric trials from the U.S. or Europe representing relevant populations with adequate randomization procedures, blinded assessments, centralized oversight, and publication in peer-reviewed journals are required. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have resectable HCC who receive TACE plus radiofrequency ablation (RFA), the evidence includes a single RCT and a systematic review. Relevant outcomes are OS, disease-specific survival, quality of life, and treatment-related mortality and morbidity. The RCT failed to show the superiority in survival benefit with combination TACE plus RFA treatment compared with surgery for HCC lesions 3 cm or smaller. Further, an ad hoc subgroup analysis showed a significant benefit for surgery in recurrence and OS in patients with lesions larger than 3 cm. It cannot be determined from this trial whether TACE plus RFA is as effective as a surgical resection for these small tumors. The systematic review, which included mostly retrospective observational studies, did not find a survival benefit with TACE plus RFA over surgery alone. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have unresectable HCC who receive TACE plus RFA, the evidence includes multiple systematic reviews and RCTs. Relevant outcomes are OS, disease-specific survival, quality of life, and treatment-related mortality and morbidity. Multiple meta-analyses and RCTs have shown a consistent benefit in survival and RFS favoring combination TACE plus RFA over RFA alone. However, results of these meta-analyses are difficult to interpret because the pooled data included heterogeneous patient populations and, in a few cases, data from a study retracted due to questions about data veracity. A larger well-conducted RCT has reported a relative reduction in the hazard of death by 44% and a 14% difference in 4-year survival favoring combination therapy. The major limitations of this trial were its lack of a TACE-alone arm and the generalizability of its findings to patient populations that have unmet needs such as those with multiple lesions larger than 3 cm and Child-Pugh class B or C. Further, this single-center trial was conducted in China, and until these results have been reproduced in patient populations representative of pathophysiology and clinical stage more commonly found in the U.S. or Europe, the results may not be generalizable. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

### **Bridge to Liver Transplant**

For individuals who have a single hepatocellular tumor less than 5 cm or no more than 3 tumors each less than 3 cm in size, absence of extrahepatic disease or vascular invasion, and Child-Pugh class A or B seeking to prevent further tumor growth and to maintain candidacy for liver transplant who receive pretransplant TACE, the evidence includes multiple small prospective studies. Relevant outcomes are OS, disease-specific survival, quality of life, and treatment-related mortality and morbidity. There is a lack of comparative trials on various locoregional treatments as a bridge therapy for liver transplantation. Multiple small prospective studies have demonstrated that TACE can prevent dropouts from the transplant list. TACE has become an accepted method to prevent tumor growth and progression while patients are on the liver transplant waiting list. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

### **Unresectable Intrahepatic Cholangiocarcinoma**

For individuals who have unresectable Intrahepatic cholangiocarcinoma who receive TACE, the evidence includes several retrospective observational studies and systematic reviews. Relevant outcomes are OS, disease-specific survival, quality of life, and treatment-related mortality and morbidity. Randomized controlled trials evaluating the benefit of adding TACE to standard of care for patients with unresectable Intrahepatic cholangiocarcinoma are lacking. Results of retrospective studies have shown a survival benefit with TACE over standard of care; however, systematic reviews comparing TACE to other locoregional therapies are conflicting. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

### **Transcatheter Arterial Chemoembolization for Symptomatic Unresectable Neuroendocrine Tumors**

For individuals who have symptomatic metastatic neuroendocrine tumors despite systemic therapy and are not candidates for surgical resection who receive TACE, the evidence includes retrospective single-cohort studies. Relevant outcomes are OS, disease-specific survival, symptoms, quality of life, and treatment-related mortality and morbidity. There is a lack of evidence from RCTs supporting the use of TACE. Uncontrolled trials have suggested that TACE reduces symptoms and tumor burden and improves hormone profiles. Generally, the response rates are over 50% and include patients with massive hepatic tumor burden. While many studies have demonstrated symptom control, survival benefits are less clear. Despite the uncertain benefit on survival, the use of TACE to palliate the symptoms associated with hepatic neuroendocrine metastases can provide a clinically meaningful improvement in net health outcome. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

### **Liver-Dominant Metastatic Uveal Melanoma**

For individuals who have liver-dominant metastatic uveal melanoma who receive TACE, the evidence includes observational studies and reviews. Relevant outcomes are OS, disease-specific survival, quality of life, and treatment-related mortality and morbidity. There is a lack of evidence from RCTs assessing the use of TACE. Noncomparative prospective and retrospective studies have reported improvements in tumor response and survival compared with historical controls. Given the very limited treatment response from systemic therapy and the rarity of this condition, the existing evidence may support conclusions that TACE meaningfully improves outcomes for patients with hepatic metastases from uveal melanoma.

The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

### **Other Unresectable Hepatic Metastases**

For individuals who have unresectable hepatic metastases from any other types of primary tumors (eg, colorectal or breast cancer) who receive TACE, the evidence includes multiple RCTs, observational studies, and systematic reviews. Relevant outcomes are OS, disease-specific survival, quality of life, and treatment-related mortality and morbidity. Multiple RCTs and numerous nonrandomized studies have compared TACE with alternatives in patients who have colorectal cancer and metastases to the liver. Nonrandomized studies have reported that TACE can stabilize disease in 40% to 60% of treated patients but whether this translates into a prolonged survival benefit relative to systemic chemotherapy alone is uncertain. Two small RCTs have reported that TACE with drug-eluting beads has resulted in statistically significant improvements in response rate and progression-free survival (PFS). Whether this translates into a prolonged survival benefit relative to systemic chemotherapy alone is uncertain. For cancers other than colorectal, the evidence is extremely limited and no conclusions can be made. Studies have assessed small numbers of patients and the results have varied due to differences in patient selection criteria and treatment regimens used. The evidence is insufficient to determine outcomes that the technology results in an improvement in the net health outcome.

## **SUPPLEMENTAL INFORMATION**

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

### **Clinical Input Received From Physician Specialty Societies and Academic Medical Centers**

In response to requests from the Blue Cross Blue Shield Association, input was received from 1 specialty medical society (2 reviewers) and 3 academic medical centers while their policy was under review in 2012. There was general agreement among reviewers that use of TACE was medically necessary for indications in the policy; however, reviewers were split for its use as a bridge to transplant. There was general support for the investigational policy statement for the use of TACE as neoadjuvant or adjuvant therapy in resectable hepatocellular carcinoma. Reviewers were split over the investigational policy statement to treat other liver metastases or for recurrent hepatocellular carcinoma. Four reviewers provided input on the use of TACE in unresectable cholangiocarcinoma; 2 reviewers considered it investigational and 2 others considered it investigational but also medically necessary, the latter citing data showing a survival benefit of TACE compared with supportive therapy.

### **Practice Guidelines and Position Statements**

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

## **National Comprehensive Cancer Network Guidelines**

### ***Hepatocellular Carcinoma***

The National Comprehensive Cancer Network (NCCN) (v.2.2024) guidelines on hepatocellular carcinoma list TACE as an option for patients who are not candidates for surgically curative treatments or as a part of strategy to bridge patients for other curative therapies.<sup>117</sup> Arterially directed therapies, including TACE, are appropriate for patients with unresectable or inoperable tumors that are not amenable to ablation therapy. Additionally, TACE in highly selected patients has been shown to be safe in the presence of limited tumor invasion of the portal vein.

The American Association for the Study of Liver Diseases 2023 guideline on hepatocellular carcinoma state that patients with Barcelona Clinic Liver Cancer Stage B HCC should receive TACE (Level 1, strong recommendation).<sup>118</sup> Both conventional TACE and drug-eluting bead TACE are mentioned, with no preference noted between these 2 modalities. The guideline also suggests using neoadjuvant locoregional therapies (which may include TACE) for bridging to liver transplant in patients with T2 lesions, in order to prevent disease progression and prevent dropouts from the waiting list. The guidelines recommend the use of locoregional therapies, including TACE, in patients with cirrhosis and T2 or T3 disease that is not amenable to resection or transplantation. The American Society of Clinical Oncology (ASCO) 2024 guideline on advanced HCC states that patients with locally advanced disease may be candidates for liver-directed therapies (including TACE); however, the guideline is focused on systemic therapy so there are no recommendations regarding TACE.<sup>119</sup>

### ***Intrahepatic Cholangiocarcinoma***

The NCCN (v.2.2024) guidelines on intrahepatic cholangiocarcinoma consider arterially directed therapies, including TACE, to be treatment options for unresectable and metastatic intrahepatic cholangiocarcinoma.<sup>117</sup>

### ***Neuroendocrine and Adrenal Tumors***

The NCCN (v.2.2022) guidelines on neuroendocrine and adrenal tumors recommend hepatic regional therapy, including arterial embolization, chemoembolization, or radioembolization, for unresectable liver metastases (category 2B).<sup>120</sup>

### ***Uveal Cancer***

The NCCN (v.1.2024) guidelines on uveal melanoma state that in patients with disease that is confined to the liver, regional liver-directed therapies such as chemoembolization, radioembolization, or immunoembolization should be considered.<sup>121</sup>

### ***Colon Cancer***

The NCCN (v.4.2024) guidelines on colon cancer recommend TACE only for clinical trials.<sup>122</sup> The (ASCO-) (2020) resource-stratified guidelines on late-stage colorectal cancer state that patients with unresectable liver metastases may receive TACE (weak recommendation).<sup>123</sup> However, this recommendation should only be implemented in centers with expertise in the technique, after multidisciplinary review, or in the context of a clinical trial. The 2022 guidelines for metastatic colorectal cancer from ASCO do not address TACE.<sup>124</sup>

### ***Breast Cancer***

The NCCN (v.4.2023) guidelines on breast cancer do not address TACE as a treatment option for breast cancer metastatic to the liver.<sup>125</sup>

### **U.S. Preventive Services Task Force Recommendations**

Not applicable.

## Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 18.

**Table 18. Summary of Key Trials**

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT06353126	A Prospective, Single Arm, Exploratory Study of Using Drug-eluting Beads Transarterial Chemoembolization Prior to SALT Liver Transplantation in the Treatment of Hepatocellular Carcinoma	40	Jul 2027
NCT03960008 <sup>a</sup>	A Randomized Multi-Center Phase III Study of Individualized Stereotactic Body Radiation Therapy (SBRT) Versus Trans-Arterial Chemoembolization (TACE) as a Bridge to Transplant in Hepatocellular Carcinoma	196	Mar 2024
NCT04143191	Sorafenib Plus Transarterial Chemoembolization Versus Sorafenib Alone as Postoperative Adjuvant Treatment for Resectable Primary Advanced Hepatocellular Carcinoma: A Phase 3, Multicenter, Randomized Controlled Trial	158	Sep 2023 (unknown status)
NCT04912258	Trans-arterial Chemoembolization With Irinotecan Drug-eluting Beads Before Liver Surgery for Patients With Primary Unresectable Colorectal Liver Metastasis: A Randomized Control Trial	80	Jun 2023 (unknown status)
NCT02724540 <sup>a</sup>	Randomized Embolization Trial for NeuroEndocrine Tumor Metastases To The Liver	162	Sept 2024
<i>Unpublished</i>			
NCT02936388	A Randomized Phase II Trial of Transarterial Radioembolisation With Yttrium-90 (SIRT) in Comparison to Transarterial Chemoembolisation With Cisplatin (TACE) in Patients With Liver Metastases From Uveal Melanoma	108	Dec 2022 (unknown status)

NCT: national clinical trial

<sup>a</sup> Denotes an industry sponsored or cosponsored clinical trial

## Government Regulations

### National/Local:

There is no specific national or local coverage determination for transcatheter arterial chemoembolization for hepatic tumors.

National Coverage Determination (NCD) for Therapeutic Embolization 20.28  
Effective date 12/15/78

Therapeutic embolization is covered when done for hemorrhage, and for other conditions amenable to treatment by the procedure, when reasonable and necessary for the individual patient. Renal embolization for the treatment of renal adenocarcinoma continues to be covered, effective December 15, 1978, as one type of therapeutic embolization to:

- Reduce tumor vascularity preoperatively
- Reduce tumor bulk in inoperable cases
- Palliate specific symptoms

*(The above Medicare information is current as of the review date for this policy. However, the coverage issues and policies maintained by the Centers for Medicare & Medicare Services [CMS, formerly HCFA] are updated and/or revised periodically. Therefore, the most current CMS information may not be contained in this document. For the most current information, the reader should contact an official Medicare source.)*

---

## Related Policies

Radioembolization for Primary and Metastatic Tumors of the Liver  
Radiofrequency Ablation of Primary or Metastatic Liver Tumors

---

## References

1. Marelli L, Stigliano R, Triantos C, et al. Transarterial therapy for hepatocellular carcinoma: which technique is more effective? A systematic review of cohort and randomized studies. *Cardiovasc Intervent Radiol*. Jan-Feb 2007;30(1):6-25. PMID 17103105
2. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Transcatheter arterial chemoembolization of hepatic tumors. TEC Assessments 2000; Volume 15, Tab 22.
3. National Cancer Institute, Surveillance Epidemiology and End Results Program. Cancer Stat Facts: Liver and Intrahepatic Bile Duct Cancer. n.d.; <https://seer.cancer.gov/statfacts/html/livibd.html> Accessed 8/15/24.
4. Qi X, Wang D, Su C, et al. Hepatic resection versus transarterial chemoembolization for the initial treatment of hepatocellular carcinoma: A systematic review and meta-analysis. *Oncotarget*. Jul 30 2015;6(21):18715-18733. PMID 26243835
5. Tian X, Dai Y, Wang DQ, et al. Transarterial chemoembolization versus hepatic resection in hepatocellular carcinoma treatment: a meta-analysis. *Drug Des Devel Ther*. 2015;9:4431-4440. PMID 26309396
6. Oliveri RS, Wetterslev J, Gluud C. Transarterial (chemo) embolisation for unresectable hepatocellular carcinoma. *Cochrane Database Syst Rev*. Mar 16 2011; (3):CD004787. PMID 21412886
7. Xie F, Zang J, Guo X et al. Comparison of transcatheter arterial chemoembolization and microsphere embolization for treatment of unresectable hepatocellular carcinoma: a meta-analysis. *J Cancer Res Clin Oncol*. Mar 2012; 138(3):455-62. PMID 22179199
8. Ahmad J, Rhee J, Carr BI. The effects of hepatic artery chemotherapy on viral hepatitis in patients with hepatocellular carcinoma. *Dig Dis Sci*. Feb 2005; 50(2): 331-5. PMID 15745096
9. Akamatsu M, Yoshida H, Obi S, et al. Evaluation of transcatheter arterial embolization prior to percutaneous tumor ablation in patients with hepatocellular carcinoma: a randomized controlled trial. *Liver Int*. Dec 2004; 24(6): 625-9. PMID 15566514
10. Bruix J, Llovet JM, Castells A, et al. Transarterial embolization versus symptomatic treatment in patients with advanced hepatocellular carcinoma: results of a randomized, controlled trial in a single institution. *Hepatology*. Jun 1998; 27(6): 1578-83. PMID 9620330
11. Cao GW, Hu S, Li G, et al. The clinical and experimental research of transhepatic arterial injection of 32P-glass microsphere therapy for hepatic carcinoma. *J Med Imaging*. 2005;15(8):678681.
12. Cao XC, Wang X, Tan J, et al. Clinical research of intra-arterial radioembolization with 32P-glass microspheres combined with chemoembolization for treatment of liver cancer. *Chin J Radiol*. 2005;39(10):10681072.

13. Carr BI, Kondragunta V, Buch SC, et al. Therapeutic equivalence in survival for hepatic arterial chemoembolization and yttrium 90 microsphere treatments in unresectable hepatocellular carcinoma: a two-cohort study. *Cancer*. Mar 01 2010; 116(5): 1305-14. PMID 20066715
14. Cheng SQ, Wu MC, Chen H, et al. [Transcatheter hepatic arterial chemoembolization and thymosin alpha1 in postoperative treatment of hepatocellular carcinoma]. *Zhonghua Zhong Liu Za Zhi*. May 2004; 26(5): 305-7. PMID 15312371
15. Doffoel M, Bonnetain F, Bouche O, et al. Multicentre randomised phase III trial comparing Tamoxifen alone or with Transarterial Lipiodol Chemoembolisation for unresectable hepatocellular carcinoma in cirrhotic patients (Federation Francophone de Cancerologie Digestive 9402). *Eur J Cancer*. Mar 2008; 44(4): 528-38. PMID 18242076
16. Du W, Lin S, Luo K, et al. Clinical analysis of TACE plus 32P-GMS in advanced hepatic carcinoma. *J Hepatobilia Surg*. 2002;10(5):351352.
17. Groupe d'Etude et de Traitement du Carcinome Hepatocellulaire. A comparison of lipiodol chemoembolization and conservative treatment for unresectable hepatocellular carcinoma. *N Engl J Med*. May 11 1995; 332(19): 1256-61. PMID 7708069
18. Hao N, Xiao X, Han X, et al. Efficacy of intra-arterial chemoembolization using drug microspheres in compare with chemoembolization in the treatment of primary hepatic carcinoma. *Tumor (Shanghai)*. 2000;20(5):375378.
19. Hou P, Guan G, Zhang X, Lu H, Wang S. Effects of intra-advanced 32P glass microspheres for advanced hepatic carcinoma. *Academic Journal of Fujian Medical University*. 2006;40(1):4850.
20. Kirchhoff TD, Rudolph KL, Layer G, et al. Chemoocclusion vs chemoperfusion for treatment of advanced hepatocellular carcinoma: a randomised trial. *Eur J Surg Oncol*. Mar 2006; 32(2): 201-7. PMID 16373084
21. Kooby DA, Egnatashvili V, Srinivasan S, et al. Comparison of yttrium-90 radioembolization and transcatheter arterial chemoembolization for the treatment of unresectable hepatocellular carcinoma. *J Vasc Interv Radiol*. Feb 2010; 21(2): 224-30. PMID 20022765
22. Lee W, Luo J, Yan Z, et al. Hepatic radioembolization with epirubicin mixed microsphere for the treatment of hepatocellular carcinoma. *J Nantong Univ (Medical Sciences)*. 2008;28(4):268270.
23. Lewandowski RJ, Kulik LM, Riaz A, et al. A comparative analysis of transarterial downstaging for hepatocellular carcinoma: chemoembolization versus radioembolization. *Am J Transplant*. Aug 2009; 9(8): 1920-8. PMID 19552767
24. Li JQ, Zhang YQ, Zhang WZ, et al. Randomized study of chemoembolization as an adjuvant therapy for primary liver carcinoma after hepatectomy. *J Cancer Res Clin Oncol*. 1995; 121(6): 364-6. PMID 7541051
25. Li Q, Wang J, Sun Y, et al. Postoperative transhepatic arterial chemoembolization and portal vein chemotherapy for patients with hepatocellular carcinoma: a randomized study with 131 cases. *Dig Surg*. 2006; 23(4): 235-40. PMID 16943671
26. Liu T, Zu M. Treatment of primary hepatic carcinoma by hepatic arterial chemoembolization with KMG microspheres and chemotherapeutic agents. *Acad Med Xuzhou*. 2005;25(2):126129.
27. Llovet JM, Real MI, Montana X, et al. Arterial embolisation or chemoembolisation versus symptomatic treatment in patients with unresectable hepatocellular carcinoma: a randomised controlled trial. *Lancet*. May 18 2002; 359(9319): 1734-9. PMID 12049862
28. Lo CM, Ngan H, Tso WK, et al. Randomized controlled trial of transarterial lipiodol chemoembolization for unresectable hepatocellular carcinoma. *Hepatology*. May 2002; 35(5): 1164-71. PMID 11981766

29. Pelletier G, Roche A, Ink O, et al. A randomized trial of hepatic arterial chemoembolization in patients with unresectable hepatocellular carcinoma. *J Hepatol.* Sep 1990; 11(2): 181-4. PMID 2174933
30. Pelletier G, Ducreux M, Gay F, et al. Treatment of unresectable hepatocellular carcinoma with lipiodol chemoembolization: a multicenter randomized trial. Groupe CHC. *J Hepatol.* Jul 1998; 29(1): 129-34. PMID 9696501
31. Salem R, Lewandowski RJ, Kulik L, et al. Radioembolization results in longer time-to-progression and reduced toxicity compared with chemoembolization in patients with hepatocellular carcinoma. *Gastroenterology.* Feb 2011; 140(2): 497-507.e2. PMID 21044630
32. Xiao E, Li D, Shen S, et al. Effect of preoperative transcatheter arterial chemoembolization on apoptosis of hepatocellular carcinoma cells. *Chin Med J.* Feb 2003; 116(2): 203-7. PMID 12775230
33. Bush DA, Smith JC, Slater JD, et al. Randomized clinical trial comparing proton beam radiation therapy with transarterial chemoembolization for hepatocellular carcinoma: results of an interim analysis. *Int J Radiat Oncol Biol Phys.* May 1 2016;95(1):477-482. PMID 27084661
34. Mabed M, Esmaeel M, El-Khodary T, et al. A randomized controlled trial of transcatheter arterial chemoembolization with lipiodol, doxorubicin and cisplatin versus intravenous doxorubicin for patients with unresectable hepatocellular carcinoma. *Eur J Cancer Care.* Sep 2009;18(5):492-499. PMID 19453695
35. Shen PC, Chang WC, Lo CH, et al. Comparison of Stereotactic Body Radiation Therapy and Transarterial Chemoembolization for Unresectable Medium-Sized Hepatocellular Carcinoma. *Int J Radiat Oncol Biol Phys.* Oct 01 2019; 105(2): 307-318. PMID 31175903
36. Biederman DM, Titano JJ, Korff RA, et al. Radiation segmentectomy versus selective chemoembolization in the treatment of early-stage hepatocellular carcinoma. *J Vasc Interv Radiol.* Jan 2018;29(1):30-37 e32. PMID 29169782
37. Molinari M, Kachura JR, Dixon E, et al. Transarterial chemoembolisation for advanced hepatocellular carcinoma: results from a North American cancer centre. *Clin Oncol (R Coll Radiol).* Nov 2006;18(9):684-692. PMID 17100154
38. Takayasu K, Arai S, Ikai I et al. Prospective cohort study of transarterial chemoembolization for unresectable hepatocellular carcinoma in 8510 patients. *Gastroenterology* Aug 2006; 131(2):461-9. PMID 16890600
39. Biselli M, Andreone P, Gramenzi A et al. Transcatheter arterial chemoembolization therapy for patients with hepatocellular carcinoma: a case-controlled study. *Clin Gastroenterol Hepatol* Sep 2005; 3(9):918-25. PMID 16234031
40. Chan KS, Tay WX, Cheo FY, et al. Preoperative transarterial chemoembolization (TACE) + liver resection versus upfront liver resection for large hepatocellular carcinoma (≥5 cm): a systematic review and meta-analysis. *Acta Chir Belg.* Dec 2023; 123(6): 601-617. PMID 37681991
41. Si T, Chen Y, Ma D, et al. Preoperative transarterial chemoembolization for resectable hepatocellular carcinoma in Asia area: a meta-analysis of random controlled trials. *Scand J Gastroenterol.* Dec 2016;51(12):1512-1519. PMID 27598831
42. Zhou Y, Zhang X, Wu L et al. Meta-analysis: preoperative transcatheter arterial chemoembolization does not improve prognosis of patients with resectable hepatocellular carcinoma. *BMC Gastroenterol* Mar 19 2013; 13:51. PMID 23509884
43. Chua TC, Liauw W, Saxena A et al. Systematic review of neoadjuvant transarterial chemoembolization for resectable hepatocellular carcinoma. *Liver Int.* Feb 2010; 30(2):166-74. PMID 19912531

44. Kaibori M, Tanigawa N, Kariya S et al. A prospective randomized controlled trial of preoperative whole-liver chemolipiodolization for hepatocellular carcinoma. *Dig Dis Sci*. May 2012; 57(5):1404-12. PMID 22271410
45. Zhou WP, Lai EC, Li AJ et al. A prospective, randomized, controlled trial of preoperative transarterial chemoembolization for resectable large hepatocellular carcinoma. *Ann Surg*. Feb 2009; 249(2):195-202. PMID 19212170
46. Cui H, Gao QQ, Li YY, et al. Influence of preventive effects of transcatheter arterial chemoembolization on primary hepatocellular carcinoma. *J Med Forum*. 2003;24:13.
47. Yamasaki S, Hasegawa H, Kinoshita H, et al. A prospective randomized trial of the preventive effect of preoperative transcatheter arterial embolization against recurrence of hepatocellular carcinoma. *Jpn J Cancer Res*. Feb 1996; 87(2): 206-11. PMID 8609071
48. Wu CC, Ho YZ, Ho WL, et al. Preoperative transcatheter arterial chemoembolization for resectable large hepatocellular carcinoma: a reappraisal. *Br J Surg*. Jan 1995; 82(1): 122-6. PMID 7881929
49. Yeh ML, Huang CI, Huang CF, et al. Neoadjuvant transcatheter arterial chemoembolization does not provide survival benefit compared to curative therapy alone in single hepatocellular carcinoma. *Kaohsiung J Med Sci*. Feb 2015;31(2):77-82. PMID 25645985
50. Choi GH, Kim DH, Kang CM et al. Is preoperative transarterial chemoembolization needed for a resectable hepatocellular carcinoma? *World J Surg*. Dec 2007; 31(12):2370-7. PMID 17912587
51. Liang L, Li C, Diao YK, et al. Survival benefits from adjuvant transcatheter arterial chemoembolization in patients undergoing liver resection for hepatocellular carcinoma: a systematic review and meta-analysis. *Therap Adv Gastroenterol*. 2020; 13: 1756284820977693. PMID 33329759
52. Liao M, Zhu Z, Wang H, et al. Adjuvant transarterial chemoembolization for patients after curative resection of hepatocellular carcinoma: a meta-analysis. *Scand J Gastroenterol*. Jun 2017;52(6-7):624-634. PMID 28276833
53. Li Q, Wang J, Sun Y et al. Efficacy of postoperative transarterial chemoembolization and portal vein chemotherapy for patients with hepatocellular carcinoma complicated by portal vein tumor thrombosis—a randomized study. *World J Surg*. Nov 2006; 30(11):2004-11. PMID 17058027
54. Zhong C, Guo RP, Li JQ, et al. A randomized controlled trial of hepatectomy with adjuvant transcatheter arterial chemoembolization versus hepatectomy alone for Stage III A hepatocellular carcinoma. *J Cancer Res Clin Oncol*. Oct 2009;135(10):1437-1445. PMID 19408012
55. Peng BG, He Q, Li JP, et al. Adjuvant transcatheter arterial chemoembolization improves efficacy of hepatectomy for patients with hepatocellular carcinoma and portal vein tumor thrombus. *Am J Surg*. Sep 2009;198(3):313-318. PMID 19285298
56. Gui CH, Baey S, D'cruz RT, et al. Trans-arterial chemoembolization + radiofrequency ablation versus surgical resection in hepatocellular carcinoma - A meta-analysis. *Eur J Surg Oncol*. May 2020; 46(5): 763-771. PMID 31937433
57. Liu H, Wang ZG, Fu SY, et al. Randomized clinical trial of chemoembolization plus radiofrequency ablation versus partial hepatectomy for hepatocellular carcinoma within the Milan criteria. *Br J Surg*. Mar 2016;103(4):348-356. PMID 26780107
58. Ako S, Nakamura S, Nouso K, et al. Transcatheter arterial chemoembolization to reduce size of hepatocellular carcinoma before radiofrequency ablation. *Acta Med Okayama*. Feb 2018;72(1):47-52. PMID 29463938
59. Haochen W, Jian W, Li S, et al. Transarterial chemoembolization plus multi-imaging-guided radiofrequency ablation for elimination of hepatocellular carcinoma nodules

- measuring 3.1 to 5.0 cm: a single-center study. *J Int Med Res.* Jul 2018;46(7): 2650-2657. PMID 29683022
60. Bholee AK, Peng K, Zhou Z, et al. Radiofrequency ablation combined with transarterial chemoembolization versus hepatectomy for patients with hepatocellular carcinoma within Milan criteria: a retrospective case-control study. *Clin Transl Oncol.* Jul 2017;19(7):844-852. PMID 28070766
  61. Lan T, Chang L, Rahmathullah MN, et al. Comparative efficacy of interventional therapies for early-stage hepatocellular carcinoma: a PRISMA-compliant systematic review and network meta-analysis. *Medicine (Baltimore).* Apr 2016;95(15):e3185. PMID 27082558
  62. Li L, Tian J, Liu P, et al. Transarterial chemoembolization combination therapy vs monotherapy in unresectable hepatocellular carcinoma: a meta-analysis. *Tumori.* Jun 02 2016;2016(3):301-310. PMID 27002950
  63. Lu Z, Wen F, Guo Q, et al. Radiofrequency ablation plus chemoembolization versus radiofrequency ablation alone for hepatocellular carcinoma: a meta-analysis of randomized-controlled trials. *Eur J Gastroenterol Hepatol.* Feb 2013;25(2):187-194. PMID 23134976
  64. Wang X, Hu Y, Ren M, et al. Efficacy and safety of radiofrequency ablation combined with transcatheter arterial chemoembolization for hepatocellular carcinomas compared with radiofrequency ablation alone: a time-to-event meta-analysis. *Korean J Radiol.* Jan-Feb 2016;17(1):93-102. PMID 26798221
  65. Peng ZW, Zhang YJ, Liang HH, et al. Recurrent hepatocellular carcinoma treated with sequential transcatheter arterial chemoembolization and RF ablation versus RF ablation alone: a prospective randomized trial. *Radiology.* Feb 2012;262(2):689-700. PMID 22157201
  66. Morimoto M, Numata K, Kondou M, et al. Midterm outcomes in patients with intermediate-sized hepatocellular carcinoma: a randomized controlled trial for determining the efficacy of radiofrequency ablation combined with transcatheter arterial chemoembolization. *Cancer.* Dec 01 2010;116(23):5452-5460. PMID 20672352
  67. Shibata T, Isoda H, Hirokawa Y, et al. Small hepatocellular carcinoma: is radiofrequency ablation combined with transcatheter arterial chemoembolization more effective than radiofrequency ablation alone for treatment? *Radiology.* Sep 2009;252(3):905-913. PMID 19567647
  68. Cheng BQ, Jia CQ, Liu CT, et al. Chemoembolization combined with radiofrequency ablation for patients with hepatocellular carcinoma larger than 3 cm: a randomized controlled trial. *JAMA.* Apr 09 2008;299(14):1669-1677. PMID 18398079
  69. DeAngelis CD, Fontanarosa PB. Retraction: Cheng B-Q, et al. Chemoembolization combined with radiofrequency ablation for patients with hepatocellular carcinoma larger than 3 cm: a randomized controlled trial. *JAMA.* 2008;299(14):1669-1677. *JAMA.* May 13 2009;301(18):1931. PMID 19380477
  70. Yi Y, Zhang Y, Wei Q, et al. Radiofrequency ablation or microwave ablation combined with transcatheter arterial chemoembolization in treatment of hepatocellular carcinoma by comparing with radiofrequency ablation alone. *Chin J Cancer Res.* Feb 2014;26(1):112-118. PMID 24653633
  71. Peng ZW, Zhang YJ, Chen MS, et al. Radiofrequency ablation with or without transcatheter arterial chemoembolization in the treatment of hepatocellular carcinoma: a prospective randomized trial. *J Clin Oncol.* Feb 01 2013;31(4):426-432. PMID 23269991
  72. Martin AP, Bartels M, Hauss J, et al. Overview of the MELD score and the UNOS adult liver allocation system. *Transplant Proc.* Dec 2007; 39(10): 3169-74. PMID 18089345
  73. Organ Procurement and Transplantation Network (OPTN). OPTN Policies. 2021;

74. Mazzaferro V, Regalia E, Doci R, et al. Liver transplantation for the treatment of small hepatocellular carcinomas in patients with cirrhosis. *N Engl J Med*. Mar 14 1996;334(11): 693-699. PMID 8594428
75. Pomfret EA, Washburn K, Wald C, et al. Report of a national conference on liver allocation in patients with hepatocellular carcinoma in the United States. *Liver Transpl*. Mar 2010;16(3):262-278. PMID 20209641
76. Butcher DA, Brandis KJ, Wang H, et al. Long-term survival and postoperative complications of pre-liver transplantation transarterial chemoembolisation in hepatocellular carcinoma: A systematic review and meta-analysis. *Eur J Surg Oncol*. Mar 2022; 48(3): 621-631. PMID 34774394
77. Si T, Chen Y, Ma D, et al. Transarterial chemoembolization prior to liver transplantation for patients with hepatocellular carcinoma: A meta-analysis. *J Gastroenterol Hepatol*. Jul 2017;32(7):1286-1294. PMID 28085213
78. Graziadei IW, Sandmueller H, Waldenberger P, et al. Chemoembolization followed by liver transplantation for hepatocellular carcinoma impedes tumor progression while on the waiting list and leads to excellent outcome. *Liver Transpl*. Jun 2003;9(6):557-563. PMID 12783395
79. Maddala YK, Stadheim L, Andrews JC, et al. Drop-out rates of patients with hepatocellular cancer listed for liver transplantation: outcome with chemoembolization. *Liver Transpl*. Mar 2004;10(3):449-455. PMID 15004776
80. Obed A, Beham A, Pullmann K et al. Patients without hepatocellular carcinoma progression after transarterial chemoembolization benefit from liver transplantation. *World J Gastroenterol* 2007; 13(5):761-7.
81. Yao FY. Liver transplantation for hepatocellular carcinoma: beyond the Milan criteria. *Am J Transplant*. Oct 2008; 8(10):1982-9. PMID 18727702
82. Gabr A, Abouchaleh N, Ali R, et al. Comparative study of post-transplant outcomes in hepatocellular carcinoma patients treated with chemoembolization or radioembolization. *Eur J Radiol*. Aug 2017;93:100-106. PMID 28668402
83. Park SY, Kim JH, Yoon HJ, et al. Transarterial chemoembolization versus supportive therapy in the palliative treatment of unresectable intrahepatic cholangiocarcinoma. *Clin Radiol*. Apr 2011;66(4):322-328. PMID 21356394
84. Seidensticker R, Seidensticker M, Doegen K, et al. Extensive Use of Interventional Therapies Improves Survival in Unresectable or Recurrent Intrahepatic Cholangiocarcinoma. *Gastroenterol Res Pract*. 2016; 2016: 8732521. PMID 26966431
85. Lv TR, Hu HJ, Liu F, et al. The effect of trans arterial chemoembolization in the management of intrahepatic cholangiocarcinoma. A systematic review and meta-analysis. *Eur J Surg Oncol*. May 2022; 48(5): 956-966. PMID 35065841
86. Edeline J, Lamarca A, McNamara MG, et al. Locoregional therapies in patients with intrahepatic cholangiocarcinoma: A systematic review and pooled analysis. *Cancer Treat Rev*. Sep 2021; 99: 102258. PMID 34252720
87. Mosconi C, Solaini L, Vara G, et al. Transarterial Chemoembolization and Radioembolization for Unresectable Intrahepatic Cholangiocarcinoma-a Systemic Review and Meta-Analysis. *Cardiovasc Intervent Radiol*. May 2021; 44(5): 728-738. PMID 33709272
88. Boehm LM, Jayakrishnan TT, Miura JT, et al. Comparative effectiveness of hepatic artery based therapies for unresectable intrahepatic cholangiocarcinoma. *J Surg Oncol*. Feb 2015;111(2):213-20. PMID 25176325

89. Knuppel M, Kubicka S, Vogel A et al. Combination of conservative and interventional therapy strategies for intra- and extrahepatic cholangiocellular carcinoma: a retrospective survival analysis. *Gastroenterol Res Pract*. 2012; 2012:190708. PMID 21776251
90. Tai E, Kennedy S, Farrell A, et al. Comparison of transarterial bland and chemoembolization for neuroendocrine tumours: a systematic review and meta-analysis. *Curr Oncol*. Dec 2020; 27(6): e537-e546. PMID 33380868
91. Nazario J, Gupta S. Transarterial liver-directed therapies of neuroendocrine hepatic metastases. *Semin Oncol*. Apr 2010; 37(2):118-26. PMID 20494704
92. Ruutiainen AT, Soulen MC, Tuite CM et al. Chemoembolization and bland embolization of neuroendocrine tumor metastases to the liver. *J Vasc Interv Radiol*. Jul 2007; 18(7):847-55. PMID 17609443
93. Gupta S, Yao JC, Ahrar KHae et al. Hepatic artery embolization and chemoembolization for treatment of patients with metastatic carcinoid tumors: the M.D. Anderson experience. *Cancer J*. Jul-Aug 2003; 9(4):241-7. PMID 12967136
94. Osborne DA, Zervos EE, Strosberg J et al. Improved outcome with cytoreduction versus embolization for symptomatic hepatic metastases of carcinoid and neuroendocrine tumors. *Ann Surg Oncol*. Apr 2006; 13(4):572-81. PMID 16511671
95. Rowcroft A, Loveday BPT, Thomson BNJ, et al. Systematic review of liver directed therapy for uveal melanoma hepatic metastases. *HPB (Oxford)*. Apr 2020; 22(4): 497-505. PMID 31791894
96. Huppert PE, Fierlbeck G, Pereira P et al. Transarterial chemoembolization of liver metastases in patients with uveal melanoma. *Eur J Radiol*. Jun 2010; 74(3):e38-44. PMID 19467811
97. Sharma KV, Gould JE, Harbour JW et al. Hepatic arterial chemoembolization for management of metastatic melanoma. *AJR Am J Roentgenol*. Jan 2008; 190(1):99-104. PMID 18094299
98. Bedikian AY, Legha SS, Mavligit G, Toummttlarot MD et al. Treatment of uveal melanoma metastatic to the liver: a review of the M. D. Anderson Cancer Center experience and prognostic factors. *Cancer*. Nov 01 1995; 76(9):1665-70. PMID 8635073
99. Patel K, Sullivan K, Berd D et al. Chemoembolization of the hepatic artery with BCNU for metastatic uveal melanoma: results of a phase II study. *Melanoma Res*. Aug 2005; 15(4):297-304. PMID 16034309
100. Sugumar K, Stitzel H, Wu V, et al. Outcomes of Hepatic Artery-Based Therapies and Systemic Multiagent Chemotherapy in Unresectable Colorectal Liver Metastases: A Systematic Review and Meta-analysis. *Ann Surg Oncol*. Jul 2024; 31(7): 4413-4426. PMID 38502296
101. Zacharias AJ, Jayakrishnan TT, Rajeev R, et al. Comparative effectiveness of hepatic artery based therapies for unresectable colorectal liver metastases: a meta-analysis. *PLoS One*. 2015;10(10):e0139940. PMID 26448327
102. Richardson AJ, Laurence JM, Lam VW. Transarterial chemoembolization with irinotecan beads in the treatment of colorectal liver metastases: systematic review. *J Vasc Interv Radiol*. Aug 2013;24(8):1209-1217. PMID 23885916
103. Swierz MJ, Storman D, Riemsma RP, et al. Transarterial (chemo)embolisation versus no intervention or placebo for liver metastases. *Cochrane Database Syst Rev*. Mar 12 2020; 3: CD009498. PMID 32163181
104. Hunt TM, Flowerdew AD, Birch SJ, et al. Prospective randomized controlled trial of hepatic arterial embolization or infusion chemotherapy with 5-fluorouracil and degradable starch microspheres for colorectal liver metastases. *Br J Surg*. Jul 1990; 77(7): 779-82. PMID 2200559

105. Eichler K, Zangos S, Mack MG, et al. First human study in treatment of unresectable liver metastases from colorectal cancer with irinotecan-loaded beads (DEBIRI). *Int J Oncol.* Oct 2012; 41(4): 1213-20. PMID 22842404
106. Martin RC, Scoggins CR, Tomalty D, et al. Irinotecan drug-eluting beads in the treatment of chemo-naïve unresectable colorectal liver metastasis with concomitant systemic fluorouracil and oxaliplatin: results of pharmacokinetics and phase I trial. *J Gastrointest Surg.* Aug 2012; 16(8): 1531-8. PMID 22528576
107. Vogl TJ, Jost A, Nour-Eldin NA, et al. Repeated transarterial chemoembolisation using different chemotherapeutic drug combinations followed by MR-guided laser-induced thermotherapy in patients with liver metastases of colorectal carcinoma. *Br J Cancer.* Mar 27 2012; 106(7): 1274-9. PMID 22382689
108. Martin RC, Joshi J, Robbins K, et al. Hepatic intra-arterial injection of drug-eluting bead, irinotecan (DEBIRI) in unresectable colorectal liver metastases refractory to systemic chemotherapy: results of multi-institutional study. *Ann Surg Oncol.* Jan 2011; 18(1): 192-8. PMID 20740319
109. Aliberti C, Fiorentini G, Muzzio PC, et al. Trans-arterial chemoembolization of metastatic colorectal carcinoma to the liver adopting DC Bead(R), drug-eluting bead loaded with irinotecan: results of a phase II clinical study. *Anticancer Res.* Dec 2011; 31(12): 4581-7. PMID 22199334
110. Fiorentini G, Aliberti C, Tilli M, et al. Intra-arterial infusion of irinotecan-loaded drug-eluting beads (DEBIRI) versus intravenous therapy (FOLFIRI) for hepatic metastases from colorectal cancer: final results of a phase III study. *Anticancer Res.* Apr 2012;32(4):1387-1395. PMID 22493375
111. Martin RC, Scoggins CR, Schreeder M, et al. Randomized controlled trial of irinotecan drug-eluting beads with simultaneous FOLFOX and bevacizumab for patients with unresectable colorectal liver-limited metastasis. *Cancer.* Oct 15 2015;121(20):3649-3658. PMID 26149602
112. Vogl TJ, Gruber T, Balzer JO, et al. Repeated transarterial chemoembolization in the treatment of liver metastases of colorectal cancer: prospective study. *Radiology.* Jan 2009;250(1):281-289. PMID 19092099
113. Vogl TJ, Mack MG, Balzer JO, et al. Liver metastases: neoadjuvant downsizing with transarterial chemoembolization before laser-induced thermotherapy. *Radiology.* Nov 2003;229(2):457-464. PMID 14500854
114. Hong K, McBride JD, Georgiades CS et al. Salvage therapy for liver-dominant colorectal metastatic adenocarcinoma: comparison between transcatheter arterial chemoembolization versus yttrium-90 radioembolization. *J Vasc Interv Radiol.* Mar 2009; 20(3):360-7. PMID 19167245
115. Rivera K, Jeyarajah DR, Washington K. Hepatectomy, RFA, and Other Liver Directed Therapies for Treatment of Breast Cancer Liver Metastasis: A Systematic Review. *Front Oncol.* 2021; 11: 643383. PMID 33842354
116. Vogl TJ, Naguib NN, Nour-Eldin NE, et al. Transarterial chemoembolization (TACE) with mitomycin C and gemcitabine for liver metastases in breast cancer. *Eur Radiol.* Jan 2010;20(1):173-180. PMID 19657653
117. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines): Hepatobiliary Cancers, Version 2.2024. Updated July 2, 2024.  
[https://www.nccn.org/professionals/physician\\_gls/pdf/hepatobiliary.pdf](https://www.nccn.org/professionals/physician_gls/pdf/hepatobiliary.pdf) Accessed 8/15/24.

118. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines): Neuroendocrine and Adrenal Tumors, Version 2.2024. Updated 8/1/24.  
[https://www.nccn.org/professionals/physician\\_gls/pdf/neuroendocrine.pdf](https://www.nccn.org/professionals/physician_gls/pdf/neuroendocrine.pdf) Accessed 8/15/24.
119. Singal AG, Llovet JM, Yarrow M, et al. AASLD Practice Guidance on prevention, diagnosis, and treatment of hepatocellular carcinoma. Hepatology. Dec 01 2023; 78(6): 1922-1965. PMID 37199193
120. Gordan JD, Kennedy EB, Abou-Alfa GK, et al. Systemic Therapy for Advanced Hepatocellular Carcinoma: ASCO Guideline Update. J Clin Oncol. May 20 2024; 42(15): 1830-1850. PMID 38502889
121. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines): Melanoma: Uveal, Version 1.2024. Updated 5/23/24.  
[https://www.nccn.org/professionals/physician\\_gls/pdf/uveal.pdf](https://www.nccn.org/professionals/physician_gls/pdf/uveal.pdf) Accessed 8/15/24.
122. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines): Colon Cancer, Version 4.2024. Updated 7/3/24.  
[https://www.nccn.org/professionals/physician\\_gls/pdf/colon.pdf](https://www.nccn.org/professionals/physician_gls/pdf/colon.pdf) Accessed 8/15/24.
123. Chiorean EG, Nandakumar G, Fadelu T, et al. Treatment of Patients With Late-Stage Colorectal Cancer: ASCO Resource-Stratified Guideline. JCO Glob Oncol. Mar 2020; 6: 414-438. PMID 32150483
124. Morris VK, Kennedy EB, Baxter NN, et al. Treatment of Metastatic Colorectal Cancer: ASCO Guideline. J Clin Oncol. Jan 20 2023; 41(3): 678-700. PMID 36252154
125. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines): Breast Cancer, Version 4.2024. Updated 7/3/24.  
[https://www.nccn.org/professionals/physician\\_gls/pdf/breast.pdf](https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf) Accessed 8/15/24.

*The articles reviewed in this research include those obtained in an Internet based literature search for relevant medical references through 8/15/24, the date the research was completed.*

### Joint BCBSM/BCN Medical Policy History

<b>Policy Effective Date</b>	<b>BCBSM Signature Date</b>	<b>BCN Signature Date</b>	<b>Comments</b>
7/16/04	7/16/04	7/2/04	Joint policy established
11/15/05	11/15/05	11/11/05	Routine maintenance
3/1/07	12/28/06	01/29/07	Routine maintenance
1/1/09	10/13/08	10/13/08	Routine maintenance
1/1/10	10/13/09	11/02/09	Routine maintenance
5/1/12	2/21/12	2/21/12	Routine maintenance; policy updated to mirror BCBSA policy
7/1/13	4/16/13	4/22/13	Routine maintenance Added treatment of unresectable cholangiocarcinoma as an exclusion. Updated references and rationale.
7/1/14	4/10/14	5/1/14	Routine maintenance Deleted CPT codes 37204 and 75894, new CPT code 37243 added
5/1/16	2/16/16	2/16/16	Routine maintenance
5/1/17	2/21/17	2/21/17	Routine maintenance
1/1/18	10/19/17	10/19/17	Routine maintenance Added "and liver function not characterized as Child-Pugh class C" to first bullet under inclusions Rationale and references updated
1/1/19	10/16/18	10/16/18	Routine maintenance
1/1/20	10/15/19		Routine maintenance
1/1/21	10/20/20		Routine maintenance References added: 8-32, 35, 45, 46, 47, 54, 70, 81, 88, 95-101, 109, 113 Added inclusion of TACE plus RFA
1/1/22	10/19/21		Routine maintenance Ref 50,85,109 added
1/1/23	10/18/22		Routine maintenance (ls) Ref 75 added
1/1/24	10/17/23		Routine maintenance (jf) Vendor Managed: NA

1/1/25	10/15/24		Routine maintenance (jf) Vendor Managed: NA Ref added: 40,85,86,87,100,118,119 and 124

Next Review Date: 4<sup>th</sup> Qtr, 2025

**BLUE CARE NETWORK BENEFIT COVERAGE**  
**POLICY: TRANSCATHETER ARTERIAL CHEMOEMBOLIZATION OF HEPATIC TUMORS**  
**(TACE)**

**I. Coverage Determination:**

<b>Commercial HMO (includes Self-Funded groups unless otherwise specified)</b>	Covered; refer to policy criteria.
<b>BCNA (Medicare Advantage)</b>	See Government Regulations section.
<b>BCN65 (Medicare Complementary)</b>	Coinsurance covered if primary Medicare covers the service.

**II. Administrative Guidelines:**

- The member's contract must be active at the time the service is rendered.
- Coverage is based on each member's certificate and is not guaranteed. Please consult the individual member's certificate for details. Additional information regarding coverage or benefits may also be obtained through customer or provider inquiry services at BCN.
- The service must be authorized by the member's PCP except for Self-Referral Option (SRO) members seeking Tier 2 coverage.
- Services must be performed by a BCN-contracted provider, if available, except for Self-Referral Option (SRO) members seeking Tier 2 coverage.
- Payment is based on BCN payment rules, individual certificate and certificate riders.
- Appropriate copayments will apply. Refer to certificate and applicable riders for detailed information.
- CPT - HCPCS codes are used for descriptive purposes only and are not a guarantee of coverage.